



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Jon Yager

UB Person Number:

5	0	2	3	6	2	3	5
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1	2	3	4	5	6	7	TOTAL	GRADE

14

10

7

20

20

6

6

2

10

94

A

1

2

3

4

5

6

7

PIAZZA

HILL

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$V = \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \xrightarrow{-2} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & 4+b \end{array} \right] \xrightarrow{\begin{smallmatrix} + \\ - \end{smallmatrix}} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 6+b \end{array} \right] \begin{array}{l} x_1 = -3x_3 \\ x_2 = 2 - x_3 \\ x_3 = x_3 \end{array}$$

$$b = -6$$



$$b) \left[\begin{array}{ccc|c} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 2 & -3 & 0 & 0 \end{array} \right] = \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{l} x_1 = -3x_3 \\ x_2 = -2x_3 \\ x_3 = x_3 \end{array}$$

$$\boxed{\text{Yes linear independent}} \xrightarrow{x_3=1} -3 \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} - 2 \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix} + \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$



This shows that these vectors are linearly dependent



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$\begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \cdot \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} \quad \checkmark$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix} \cdot \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} ?$$

$$C = B \cdot (A^T)^{-1} \quad (A^T)^{-1} = B$$

$$C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 3 & 2 \\ -1 & 2 & 3 \end{bmatrix}$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

$$T = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix} \quad \checkmark$$

$$b) \quad \left[\begin{array}{cc|c} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 1 & 9 \\ 0 & -1 & -3 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & 0 & 7 \\ 0 & 1 & 9 \\ 0 & 0 & 0 \end{array} \right]$$

$$u = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$x_1 = 7 \\ x_2 = 9$$

$$u = \begin{bmatrix} 7 \\ 9 \end{bmatrix} \quad \checkmark$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -2 \\ 0 & 2 & 4 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -2 \\ 0 & 2 & 4 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - \text{pivot in every column}$$

$T_A(v) = Av$ is one-to-one

T_A is not one-to-one

$x_3 = \text{free}$

$$x_1 = 2x_3$$

$$x_2 = -2x_3$$

$$v_1 = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}$$

$$v_2 = \begin{bmatrix} -2 \\ 2 \\ -1 \end{bmatrix}$$

$$x_3 = 1$$

$$x_3 = -1$$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

$$u = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad v = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad w = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

~~False~~

$$w + u = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

but there is no λ value where $\lambda w = \text{Span}(u, v)$

$$w = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} = -u + v \quad \text{so } w \in \text{Span}(u, v)$$

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

✓ True linear independent means vectors are not multiples
so removing one will not affect linear independency



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

False

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix} \quad \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

Au and Av are dependent
 $\begin{bmatrix} 1 \\ 2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ are independent

✓

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

True ✓ a transform will only move a vector, not change it so if u is in the span it will also be in the transform span why?

$$T = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \perp \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

?