



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

David Flores

UB Person Number:

5	0	2	3	6	3	8	7
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1 2 3 4 5 6 7 TOTAL GRADE

--	--	--	--	--	--	--	--	--

19

10

10

20

20

2

2

--

--

83

B+

1

2

3

4

5

6

7

PIAZZA

HILL

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

a) $(-2) \rightarrow \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right]$

$(-1) \rightarrow \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & b+4 \end{array} \right]$

$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & b+2 \end{array} \right] \quad b+6$

$b = -2$ or else no solution
-6

b) Set $\{v_1, v_2, v_3\}$ is linearly dependent because if $b = -2$, and
Since there is no leading one (pivot position) in third column,
then there is a free variable x_3 meaning infinite solutions \Rightarrow linearly dependent



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$A = \begin{array}{l} (-1) \\ + \end{array} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \downarrow$$

$$\begin{array}{l} (-2) \\ + \end{array} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \downarrow$$

$$+ \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \downarrow$$

$$\begin{array}{l} + \\ (-2) \end{array} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \downarrow$$

$$+ \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & -3 & 4 & -2 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \downarrow$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \rightarrow A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} \checkmark$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix}$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ -1 & 0 & 2 & 4 & 5 & 4 \\ 2 & 1 & -1 & 3 & 2 & 1 \end{array} \right]$$

$$\left(\begin{array}{l} R_2 + R_1 \\ R_3 - 2R_1 \end{array} \right) \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 2 & 1 & -1 & 3 & 2 & 1 \end{array} \right]$$

$$\left(\begin{array}{l} R_3 - R_2 \end{array} \right) \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 0 & -1 & -1 & 1 & -2 & -5 \end{array} \right]$$

$$\left(\begin{array}{l} R_3 + R_2 \end{array} \right) \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 0 & 0 & 1 & 6 & 5 & 2 \end{array} \right]$$

$$\left(\begin{array}{l} R_1 - R_2 \\ R_1 - 3R_3 \end{array} \right) \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 0 & 0 & 1 & 6 & 5 & 2 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 8 & 5 & 0 \\ 0 & 1 & 0 & -7 & -3 & 3 \\ 0 & 0 & 1 & 6 & 5 & 2 \end{array} \right]$$

$$\rightarrow C = \begin{bmatrix} 8 & 5 & 0 \\ -7 & -3 & 3 \\ 6 & 5 & 2 \end{bmatrix} \quad \checkmark$$

Simpler:

$$C = (A^T)^{-1} \cdot B$$

$$= (A^{-1})^T \cdot B$$

Then use A^{-1} from problem 2,



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

$$a) T(e_1) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 - 0 \\ 1 + 0 \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$T(e_2) = T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 - 2 \\ 0 + 1 \\ 0 - 3 \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$$

Standard matrix A :

$$A = [T(e_1) \ T(e_2)]$$

$$A = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix} \checkmark$$

$$b) \begin{array}{c} u_1 \quad u_2 \\ (-1) \rightarrow \begin{bmatrix} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{bmatrix} \end{array}$$

$$(-1) \rightarrow \begin{bmatrix} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 1 & -3 & -2 \end{bmatrix}$$

$$(-1) \rightarrow \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 1 & -3 & -2 \end{bmatrix}$$

$$+ \rightarrow \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & -1 & -3 \end{bmatrix}$$

$$(2) \rightarrow \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$u = \begin{bmatrix} 7 \\ 3 \end{bmatrix} \checkmark$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix} (v_2)$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} (v_2)$

a) $\begin{pmatrix} (-2) \end{pmatrix} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 3 & 4 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
 \uparrow
 one-to-one ✓

b) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 3 & 4 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$
 not one-to-one ✓

b) $T_A(v_1) = T_A(v_2)$

$v_1 = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \quad v_2 = \begin{bmatrix} 4 \\ -4 \\ 2 \end{bmatrix}$

Try
 $v_1 = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}$

$v_2 = \begin{bmatrix} 4 \\ -4 \\ 2 \end{bmatrix}$

$Av_1 = Av_2$

$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \cdot \begin{bmatrix} 4 \\ -4 \\ 2 \end{bmatrix}$

$\begin{bmatrix} 2 - 2 + 0 \\ 0 - 4 + 4 \\ 6 - 8 + 2 \end{bmatrix} = \begin{bmatrix} 4 - 4 + 0 \\ 0 - 8 + 8 \\ 12 - 16 + 4 \end{bmatrix}$

$\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \checkmark$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

True, [✓] if a vector is in span of other vectors when modified, then that vector is in span itself. | ?

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

True, [✓] in \mathbb{R}^3 if $\{u, v, w\}$ has one particular solution, then the vectors are linearly independent with one another always. | ?



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

~~True~~, if Au, Av are linearly dependent, then
 u, v are always zero vectors, which are linearly dependent.
 ?

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

True ✓, linear transformation is in span of other
 trans. if the vector of lin. trans is in span of
 other vectors. | why?