



MTH 309T LINEAR ALGEBRA EXAM 1

October 3, 2019

Name:

Alexander T. Nowasell

UB Person Number:

5	0	2	9	0	3	2	6
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1	2	3	4	5	6	7	TOTAL	GRADE

0

--

--

--

--

--

--

0

nan

1

2

3

4

5

6

7

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 2 & -3 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 1 & -2 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 2 & -3 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & -4 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & -1 & -2 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & b-2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & b \end{bmatrix}$$

for $\begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$ in span



~~linearly independent~~

~~linearly independent~~

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced matrix

b) No

linearly dependent

$$\begin{bmatrix} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

~~$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$~~

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$

Check:

$$\begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \cdot \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{bmatrix}$$

Reduction

$$\begin{bmatrix} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 1 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 1 \\ 0 & 0 & 1 & 2 & -2 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 0 & -3 & 4 & 2 \\ 0 & 1 & 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 2 & -2 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & -2 & 3 & 2 \\ 0 & 1 & 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 2 & -2 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & 1 \\ 0 & 1 & -1 & -1 \\ 0 & 0 & 1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 0 & -3 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} -2 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 0 & -2 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & 0 \\ 1 & 0 & 1 & 1 \\ 0 & 2 & -1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & 0 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & -2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 0 & 4 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -2 \end{bmatrix}$$



12
34
56

135
246

3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

~~Matrix~~ ~~Matrix~~

$$A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix}$$

$$A^{T^{-1}} \cdot B = C$$

$$A^{T^{-1}} = \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix}$$

$$\begin{aligned} -2+4+6 &= -7 \\ 3-4-6 &= -7 \\ -1+4+3 &= 6 \end{aligned}$$

~~Matrix~~

$$\begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

=

$$C = \begin{bmatrix} 8 & 5 & 0 \\ -7 & -3 & 3 \\ 6 & 5 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 & 1 \\ -1 & 0 & 2 & 0 \\ 2 & 1 & -1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 2 & 1 \\ 0 & -1 & -1 & -2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 3 \end{bmatrix}$$

$$-4+5+4=5$$

$$6-5-4=-3$$

$$-2+5+2=5$$

$$-6+4+2=0$$

$$9-4-2=3$$

$$-3+4+1=2$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.



b) $x_1 = 7$
 $x_2 = 3$

$\begin{bmatrix} 7 \\ 3 \end{bmatrix}$



a) $\begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$



$$\begin{bmatrix} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & 1 & -3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & -1 & -3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & -1 & -3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & -3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & -18 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 1 \end{bmatrix}$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 4 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

1 0 0 | Pivot
0 1 0 | Pivot
0 0 1 | Pivot

one to one

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$$

0 0 0 | Not Pivot

Not one-to-one

$$v_1 = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}$$

$$v_2 = \begin{bmatrix} 4 \\ -4 \\ 2 \end{bmatrix}$$

$$\begin{aligned} x_1 &= 2x_3 \\ x_2 &= -2x_3 \end{aligned}$$

$$\begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix} \quad \begin{matrix} u & v & w \end{matrix}$$

$$w + u \in \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

True,
if double plus
u kept it the
Span of u, v ,
the w by itself
should be in the
Span.

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

True,
because if, $\{u, v, w\}$
was linearly independent, then
making w zero didn't change it
to be dependent, so removing
 w entirely changes nothing.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

True The
Transformation
the
the $\times 2$
Matrix keep
the li

They are
dependent after
the transformation,
they put in

True

Putting to linearly dependant vectors through transformation, they must stay dependant when put together with more dependants.

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^2$ are vectors such that \mathbf{u} is in $\text{Span}(\mathbf{v}, \mathbf{w})$ then $T(\mathbf{u})$ must be in $\text{Span}(T(\mathbf{v}), T(\mathbf{w}))$.

True, the ^{linear} transformation must be one to one and onto, so the span would be kept intact.