



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Daniel Kane

UB Person Number:

5	0	2	3	8	7	0	6
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1 2 3 4 5 6 7 TOTAL GRADE

--	--	--	--	--	--	--	--	--

14

10

10

20

14

10

6

2

--

86

B+

1

2

3

4

5

6

7

PIAZZA

HILL

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

a) ~~$v_1 + v_3 = w \mid b = 2$~~

~~$-v_1 + v_3 = w$~~

$2v_2 = \begin{bmatrix} -2 \\ 2 \\ -6 \end{bmatrix}, b = -6$

$b = -6$ ✓

$-3v_1 + v_3 = \begin{bmatrix} -2 \\ 2 \\ -6 \end{bmatrix}, b = -6$

b) The set $\{v_1, v_2, v_3\}$ ~~is linearly independent~~, because the vectors cannot linearly combine to cancel one another
~~cannot cancel all terms~~

$\begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$

cannot cancel all terms

$\begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}$

no scalar multiple

They can. E.g. $3v_1 + 2v_2 - v_3 = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -1 & 1 & -1 & 0 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & -1 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & -1 & 1 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & -1 & -2 & 2 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$[A|I] \rightarrow [I|A^{-1}]$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} \quad \checkmark$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix}$$

$$A^T C = B$$

$$C = (A^T)^{-1} B \quad (A^T)^{-1} = (A^{-1})^T = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}^T = \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix} \quad \checkmark$$

$$C = \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} = \begin{bmatrix} (-2+4+6) & (-4+5+4) & (-6+4+2) \\ (3-4-6) & (6-5-4) & (9-4-2) \\ (-1+4+3) & (-2+5+2) & (-3+4+1) \end{bmatrix}$$

$$C = \begin{bmatrix} 8 & 5 & 0 \\ -7 & -3 & 3 \\ 6 & 5 & 2 \end{bmatrix} \quad \checkmark$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

a) $T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$
 $T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$ $T = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$ ✓

b) $1 = x_1 - 2x_2$
 $10 = x_1 + x_2$
 $-2 = x_1 - 3x_2$

$$\begin{bmatrix} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 1 & 3 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 1 & -3 & -2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & -3 & -9 \end{bmatrix}$$

$-7 \begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix} \begin{matrix} x_1 = 7 \\ x_2 = 3 \end{matrix}$ $u = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$ ✓



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

a) One to one \rightarrow pivot in every column

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 4 \\ 0 & 1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ A is one-to-one } \checkmark$$

b) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ not one-to-one \checkmark

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 + x_2 \\ 2x_2 + 4x_3 \\ 3x_1 + 4x_2 + 2x_3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ 0 & 6 \\ 3 & 6 \end{bmatrix} \begin{matrix} x_1 = 1 \\ x_2 = 1 \end{matrix} \begin{bmatrix} 2 \\ 6 \\ 9 \end{bmatrix}$$

$$\begin{matrix} x_1 = 2 \\ x_2 = 1 \end{matrix} \begin{bmatrix} 3 \\ 6 \\ 12 \end{bmatrix}$$

No such vectors exist, since $\begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ 4 \\ 2 \end{bmatrix}$ are linearly independent

Since T_A is not one-to-one these vectors must exist.

Try e.g. $v_1 = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}, v_2 = \begin{bmatrix} 4 \\ -4 \\ 2 \end{bmatrix}$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

True

For any equation $w + u = a_1 u + a_2 v$, $w = a_1 u - u + a_2 v = (a_1 - 1)u + a_2 v$
 $w \in \text{Span}(u, v)$



b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

True

If $\{u, v\}$ were dependent, then $a_1 u + a_2 v + 0w = 0$ would prove $\{u, v, w\}$ to be dependent





7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

False

$$A = \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \quad v_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad v_2 = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

$$Av_1 = \begin{bmatrix} -1 \\ 0 \end{bmatrix} \quad Av_2 = \begin{bmatrix} -3 \\ 0 \end{bmatrix}$$

v_1, v_2 are independent

Av_1, Av_2 are dependent



b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

$$u = a_1 v + a_2 w$$

True

~~Their transformations are the same~~

$u, a_1 v + a_2 w$
are equivalent
vectors

← actually equal
(by your assumption)

∴
Their transformations
are the same

| so?

