



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Justin W. Allen

UB Person Number:

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 5 | 0 | 2 | 9 | 4 | 9 | 4 | 8 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 |
| 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 |

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1 2 3 4 5 6 7 TOTAL GRADE

| | | | | | | | | |
|--|--|--|--|--|--|--|--|--|
| | | | | | | | | |
|--|--|--|--|--|--|--|--|--|

--

--

--

--

--

--

--

0

nan

1

2

3

4

5

6

7

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$\begin{array}{ccc|c} v_1 & v_2 & v_3 & \\ \hline 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \xrightarrow{R_3 = R_3 - 2R_1} \begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & (4+b) \end{array}$$

$$R_1 = R_1 + R_2 \quad \begin{bmatrix} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & (4+b) \end{bmatrix} \quad R_3 = R_3 + R_2 \quad \begin{bmatrix} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & (6+b) \end{bmatrix}$$

a. $w \in \text{Span}(v_1, v_2, v_3)$ when $b = -6$

$$b. \quad \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 2 & -3 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -1 \\ & 1 & -1 \end{bmatrix}$$

The set is not linearly independent, because the v_3 column is a free variable, so there is not a leading one in each column



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 = R_2 - R_1}$$

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right]$$

$$R_1 = R_1 + R_2$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_3 = R_3 - 2R_2}$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$R_2 = R_2 + R_3$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$R_1 = R_1 - R_3$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

$$A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix}$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ -1 & 0 & 2 & 0 & 1 & 0 \\ 2 & 1 & -1 & 0 & 0 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 1 & 1 & 0 \\ 2 & 1 & -1 & 0 & 0 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 1 & 1 & 0 \\ 0 & -1 & -1 & -2 & 0 & 1 \end{array} \right]$$

$$\begin{array}{ccc|ccc} 1 & 0 & -2 & 0 & -1 & 0 \\ 0 & 1 & 2 & 1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array}$$

$$\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 1 & 2 \\ 0 & 1 & 0 & 3 & -1 & -2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array}$$

$$(A^T)^{-1} = \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix}$$

$$(A^T)^{-1} A^T C = (A^T)^{-1} B$$

$$\begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

$$\begin{array}{ccc} (-2 + 4 + 6) & (-4 + 5 + 4) & (-6 + 4 + 2) \\ (3 + -4 - 6) & (6 - 5 - 4) & (9 - 4 - 2) \\ (-1 + 4 + 3) & (2 + 5 + 2) & (-3 + 4 + 1) \end{array}$$

$$\begin{bmatrix} 8 & 5 & 0 \\ -7 & -3 & 3 \\ 6 & 9 & 2 \end{bmatrix}$$

$$C = \begin{bmatrix} 8 & 5 & 0 \\ -7 & -3 & 3 \\ 6 & 9 & 2 \end{bmatrix}$$



$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

$$a. \quad T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$$

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$$

$$T(e_1) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$T(e_2) = \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 1 & -3 & -2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & -1 & -3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & -1 & -3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$x_1 = 7$$

$$x_2 = 3$$

$$u = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

$$\begin{array}{ccc} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{array} \xrightarrow{R_3 = R_3 - 3R_1}$$

$$\begin{array}{ccc} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 4 \end{array}$$

$$\begin{array}{ccc} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 4 \end{array}$$

$$\begin{array}{ccc} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 1 & 4 \end{array}$$

$$\begin{array}{ccc} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 2 \end{array}$$

$$\begin{array}{ccc} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{array}$$

A is one-to-one, because there is a leading one in each column.

b. $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

$$\begin{array}{ccc} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \end{array}$$

$$\begin{array}{ccc} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{array}$$

The matrix is not one-to-one, because the third column does not have a leading one.

$$T_A(v_1) = T_A(v_2) \quad T_A(v_1 - v_2) = 0$$

$$\begin{array}{cccc} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array}$$

$$\begin{array}{cccc} 1 & 0 & -2 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array}$$

$$\begin{array}{cccc} 1 & 1 & 0 & 0 \\ 0 & 2 & 4 & 0 \\ 3 & 4 & 2 & 0 \end{array}$$

$$\begin{array}{cccc} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 1 & 2 & 0 \end{array}$$

$$x_1 - 2x_3 = 0$$

$$x_1 = 2x_3$$

$$x_2 + 2x_3 = 0$$

$$x_2 = -2x_3$$

$$x_3 = x_3$$

$$x_3 = x_3$$

$x_3 \text{ Nul } A \in \text{Span} \left(\begin{bmatrix} 2 \\ -2 \\ 0 \end{bmatrix} \right)$

$$v_1 = \begin{bmatrix} 2 \\ -2 \\ 0 \end{bmatrix} \quad v_2 = \begin{bmatrix} -2 \\ 2 \\ 0 \end{bmatrix}$$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

This statement is false, because $w + u$ will only span (u, v) if $w + u$ is in the null set of (u, v)

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

This statement is true.

$$u = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad v = \begin{bmatrix} 4 \\ 1 \\ 0 \end{bmatrix} \quad w = \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix}$$

If $\{u, v, w\}$ is linearly independent then each column in the aug matrix must have a leading one. If you remove one vector (column), each column will still have a leading one



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

$$\begin{bmatrix} 2 & 4 \\ 6 & 8 \end{bmatrix}$$

This statement is false

$$\left| \begin{array}{ccc} 1 & 1 & 2 \\ 0 & 2 & 4 \end{array} \right|$$

$$\left| \begin{array}{ccc} 1 & 1 & 2 \\ 0 & 2 & 4 \end{array} \right|$$

$$\left| \begin{array}{ccc} 1 & 1 & 2 \\ 0 & 2 & 4 \end{array} \right|$$

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

$T(u)$ must be in $\text{Span}(T(v), T(w))$, be