4. Consider the matrix

$$A = \begin{bmatrix} 1 & 8 & 4 \\ -2 & 11 & 4 \\ 2 & -8 & -1 \end{bmatrix}$$

Knowing that eigenvalues of A are $\lambda_1 = 3$ and $\lambda_2 = 5$ diagonalize this matrix; that is, find a diagonal matrix D and an invertible matrix P such that

$$A = PDP^{-1}$$

Note: you do not need to compute P^{-1} .

$$Null(A-\int_{1}I) = \begin{bmatrix} -2 & 8 & 4 & 0 \\ -2 & 8 & 4 & 0 \\ 2 & -8 & -4 & 0 \end{bmatrix} \xrightarrow{-\frac{1}{2}} \begin{bmatrix} -2 & 8 & 4 & 0 \\ 0 & 0 & 0 & 6 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{=} \begin{bmatrix} 1 & -4 & -2 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} x_{1} = 4x_{2} + 2x_{3} \\ x_{2} \\ x_{3} \end{bmatrix}$$

$$X_{2} \begin{bmatrix} 4 \\ 1 \\ 0 \end{bmatrix} + X_{3} \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}$$

$$|V||(A-h_2I) = \begin{vmatrix} -2 & 6 & 4 & 0 \\ -2 & 6 & 4 & 0 \\ +2 & 8 & -6 & 0 \end{vmatrix} \xrightarrow{\chi_2} \begin{vmatrix} 1 & -2 & -1 & 0 \\ -2 & 6 & 4 & 6 \\ 2 & -8 & -6 & 0 \end{vmatrix} \rightarrow \begin{bmatrix} 1-2 & -1 & 0 \\ 0 & 2 & 2 \\ 0-4 & -4 & 0 \end{bmatrix} \xrightarrow{\chi_2} \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} x_1 = x_3 \\ x_2 = x_3 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_3 \\ -1 \\ 1 \end{bmatrix}$$

$$P = \begin{bmatrix} 4 & 2 & -1 \\ 1 & 0 & -1 \\ 0 & 1 & 1 \end{bmatrix}$$