



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

PRANSHU TEOTIA

UB Person Number:

5	0	2	1	8	5	6	0
0	●	0	0	0	0	0	●
1	1	1	●	1	1	1	1
2	2	●	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
●	5	5	5	5	●	5	5
6	6	6	6	6	6	●	6
7	7	7	7	7	7	7	7
8	8	8	8	●	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1	2	3	4	5	6	7	TOTAL	GRADE

--

10

6

9

--

--

--

25

nan

1

2

3

4

5

6

7

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

- a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.
 b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$a) \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \xrightarrow{R_3 = R_3 - 2R_1} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & 4+b \end{array} \right] \xrightarrow{R_3 = R_3 + R_2} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 6+b \end{array} \right]$$

$$R_1 = R_1 + R_2 \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 6+b \end{array} \right] \quad \underline{b = -6} \rightarrow \text{For all other values of } b \text{ there will not be a solution.}$$

$$\left[\begin{array}{ccc} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 2 & -3 & 0 \end{array} \right] \xrightarrow{R_3 = R_3 - 2R_1} \left[\begin{array}{ccc} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 0 & -1 & -2 \end{array} \right] \xrightarrow{R_3 = R_3 + R_2} \left[\begin{array}{ccc} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{array} \right]$$

$$R_1 = R_1 + R_2 \rightarrow \begin{array}{c} x_1 \quad x_2 \quad x_3 \\ \left[\begin{array}{ccc} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{array} \right] \\ \quad \quad \quad \uparrow \\ \quad \quad \quad \text{Free variable} \end{array}$$

~~Since~~ Since, x_3 is a free variable, the set $\{v_1, v_2, v_3\}$ is NOT linearly independent



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{\substack{R_2 = -R_1 + R_2}} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right]$$

$$\downarrow R_1 = R_1 + R_2$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -2 & -2 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xleftarrow{R_3 = 2R_2 + R_3}$$

$$\downarrow R_2 = R_2 + R_3$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \xrightarrow{R_1 = R_1 - R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A^T = \begin{bmatrix} 1 & 4 & 3 \\ 2 & 5 & 2 \\ 3 & 4 & 1 \end{bmatrix} \cdot \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 4 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{cases} a + 4d + 3g = 1 \\ 2a + 5d + 2g = 4 \\ 3a + 4d + g = 3 \end{cases} \quad \begin{cases} b + 4e + 3h = 2 \\ 2b + 5e + 2h = 5 \\ 3b + 4e + h = 4 \end{cases} \quad \begin{cases} c + 4f + 3i = 3 \\ 2c + 5f + 2i = 4 \\ 3c + 4f + i = 1 \end{cases}$$

~~$$\begin{cases} a = -1 \\ d = 2 \\ g = -2 \end{cases}$$~~

$$a = -1, d = 2, g = -2$$

$$b = -\frac{3}{4}, e = 2, h = \frac{7}{4} \quad c = -2, f = 2, i = -1$$

$$C = \begin{bmatrix} -1 & -\frac{3}{4} & -2 \\ 2 & 2 & 2 \\ -2 & -\frac{7}{4} & -1 \end{bmatrix}$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

$$\begin{aligned} x_1 - 2x_2 &= 1 \\ x_1 + x_2 &= 10 \\ x_1 - 3x_2 &= -2 \end{aligned} \quad \text{we get } x_1 = 7, x_2 = 3$$

(b) $u = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$

$$\begin{bmatrix} a & b \\ c & d \\ e & f \end{bmatrix} \begin{bmatrix} 7 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix} \rightarrow \begin{aligned} 7a + 3b &= 1 \\ 7c + 3d &= 10 \\ 7e + 3f &= -2 \end{aligned}$$

$3 \times 2 \quad \quad 2 \times 1$

Standard Matrix of $T = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$

(a)



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

a) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix} \xrightarrow{R_2 = R_2/2} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 3 & 4 & 4 \end{bmatrix} \xrightarrow{R_3 = R_3 - 3R_1} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 4 \end{bmatrix}$

$\xrightarrow{R_3 = R_3 - R_2} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 2 \end{bmatrix} \xrightarrow{\begin{matrix} R_2 = R_2 - R_3 \\ \text{and} \\ R_3 = R_3/2 \end{matrix}} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_1 = R_1 - R_2} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

We have pivot pos. in every row so matrix $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$ is one-to-one.

b) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \xrightarrow{R_2 = R_2/2} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 3 & 4 & 2 \end{bmatrix} \xrightarrow{R_3 = R_3 - 3R_1} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \end{bmatrix} \xrightarrow{R_3 = R_3 - R_2} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$

$\xrightarrow{R_1 = R_1 - R_2} \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$

Since every col is not a pivot col matrix

$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$ is not one-to-one.

$v_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad v_2 = \begin{bmatrix} -5 \\ 8 \\ 0 \end{bmatrix}$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

TRUE

If $w + u = v$, then $w = v - u$

v is in the $\text{span}(u, v)$ and $(v - u)$ is also in the $\text{span}(u, v)$

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

TRUE

The set of u, v, w , in that each column ~~is~~ is a pivot column

So, in the set of $\{u, v\}$ each column will also be a pivot column.



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

FALSE If $A = \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$ and $u = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ $v = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
 then $Au \rightarrow \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$ is linearly dependent ~~but u is linearly~~
~~independent.~~
 ~~$A = \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$~~ ~~TRUE~~ the but $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ is linearly
 independent.

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

TRUE

$$u = v + w$$

$$T(u) = T(v + w)$$

$$\hookrightarrow \underline{T(v) + T(w)}$$

So, $T(u)$ is in $\text{Span}(T(v), T(w))$