



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Shawn Mammen

UB Person Number:

5	0	1	9	0	2	2	7
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1 2 3 4 5 6 7 TOTAL GRADE

--	--	--	--	--	--	--	--	--

--

--

--

--

--

--

--

0

nan

1

2

3

4

5

6

7

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

a)

$$\begin{array}{c} \begin{array}{cccc} 2 & -2 & 2 & -4 \end{array} \\ \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 - 2R_1} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & b-4 \end{array} \right] \xrightarrow{\begin{array}{l} R_3 \rightarrow R_3 + R_2 \\ R_3 \rightarrow R_3 \end{array}} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & b-2 \end{array} \right]$$

when $b=2$

b) The set isn't linearly independent because since the span is made up of linear combinations of the vectors all the vectors in the span are linearly dependent.



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} \quad \begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 1 & 0 & 1 & | & 0 & 1 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & | & -2 & 3 & -1 \\ 0 & 1 & 0 & | & 1 & -1 & 1 \\ 0 & 0 & 1 & | & 2 & -2 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 1 & 0 & 1 & | & 0 & 1 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_2 \rightarrow R_2 - R_1} \begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 0 & 1 & -1 & | & -1 & 1 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 0 & 1 & -1 & | & -1 & 1 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_1 \rightarrow R_1 + R_2} \begin{bmatrix} 1 & 0 & 1 & | & 0 & 1 & 0 \\ 0 & 1 & -1 & | & -1 & 1 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_3 \rightarrow R_3 - 2R_2} \begin{bmatrix} 1 & 0 & 1 & | & 0 & 1 & 0 \\ 0 & 1 & -1 & | & -1 & 1 & 0 \\ 0 & 0 & 1 & | & 2 & -2 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 1 & | & -2 & 3 & -1 \\ 0 & 1 & 0 & | & 1 & -1 & 1 \\ 0 & 0 & 1 & | & 2 & -2 & 1 \end{bmatrix} \xrightarrow{\begin{matrix} R_2 \rightarrow R_2 + R_3 \\ R_1 \rightarrow R_1 - R_3 \end{matrix}} \begin{bmatrix} 1 & 0 & 1 & | & 0 & 1 & 0 \\ 0 & 1 & -1 & | & -1 & 1 & 0 \\ 0 & 0 & 1 & | & 2 & -2 & 1 \end{bmatrix}$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix}$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ -1 & 0 & 2 & 4 & 5 & 4 \\ 2 & 1 & -1 & 3 & 2 & 1 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + R_1} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 2 & 1 & -1 & 3 & 2 & 1 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 - 2R_1} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 0 & -1 & -1 & 1 & -2 & -5 \end{array} \right]$$

$$\downarrow R_3 \rightarrow R_3 + R_2$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 0 & 0 & 1 & 6 & 5 & 2 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 - 2R_3} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ 0 & 1 & 0 & -7 & -3 & 3 \\ 0 & 0 & 1 & 6 & 5 & 2 \end{array} \right] \xrightarrow{R_1 \rightarrow R_1 - R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 8 & 5 & 0 \\ 0 & 1 & 0 & -7 & -3 & 3 \\ 0 & 0 & 1 & 6 & 5 & 2 \end{array} \right]$$

$$C = \begin{bmatrix} 8 & 5 & 0 \\ -7 & -3 & 3 \\ 6 & 5 & 2 \end{bmatrix}$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors \mathbf{u} satisfying $T(\mathbf{u}) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix} \xrightarrow{R_3 \rightarrow R_3 - 3R_1} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 4 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \xrightarrow{\begin{matrix} R_3 - 3R_1 \\ R_2 \rightarrow \frac{1}{2}R_2 \\ R_3 \rightarrow R_3 - R_2 \end{matrix}} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 2 \end{bmatrix} \downarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

T_A not one-to-one



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

True, because if we add the vectors together in a vector equation and the resulting vector is in the span, then that means that the vector that was added is in the span as it is a linear combination.

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

true, because if one of the vectors is linearly dependent then the whole set would have been linearly dependent but it's not.



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

true

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

true, if all the vectors undergo a linear transformation then they will still share the relationship of being linear combinations.