



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Alvin Tsang

UB Person Number:

5	0	2	5	6	5	4	8
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1	2	3	4	5	6	7	TOTAL	GRADE

--

--

--

--

--

--

--

0

nan

1

2

3

4

5

6

7

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right]$$

$$\downarrow$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 1 & 1 & -2-b \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 3 & -4 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 2+\frac{b}{2} \end{array} \right]$$

$$2 + \frac{b}{2} = 0$$

$$\frac{b}{2} = -2$$

$$b = -4$$

The set $\{v_1, v_2, v_3\}$ are linearly independent because v_1, v_2, v_3 are not scalar multiples of each other.



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & -1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & -1 & -2 & 2 & -1 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$A^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \quad A^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$

$$(A^T)^{-1} = (A^{-1})^T$$

$$(A^{-1})^T C = B$$

$$(A^T)^{-1} A^T C = (A^T)^{-1} B$$

$$I = (A^{-1})^T B$$

$$= \begin{bmatrix} 0 & 1 & 0 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

$$C = \begin{bmatrix} 4 & 5 & 4 \\ 0 & -1 & 0 \\ -3 & -2 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} + \begin{bmatrix} 4 \\ -4 \\ -8 \end{bmatrix} + \begin{bmatrix} 0 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \\ -3 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 2 \\ 4 \end{bmatrix} + \begin{bmatrix} 5 \\ -5 \\ -10 \end{bmatrix} + \begin{bmatrix} 0 \\ 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ -1 \\ -4 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 4 \\ -4 \\ -8 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \\ -7 \end{bmatrix}$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

Standard matrix $T = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$

$$T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{array} \right]$$

↓

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & 1 & 3 \end{array} \right]$$

$$\left[\begin{array}{cc|c} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 3 & 9 \end{array} \right]$$

$$\left[\begin{array}{cc|c} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right]$$

$$\left[\begin{array}{cc|c} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right]$$

$$x_1 - 2x_2 = 1$$

$$x_1 + x_2 = 10$$

$$x_1 - 3x_2 = -2$$

$$-3x_2 = -9$$

$$x_2 = 3$$

$$x_1 + 3 = 10$$

$$x_1 = 7$$

$$u = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$$

↓

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 1 \end{bmatrix}$$

↓

$$\begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 1/2 \end{bmatrix}$$

↓

$$\begin{bmatrix} \textcircled{1} & 0 & 0 \\ 0 & \textcircled{1} & 0 \\ 0 & 0 & \textcircled{1} \end{bmatrix}$$

One-to-one because pivot position in every column

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} \textcircled{1} & 0 & -2 \\ 0 & \textcircled{1} & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

Not one-to-one:

$$T_A(v_1) = T_A(v_2)$$

$$T_A(v_1) - T_A(v_2) = 0$$

$$A(v_1 - v_2) = 0$$

$$A(v_1 - v_2) = 0$$

$$A(v_1 - v_2) = 0$$

$$\begin{bmatrix} -2 \\ 2 \\ 0 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

$$v_1 = \begin{bmatrix} -2 \\ 2 \\ 0 \end{bmatrix} \quad v_2 = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

$$w + u = c_1 u + c_2 v$$

$$w = c_1 u + c_2 v - u$$

True because if $w \in \text{Span}(u, v)$, according to the definition, the set of vectors is linearly dependent. All the vectors are just linear combinations of each other, so w is in the span of u, v .

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

True because if they are all linearly independent, they can't be changed to be linearly dependent. There is no relationship between u, v, w , so if you take one out, it would not change the linear independence.



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

False $u = \begin{bmatrix} 1 \\ 0 \end{bmatrix} v = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

Not linearly dependent

$$Au = \begin{bmatrix} 1+0 \\ 1+0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$Av = \begin{bmatrix} 0+1 \\ 0+1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Linearly dependent

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

True $c_1 v + c_2 w = u$

$$T(c_1 v + c_2 w) = T(u)$$

$$c_1 T(v) + c_2 T(w) = T(u)$$

$$A(c_1 v + c_2 w) = T(u)$$

$$T_A(c_1 v) + T_A(c_2 w) = T(u)$$

$$T(c_1 v + c_2 w) = T(u) \quad ; \quad c_1 v + c_2 w = u$$