



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Raymond Groesbeck

UB Person Number:

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 5 | 0 | 2 | 8 | 6 | 2 | 3 | 9 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 |
| 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 |

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1 2 3 4 5 6 7 TOTAL GRADE

| | | | | | | | | |
|--|--|--|--|--|--|--|--|--|
| | | | | | | | | |
|--|--|--|--|--|--|--|--|--|

16

8

7

9

8

10

7

65

nan

1

2

3

4

5

6

7

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2v_1 \\ 2v_2 \\ bv_3 \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$b) \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 2 & -3 & 0 \end{bmatrix} \begin{matrix} \leftarrow -2 \\ \leftarrow -2 \end{matrix}$$

$$\begin{bmatrix} -2 & -2 \\ 0 & 1 \\ -4 & -3 \end{bmatrix}$$

$$\downarrow$$

$$\begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 0 & -1 & 0 \end{bmatrix} \begin{matrix} \leftarrow +1 \\ \leftarrow +1 \end{matrix}$$

$$\begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 2 \end{bmatrix} \begin{matrix} \leftarrow +1/2 \\ \leftarrow +1/2 \end{matrix}$$

$$\begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{matrix} \leftarrow +1 \\ \leftarrow +1 \end{matrix}$$

$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ is linearly independent because each part of the matrix has a pivot point



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} \cdot -1 \\ \\ \end{array}$$

↓

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} \\ \cdot -2 \\ \end{array}$$

↓

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \begin{array}{l} \cdot -1 \\ \cdot -2 \\ \cdot 1 \end{array}$$

↓

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 0 & -3 & 4 & -2 \\ 0 & 1 & 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \cdot 1$$

↓

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 2 & -1 \\ 0 & 1 & 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$\boxed{\begin{bmatrix} -2 & 2 & -1 \\ 1 & -2 & 1 \\ 2 & -2 & 1 \end{bmatrix} = A^{-1}}$$

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix} \cdot \begin{bmatrix} C \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 & 1 & 0 & 0 \\ -1 & 0 & 2 & 0 & 1 & 0 \\ 2 & 1 & -1 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{\substack{R_2 + R_1 \\ R_3 - 2R_1}} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \xrightarrow{\substack{R_2 - 4R_1 \\ R_3 - 3R_1}} \begin{bmatrix} 1 & 2 & -6 \\ 13 & -13 & 13 \\ 12 & -12 & 6 \end{bmatrix} = C$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 1 & 1 & 0 \\ 0 & -1 & -1 & -2 & 0 & 1 \end{array} \right] \xrightarrow{R_3 + R_2}$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \xrightarrow{-2}$$

$$\begin{bmatrix} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 3 & -1 & -2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{bmatrix} \xrightarrow{-1}$$

$$A^{-1} = \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix}$$

$$C = \begin{bmatrix} 12 & -6 \\ 13 & 13 \\ 12 & 6 \end{bmatrix}$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

a) $\begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$

$$\begin{cases} x_1 = 2x_2 + 1 \\ x_2 = 3 \end{cases}$$

b) $\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{array} \right] \begin{matrix} \\ \cdot -1 \\ \cdot -1 \end{matrix}$

$$\begin{bmatrix} x_1 = 7 \\ x_2 = 3 \end{bmatrix}$$

$$\downarrow$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & -1 & -3 \end{array} \right] \cdot \frac{1}{3}$$

$$u = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$$

$$\downarrow$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & -1 & -3 \end{array} \right] \cdot 1$$

$$\downarrow$$

$$\begin{bmatrix} x_1 & x_2 \\ 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

a) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix} \xrightarrow{\cdot -3}$

\downarrow
 $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 4 \end{bmatrix} \xrightarrow{\cdot 1/2}$

\downarrow
 $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 4 \end{bmatrix} \xrightarrow{\cdot -1}$

\downarrow
 $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 2 \end{bmatrix} \xrightarrow{\cdot -1/2}$

\downarrow
 $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{\cdot -1}$

\downarrow
 $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

is one to one

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \xrightarrow{\cdot -3}$

\downarrow
 $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 2 \end{bmatrix} \xrightarrow{\cdot 1/2}$

\downarrow
 $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{\cdot -1/2}$

\downarrow
 $\begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$
 free

$x_1 = 2x_3$

$x_2 = -2x_3$

$x_3 = x_3$

1

3

Not one to one

b



Two vectors

$v_1 = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \quad v_2 = \begin{bmatrix} 6 \\ -6 \\ 3 \end{bmatrix}$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

False,

Counter example

$$u = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \quad v = \begin{bmatrix} 3 \\ 3 \\ 2 \end{bmatrix} \quad w = \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}$$

$$u + w = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ 2 \end{bmatrix} \in \text{Span}\left(\begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 2 \end{bmatrix}\right)$$

$$\begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix} \notin \text{Span}\left(\begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 2 \end{bmatrix}\right)$$

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

True because in order to be independent there must be no dependency on another vector, so if a vector were to leave the group that vector would still be independent.



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

False

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

False

$$v = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad w = \begin{bmatrix} 0 \\ 2 \end{bmatrix} \quad u = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad T = \begin{bmatrix} 5 & 7 \end{bmatrix}$$