



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Judy Mei

UB Person Number:

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 5 | 0 | 2 | 3 | 7 | 0 | 3 | 2 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 6 | 6 | 6 | 6 | 6 | 6 | 6 | 6 |
| 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 |

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1 2 3 4 5 6 7 TOTAL GRADE

| | | | | | | | | |
|--|--|--|--|--|--|--|--|--|
| | | | | | | | | |
|--|--|--|--|--|--|--|--|--|

10

7

1

10

0

5

2

--

--

35

F

1

2

3

4

5

6

7

PIAZZA

HILL

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 2 & -3 & 0 & 0 \end{array} \right] \xrightarrow{R_3 - 2R_1} \left[\begin{array}{ccc|c} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & -1 & -2 & 0 \end{array} \right] \xrightarrow{R_2 + R_1} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & -1 & -2 & 0 \end{array} \right] \xrightarrow{R_3 + R_2} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\xrightarrow{R_1 + (-R_2)} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \xrightarrow{\text{restore } R_3} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & -1 & -2 & 0 \end{array} \right] \xrightarrow{R_3 + R_2} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

b) It is linearly dependent since the reduced form contains a free variable meaning it has infinitely many solutions. ✓

a) ~~$b = 2$~~ ← where did this come from?
 $b = -6$



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & x_1 & y_1 & z_1 \\ 1 & 0 & 1 & x_2 & y_2 & z_2 \\ 0 & 2 & -1 & x_3 & y_3 & z_3 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{array} \right] \quad \begin{array}{l} -(+1) \\ 1-1=0 \\ 0+0=0 \end{array}$$

$$\begin{array}{l} x_1 - x_2 + 2x_3 \\ x_1 + 0x_2 + x_3 \\ 0x_1 + 2x_2 + 2x_3 \end{array} \left| \begin{array}{l} 1 \\ 0 \\ 0 \end{array} \right. \rightarrow \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 + (-R_1)} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \quad \begin{array}{l} 0 \\ 1-1=0 \\ 0+0=0 \end{array}$$

$$\xrightarrow{R_1 + R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_3 + (-2R_2)} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & -3 & 2 & -2 & 1 \end{array} \right] \xrightarrow{\times 1/3} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -2/3 & 2/3 & 1/3 \end{array} \right] \quad \begin{array}{l} 0 \\ 0 \\ -1+1=0 \end{array}$$

$$\xrightarrow{R_2 + R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -5/3 & 5/3 & 1/3 \\ 0 & 0 & 1 & -2/3 & 2/3 & 1/3 \end{array} \right] \xrightarrow{R_1 + (-R_3)} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 2/3 & -1/3 & -1/3 \\ 0 & 1 & 0 & -5/3 & 5/3 & 1/3 \\ 0 & 0 & 1 & -2/3 & 2/3 & 1/3 \end{array} \right] \quad \begin{array}{l} 0 \\ -5/3 \\ -2/3 \end{array}$$

This is the long way to solve this...

$$A^{-1} = \begin{bmatrix} 2/3 & -1/3 & -1/3 \\ -5/3 & 5/3 & 1/3 \\ -2/3 & 2/3 & 1/3 \end{bmatrix}$$

where did this come from?



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \quad Ax = B$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \rightarrow A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix} \quad \checkmark$$

|

?

.

|



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

a) $A = [T(e_1) \ T(e_2)]$ where $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

$T(e_1) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

$T(e_2) = T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$

$A = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$ ✓

b) ?



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(\mathbf{v}) = A\mathbf{v}$ is one-to-one or not. If T_A is not one-to-one, find two vectors \mathbf{v}_1 and \mathbf{v}_2 such that $T_A(\mathbf{v}_1) = T_A(\mathbf{v}_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

?



0

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

True. \leftarrow why?

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

True. In order to be linearly independent, each column is a pivot column. So if you do set $\{u, v\}$ they will both still have a pivot column in each column. ✓



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

False, ✓ ← why?

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

True, ✓ ← why?