

October 3, 2019

Brandon Hosken

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

5	0	1	8	4	5	0	3
0	●	0	0	0	0	●	0
1	1	●	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	●
4	4	4	4	●	4	4	4
●	5	5	5	5	●	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	●	8	8	8	8
9	9	9	9	9	9	9	9

[illegible]



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

- a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.
 b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -3 & 0 & b \end{array} \right] \xrightarrow{R_3 + 3R_2}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 6 & b+6 \end{array} \right] \xrightarrow{R_3 \div 6}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 1 & \frac{b+6}{6} \end{array} \right] \xrightarrow{R_1 + R_2}$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 1 & \frac{b+6}{6} \end{array} \right] \xrightarrow{R_1 - 3R_3}$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & -\frac{b+6}{2} \\ 0 & 1 & 0 & \frac{b+6}{3} \\ 0 & 0 & 1 & \frac{b+6}{6} \end{array} \right]$$

x_3 is free

~~$b=0$~~
 a) The only value of b that allows w to be in $\text{Span}(v_1, v_2, v_3)$ is ~~0~~ -6 because if it isn't zero there is no solution to the augmented matrix after row reduction.

b) The set $\{v_1, v_2, v_3\}$ is not linearly independent because x_3 is a free variable meaning it has infinitely many solutions. ✓



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

~~$A^{-1} = [w_1, w_2, w_3]$~~

$$w_1 = \begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 1 & 0 & 1 & | & 0 & 1 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \end{bmatrix}$$

$$\downarrow$$

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 0 & 2 & -1 & | & 0 & 1 & 0 \\ 0 & 0 & 1 & | & 0 & 0 & 1 \end{bmatrix} \cdot (-1)$$

$$\downarrow$$

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 0 & 2 & -1 & | & 0 & 1 & 0 \\ 0 & 1 & -1 & | & 0 & -1 & 1 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 1 & 0 & 1 & | & 0 & 1 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \end{bmatrix}$$

$$\downarrow$$

$$A^{-1} = \begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \\ 1 & 0 & 1 & | & 0 & 1 & 0 \end{bmatrix} \cdot (-1)$$

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \\ 0 & 1 & -1 & | & -1 & 1 & 1 \end{bmatrix} \cdot (-1)$$

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 0 & 2 & -1 & | & 0 & 0 & 1 \\ 0 & 0 & 1 & | & 0 & 1 & 0 \end{bmatrix} \cdot (-1)$$

$$\downarrow$$

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 0 & 1 & 1 & | & 1 & 0 & 1 \\ 0 & 0 & 1 & | & 0 & 1 & 0 \end{bmatrix} \cdot (-1)$$

$$\downarrow$$

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 & 0 & 0 \\ 0 & 1 & 0 & | & 1 & -1 & 1 \\ 0 & 0 & 1 & | & 0 & 1 & 0 \end{bmatrix} \cdot (-2)$$

$$\downarrow$$

$$\begin{bmatrix} 1 & -1 & 0 & | & 1 & -2 & 0 \\ 0 & 1 & 0 & | & 1 & -1 & 1 \\ 0 & 0 & 1 & | & 0 & 1 & 0 \end{bmatrix} \cdot (-1)$$

$$\begin{bmatrix} 1 & 0 & 0 & | & 2 & -3 & 1 \\ 0 & 1 & 0 & | & 1 & -1 & 1 \\ 0 & 0 & 1 & | & 0 & 1 & 0 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} 2 & -3 & 1 \\ 1 & -1 & 1 \\ \text{---} & \text{---} & \text{---} \end{bmatrix}$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 0 \\ 0 & 2 & -1 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 2 \\ 2 & 1 & -1 \end{bmatrix} \quad \checkmark C = (A^T)^{-1} \cdot B$$

$$(A^T)^{-1} = \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ -1 & 1 & 2 & 0 & 1 & 0 \\ 2 & 1 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{\cdot -2}$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ -1 & 1 & 2 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 + R_3}$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 - R_3}$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 1 & 1 \end{array} \right] \xrightarrow{\cdot -2}$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 2 \\ 0 & 0 & -1 & 0 & 1 & 1 \end{array} \right] \xrightarrow{\cdot -1}$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 2 \\ 0 & 0 & 1 & 0 & 1 & 1 \end{array} \right] \xrightarrow{R_2 - 2R_3}$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 2 \\ 0 & 0 & 1 & 0 & 1 & 1 \end{array} \right] \xrightarrow{R_2 - 2R_3}$$

$$C = B \cdot (A^T)^{-1} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 2 \\ -1 & 1 & 1 \end{bmatrix}$$

$$C = [Bv_1 \ Bv_2 \ Bv_3]$$

$$C = \begin{bmatrix} 1 & 0 & 1 \\ -1 & -5 & 2 \\ -1 & -4 & 3 \end{bmatrix}$$

There are mistakes in matrix multiplication too.

Simpler: $(A^T)^{-1} = (A^{-1})^T$
Then use problem 2.



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

a)

$$A = [T(e_1) \ T(e_2)] \quad e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$T(e_1) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 - 2(0) \\ 1 + 0 \\ 1 - 3(0) \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$T(e_2) = T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 - 2(1) \\ 0 + 1 \\ 0 - 3(1) \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$$

b)

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{array} \right] \begin{matrix} \\ \cdot -1 \\ \cdot -1 \end{matrix}$$

$$\downarrow$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 0 & -4 & -12 \end{array} \right] \begin{matrix} \\ \cdot -1 \\ \cdot -1 \end{matrix}$$

$$\downarrow$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & -4 & -12 \end{array} \right] \begin{matrix} \\ \cdot -1 \\ \cdot -1 \end{matrix}$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & 0 & -14 \end{array} \right] \begin{matrix} \\ \cdot 1/3 \\ \cdot 1/3 \end{matrix}$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & -14 \end{array} \right] \begin{matrix} \\ \cdot 2 \\ \cdot 1/3 \end{matrix}$$

$$\left[\begin{array}{cc|c} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & -14 \end{array} \right] \begin{matrix} \\ \\ \cdot -1/14 \end{matrix}$$

\therefore There are no solutions for u that satisfies $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 4 \end{bmatrix} \cdot 1/2$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 4 \end{bmatrix} \cdot 1/2$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 2 \end{bmatrix} \cdot 1/2$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \cdot 5-2$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \cdot 5-1$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$\therefore A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$ is one-to-one

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 2 \end{bmatrix} \cdot 1/2$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \end{bmatrix} \cdot -1$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \cdot 5$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

$\therefore A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$ is not one-to-one as it does not have pivot position in every row.

v_1, v_2 ?



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

~~False, because~~

? True, because $w+u$ is some linear combination of u that results in the ~~span~~ the new vector being in the span, for that to occur by the definition of a linear combination w must also be in the $\text{span}(u, v)$.

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

? ~~True~~ False, because the row reduction of the vectors in an augmented matrix require $n \times n$ to be linearly independent so without the 3rd vector in \mathbb{R}^3 , it is impossible to have only 1 solution.



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

~~True~~, because A is just a linear transformation so it will still return the properties of having infinitely many solutions to the vector. ?

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

? True, because you are applying the same transformation to all 3 vectors so the span will stay the same as before meaning $T(u)$ is in $\text{Span}(T(v), T(w))$