



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Robert Amm

UB Person Number:

5	0	2	6	4	6	1	1
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1	2	3	4	5	6	7	TOTAL	GRADE

20

10

2

20

20

0

0

--

--

72

B-

1

2

3

4

5

6

7

PIAZZA

HILL

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$a) \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \xrightarrow{-2} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & b+4 \end{array} \right] \xrightarrow{.1} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & b+6 \end{array} \right] \xrightarrow{.1} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & b+6 \end{array} \right]$$

$w \in \text{Span}(v_1, v_2, v_3)$ if and only if $b+6=0$ $b=-6$ ✓

b) set is linearly independent if and only if

$$x_1 v_1 + x_2 v_2 + x_3 v_3 = 0 \quad \text{has only trivial solution } (x_1 = x_2 = x_3 = 0)$$

$$\left[\begin{array}{cccc} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 2 & -3 & 0 & 0 \end{array} \right] \xrightarrow{-2} \left[\begin{array}{cccc} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & -1 & -2 & 0 \end{array} \right] \xrightarrow{.1} \left[\begin{array}{cccc} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \xrightarrow{.1} \left[\begin{array}{cccc} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\begin{cases} x_1 = -3x_3 \\ x_2 = -2x_3 \\ x_3 = x_3 \end{cases}$$

infinite solutions $x_1 \neq x_3 \neq x_2$

so set is not linearly independent ✓



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 - R_1} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_3 - 2R_2} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \xrightarrow{R_2 \cdot (-1)} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \xrightarrow{R_1 + R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 2 & -1 & 0 \\ 0 & 1 & -1 & 1 & -1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \xrightarrow{R_1 - R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & -1 & 1 & -1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \xrightarrow{R_2 + R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & 0 & 3 & -3 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & 0 & 3 & -3 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \xrightarrow{R_2 - 3R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -4 & 5 & -4 \\ 0 & 1 & 0 & 3 & -3 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \xrightarrow{R_2 + 4R_3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -4 & 5 & -4 \\ 0 & 1 & 0 & 11 & -5 & 5 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} \quad \checkmark$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix} \checkmark$$

$$B_{11} = [1 \ 1 \ 0] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = 1 \Rightarrow 2c_1$$

$$A^T C = B$$

?



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

a) $A = [T(e_1) \ T(e_2)] = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$ ✓ standard matrix of T

$$T(e_1) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 - 2(0) \\ 1 + 0 \\ 1 - 3(0) \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \quad T(e_2) = T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 - 2(1) \\ 0 + 1 \\ 0 - 3(1) \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$$

b) $\begin{bmatrix} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{bmatrix} \xrightarrow{\cdot -1} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 1 & -3 & -2 \end{bmatrix} \xrightarrow{\cdot -1} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & -1 & -3 \end{bmatrix} \xrightarrow{\cdot \frac{1}{3}} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{\cdot \frac{1}{3}} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{\cdot 2} \begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow \begin{cases} x_1 = 7 \\ x_2 = 3 \end{cases} \quad u = \begin{bmatrix} 7 \\ 3 \end{bmatrix} \quad \checkmark$$

$$\text{check } \begin{bmatrix} 7 - 2(3) \\ 7 + 3 \\ 7 - 3(3) \end{bmatrix} = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

a) $\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 2 & 4 & 0 \\ 3 & 4 & 4 & 0 \end{array} \right] \xrightarrow{-3R_1} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 2 & 4 & 0 \\ 0 & 1 & 4 & 0 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 4 & 0 \\ 0 & 2 & 4 & 0 \end{array} \right] \xrightarrow{-2R_2} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 4 & 0 \\ 0 & 0 & -4 & 0 \end{array} \right] \xrightarrow{\cdot (-1/4)} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 4 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right] \xrightarrow{-4R_3} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right] \xrightarrow{-R_1} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right] \xrightarrow{\cdot (-1)} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$

pivot in all columns - one-to-one ✓

b) $\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 2 & 4 & 0 \\ 3 & 4 & 2 & 0 \end{array} \right] \xrightarrow{-3R_1} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 2 & 4 & 0 \\ 0 & 1 & 2 & 0 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 2 & 4 & 0 \end{array} \right] \xrightarrow{-2R_2} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \xrightarrow{-R_1} \left[\begin{array}{ccc|c} 1 & 0 & -2 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$

Some columns missing pivots - not one-to-one ✓

$\left[\begin{array}{ccc|c} 1 & 0 & -2 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \begin{cases} x_1 = 2x_3 \\ x_2 = -2x_3 \\ x_3 = x_3 \end{cases} \quad Nul(A) = \text{span} \left(\begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \right)$

let $v_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ $T_A(v_1) = T_A(v_2)$ if and only if $v_1 = v_2 + n$ for $n \in Nul(A)$

$v_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix}$ $v_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ $v_2 = \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix}$ ✓



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

?

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

?



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.



0 1 2 3 4 5 6 7 8 9 10