



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Paul Seungyeol Ko

UB Person Number:

5	0	1	3	9	3	6	3
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1	2	3	4	5	6	7	TOTAL	GRADE

15

10

7

20

20

7

5

--

10

92

A-

1

2

3

4

5

6

7

PIAZZA

HILL

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

a) $c_1 v_1 + c_2 v_2 + c_3 v_3 = w$
Aug mat

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \xrightarrow{\text{③} - 2\text{①}} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & b+4 \end{array} \right] \xrightarrow{\text{③} + \text{②}} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & b+6 \end{array} \right]$$

$$\boxed{b \neq -6}$$

$$\boxed{b = -6}$$

b) $\begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 2 & -3 & 0 \end{bmatrix} \xrightarrow{\text{row red}} \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$ $x_3 = \text{free}$
 $x_1 = -3x_3$
 $x_2 = -2x_3$

Equation $x_1 v_1 + x_2 v_2 + x_3 v_3 = 0$ has infinitely many solution

So $\{v_1, v_2, v_3\}$ is linearly dependent





2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} . $A|I$

$$\begin{aligned} & \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{\textcircled{2}-\textcircled{1}} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{\textcircled{3}-2\textcircled{2}} \\ & \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \xrightarrow{\textcircled{1}+\textcircled{2}-\textcircled{3}} \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \xrightarrow{\textcircled{2}+\textcircled{3}} \\ & \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] I|A^{-1} \end{aligned}$$

Check

$$A \cdot A^{-1} = I$$

$$\begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \cdot \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} \quad \checkmark$$

Ans



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$\begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix} \begin{bmatrix} c_1 & c_2 & c_3 \\ c_4 & c_5 & c_6 \\ c_7 & c_8 & c_9 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$

$$A^T \cdot (A^T)^{-1} C = B \cdot (A^T)^{-1} \text{ since } (A^T)^{-1} = (A^{-1})^T$$

$$I C = B \cdot (A^{-1})^T$$

$$C = B \cdot (A^{-1})^T = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & 2 \\ -1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 3 & 2 \\ -1 & 2 & 3 \end{bmatrix}$$

$$C = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 3 & 2 \\ -1 & 2 & 3 \end{bmatrix}$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

$$(a) \quad A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 \\ a_{21}x_1 + a_{22}x_2 \\ a_{31}x_1 + a_{32}x_2 \end{bmatrix} = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

$$a_{11} = 1 \quad a_{12} = -2 \quad a_{21} = 1 \quad a_{22} = 1 \quad a_{31} = 1 \quad a_{32} = -3$$

$$A = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix} \quad \checkmark$$

$$(b) \quad T(u) = Au = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} u_1 - 2u_2 \\ u_1 + u_2 \\ u_1 - 3u_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{array} \right] \xrightarrow{\text{row red}} \begin{array}{cc|c} u_1 & u_2 & \\ \hline 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array}$$

$$u_1 = 7 \\ u_2 = 3$$

$$u = \begin{bmatrix} 7 \\ 3 \end{bmatrix} \quad \checkmark$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

(a) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix} \xrightarrow{\text{row red}} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Since $\text{Nul}(A) = \{0\}$

A is one to one

(b) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \xrightarrow{\text{row red}} \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$

$\left[\begin{array}{ccc|c} 1 & 0 & -2 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$

$x_1 - 2x_3 = 0$

$x_2 + 2x_3 = 0$

$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \\ 1 \end{bmatrix} x_3$

$\text{Nul}(A) = \text{Span}\left\{ \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \right\}$

$v_1 = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \quad v_2 = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$

$T_A(v_1) = 0$

$T_A(v_2) = 0$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

True

$$\text{If } w + u = c_1 u + c_2 v$$

$$w = c_1 u + c_2 v - u = c_3 u + c_2 v$$

$$\text{Thus } w \in \text{Span}(u, v)$$



b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

True

$$\text{if } u = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad v = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad w = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$\{u, v\}$ has to be lin ind for $\{u, v, w\}$ to be lin ind

why?



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

(1) ~~True~~ set of vectors have to be linearly dependent
for Au, Av to be linearly dependent

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

$$c_1 v + c_2 w = c_3 u$$

Since linear trans

$$T(c_1 v) + T(c_2 w) = T(c_3 u) \Rightarrow c_1 T(v) + c_2 T(w) = T(c_3 u)$$

Thus $T(u) \in \text{Span}(T(v), T(w))$



so ... TRUE.