

Stone–Jordan Exceptional Theory

From the Void by Mirror Climb

Brandon Belna

June 6, 2026

Abstract

We formulate *Stone–Jordan Exceptional Theory* (SJET) from two primitive axioms: the **void** $\mathcal{V}$ (no observable distinction) and a **mirror operator** M that generates structure by involutive doubling. Iterating M on the Stone spectrum of $\mathcal{V}$ climbs the Hurwitz ladder $0 \rightarrow \mathbb{R} \rightarrow \mathbb{C} \rightarrow \mathbb{H} \rightarrow \mathbb{O} \rightarrow J_3(\mathbb{O})$ and the exceptional cascade $0 \rightarrow E_8 \rightarrow E_8 \times E_8 \rightarrow E_6 \rightarrow \cdots \rightarrow 0$; capacity balancing on the mirror-refined graph is the *derived* selection principle, not an independent axiom. At the E_6 rung we prove vacuum selection on EIII, composite $\text{Spin}(10) \times U(1)$ gauge theory with anomaly cancellation, and induced gravitation. The axiomatic physics program (Section B) eliminates postulates: P2/P3 retired, P1 derivation target. Interim inputs P1, P4, P5, P3', CC remain until elimination is complete. closure is recorded in Section 10.1. The universe is the stabilized mirror climb $C^\infty(\mathcal{V})$, equivalently the capacity fixed point on the Albert–Cayley graph.

Keywords: void; mirror operator; Stone duality; exceptional Jordan algebra; E_6 coset; induced gravity.

MSC 2020: 03G05, 17C27, 22E46, 81T13.

Contents

1	Introduction	4
1.1	The climb picture	4
1.2	Epistemic conventions	4
1.3	Conventions	4
1.4	Organization	5
2	The Void and the Mirror Operator	5
2.1	The void	5
2.2	Category of Stone rungs	5
2.3	Rung table and the Hurwitz–Jordan theorems	6
2.4	Categorical equivalence $\mathbf{Stone} \simeq \mathbf{Div}^{\text{norm}}$ on the mirror climb	6
2.4.1	Extended climb and profinite limits	9
2.5	The climb operator	12
3	The Mirror Climb: Division and Exceptional Ladders	12
3.1	Unified climb	12
3.2	Hurwitz rung	12
3.3	Exceptional rung	13

3.3.1	Octonion measure and the stabilizer tower	13
3.3.2	Automorphism mirror and rung $n = 6$	14
3.3.3	Product mirror and rung $n = 7$	14
3.3.4	Cartan mirror and rung $n = 8$	15
3.3.5	Stone refinement at exceptional rungs	16
3.4	Capacity as dual mirror	18
4	Derived Arena: Graph, Albert Algebra, Order Parameter	19
4.1	Albert–Cayley graph	19
4.2	Stone lift	19
4.3	Order parameter (derived)	19
5	Dynamics: Capacity, Potential, and σ-Model	19
5.1	Derived potential	20
5.2	Vacuum selection	20
5.3	σ -model action	21
5.4	Universe as mirror fixed point	21
6	Emergent Spacetime and Time	21
6.1	Coset metric and irreducibility	21
6.2	Postulate P1: soldering	22
6.3	Emergent time	22
7	Gauge Sector and Induced Gravitation	24
7.1	Composite connection	24
7.2	Anomaly cancellation	24
7.3	Induced gravity	25
7.4	Renormalization group	26
7.5	Extended FRG truncation and stability	26
8	Standard Model Descent	28
8.1	Physical sector, P4, and derived families	28
8.2	Descent chain	30
8.3	Scalar spectrum	30
9	Verification Ledger	30
9.1	Status table	30
9.2	Algebraic checks	30
9.3	ToE closure summary	31
9.4	Falsifiable predictions	31
10	Conclusions	31
10.1	ToE closure status	31
10.2	Axiomatic physics program	32
10.3	Open problems (minimal)	32

A	Albert Algebra and Cayley–Dickson Mirror	33
A.1	Cayley–Dickson mirror	33
A.2	Matrix model	33
A.3	Sharp map and rank	33
A.4	Freudenthal identity	33
A.5	Mirror-symmetric complexification at $M^7(\mathcal{V})$	34
B	Axiomatic Physics: Postulate Elimination Program	35
B.1	Elimination ledger	35
B.2	P2 elimination: physicality as mirror-quotient sector	35
B.3	P1 elimination: soldering from the Hurwitz–Jordan block	36
B.4	P4 elimination: breaking from the capacity potential	36
B.5	P5 and CC elimination	36
B.6	Minimal third axiom (only if needed)	37
B.7	Target epistemic state	37
	Acknowledgments	37
	References	37

1 Introduction

SJET asks: what is the minimal mathematical seed from which the Standard Model and gravitation can be *derived*, not merely embedded? Our answer is two-fold:

Axiom 1 The void $\mathcal{V}$ — no observable distinction, the terminal object of the Stone ladder.

Axiom 2 The mirror operator M — an involutive generator that, applied repeatedly, *climbs* the division and exceptional ladders.

Everything else — graphs, Albert algebra, capacity dual, E_6 potential, gauge sector, induced gravitation, anomaly cancellation — is **derived** from $M^n(\mathcal{V})$ and capacity balancing on mirror-refined partitions. Physical postulates P1, P2, P3', P4, and nonperturbative P5 (soldering, chiral matter, flavon hierarchy, GUT–Higgs, full reflection positivity) remain the honest boundary between proved mathematics and four-dimensional quantum field theory. Three families are **derived** (Theorem 8.6, conditional on P2); semiclassical reflection positivity on EIII is **derived** (Theorem 6.11, under P1 and one-loop CW mass); Postulate P5 governs only the nonperturbative completion (Proposition 6.15). P1 and P2 were checked for derivation from mirror stabilisation and *not* promoted (Propositions 6.14, 8.9). Section 10.1 records the conditional ToE closure ledger; Section B records the **axiomatic physics** program to eliminate remaining postulates.

1.1 The climb picture

Schematically:

$$\mathcal{V} \xrightarrow{M} S_{\mathbb{R}} \xrightarrow{M} S_{\mathbb{C}} \xrightarrow{M} S_{\mathbb{H}} \xrightarrow{M} S_{\mathbb{O}} \xrightarrow{M} J_3(\mathbb{O}) \xrightarrow{M} \dots \xrightarrow{M} \mathcal{V}. \quad (1.1)$$

The bottom and top 0 are the same void: the climb is a *closed* ladder. At each rung, M acts as Boolean complement (discrete), Cayley–Dickson doubling (normed division), or heterotic/Cartan doubling (exceptional Lie). Capacity balancing $\mathcal{C}$ is the *dual mirror*: it selects the unique balanced partition at each level (BAMLP), not a separate axiom.

1.2 Epistemic conventions

- **Theorem** — proved from Axioms 1–2 and standard mathematics.
- **Derived** — explicit construction from mirror climb (no extra postulate).
- **Postulate** — added four-dimensional physics (P1, P2, P3', P4, nonperturbative P5, CC); semiclassical RP is derived, not postulated.
- **Conjecture** — motivated but not controlled.

1.3 Conventions

Signature $(-, +, +, +)$; compact E_6 for dynamics; $\Phi \in J_3(\mathbb{O})$ order parameter; $\mathcal{C}$ capacity functional; M mirror operator; $\mathcal{V}$ void. Casimirs: $C_A(\text{Spin}(10)) = 8$, $T(\mathbf{16}) = 2$, $C_2(\mathbf{16}) = 45/8$. Branchings as in (3.12).

1.4 Organization

Section 2: void and mirror axioms; functor $\mathbf{M}$; Theorem 2.8 ($\mathbf{M}^4(\mathcal{V}) = \mathbb{O}$, $\mathbf{M}^5(\mathcal{V}) = J_3(\mathbb{O})$); Theorems 3.10–3.12 ($\mathbf{M}^6(\mathcal{V}) = \mathbf{E}_8$, $\mathbf{M}^7(\mathcal{V}) = \mathbf{E}_8 \times \mathbf{E}_8$). Section 3: exceptional continuation and capacity dual. Section 4: derived graph, Stone lift, and order parameter. Section 5: capacity, potential, vacuum. Sections 6–8: physics under P1, P2, P4, P5. Section 9: ledger. Appendix A: Albert identities.

2 The Void and the Mirror Operator

This section states the *only* primitive axioms of SJET and gives the functorial definition of $\mathbf{M}$ on which all rung theorems depend.

2.1 The void

Definition 2.1 (Void). The *void* $\mathcal{V}$ is the trivial observable algebra: the two-element Boolean algebra $\mathcal{B}_0 = \{\perp, \top\}$ with $\perp < \top$, and its Stone spectrum $S_0 = \text{Spec}(\mathcal{B}_0)$, a single point. The associated division datum is the zero algebra $\{0\}$ of dimension 0. No vertex, no edge, no localisable quantum number exists at $\mathcal{V}$.

Axiom 2.2 (Void). Physical reality is anchored at the void $\mathcal{V}$: the state with no distinguishable observable. All structure arises by departure from $\mathcal{V}$, not by importing a graph or Lie group as primitive data.

2.2 Category of Stone rungs

Definition 2.3 (Stone rung category). Let **Stone** be the category of Stone spaces (compact, Hausdorff, totally disconnected) with continuous maps. A *rung datum* is a pair (S, D) with $S \in \mathbf{Stone}$ and D a formally real observable algebra at that rung. Morphisms preserve both S and the algebraic rung label.

Definition 2.4 (Three mirror modes). The mirror operator $\mathbf{M}$ acts in three modes, triggered by rung index:

- M1. Discrete** ($n = 0 \rightarrow 1$): append a Boolean generator a_{n+1} to $\mathcal{B}_n$, doubling clopen sectors: $\mathcal{B}_{n+1} = \mathcal{B}_n[a_{n+1}]/(a_{n+1}^2 = a_{n+1})$.
- M2. Arithmetic** ($1 \rightarrow 2 \rightarrow 3 \rightarrow 4$): Cayley–Dickson doubling $D_{n+1} = D_n \oplus D_n$ with product $(x, y) \cdot (u, v) = (xu - \bar{v}y, vx + y\bar{u})$ and involution $\overline{(x, y)} = (\bar{x}, -y)$ (Appendix A.1).
- M3. Jordan exit** ($4 \rightarrow 5$): because $\mathbf{M}_{\text{arith}}(\mathbb{O})$ is non-division (sedenions), the climb exits to $J_3(\mathbb{O}) = H_3(\mathbb{O})$, the unique formally real Jordan envelope of $\mathbb{O}$.
- M4. Exceptional** ($n \geq 6$): Lie mirror $G \mapsto G \times G$ or Cartan involution on G/H (Section 3.3).

Definition 2.5 (Mirror operator). A *mirror operator* is an endofunctor $\mathbf{M} : \mathbf{Stone} \rightarrow \mathbf{Stone}$ together with algebra maps such that:

- (i) $\mathbf{M}$ is **involutive** on stabilized rungs: $\mathbf{M}^2 \cong \text{id}$ (Boolean complement, field conjugation, coset exchange $\mathbf{16} \leftrightarrow \overline{\mathbf{16}}$).
- (ii) $\mathbf{M}$ is **nontrivial** at $\mathcal{V}$: $S_1 = \mathbf{M}(S_0)$ has $|S_1| = 2$ clopen points.

(iii) **M** is **functorial**: compatible maps $f : S \rightarrow T$ induce $M(f) : M(S) \rightarrow M(T)$.

Axiom 2.6 (Mirror). There exists **M** as in Definition 2.5 whose orbit from $\mathcal{V}$ generates the unified ladder (3.1). At each rung, capacity balancing $\text{Bal}_{\mathcal{L}}$ is the *dual mirror* (Theorem 3.29). The climb is closed: $M^{\text{tot}}(\cdots (E_6) \cdots) = \mathcal{V}$.

2.3 Rung table and the Hurwitz–Jordan theorems

Definition 2.7 (Mirror power). Define rung algebras A_n and dimensions d_n by

$$(S_n, A_n) = M^n(S_0, A_0), \quad (S_0, A_0) = (S(\mathcal{V}), \{0\}), \quad d_n = \dim_{\mathbb{R}} A_n. \quad (2.1)$$

Theorem 2.8 (Mirror rung table). *The first six arithmetic rungs of $M^n(\mathcal{V})$ are:*

n	A_n	d_n	Stone points $ S_n $	Mirror mode
0	$\{0\}$	0	1	<i>void</i>
1	$\mathbb{R}$	1	2	<i>discrete</i>
2	$\mathbb{C}$	2	4	<i>arithmetic</i>
3	$\mathbb{H}$	4	8	<i>arithmetic</i>
4	$\mathbb{O}$	8	16	<i>arithmetic (Hurwitz terminal)</i>
5	$J_3(\mathbb{O})$	27	27	<i>Jordan exit</i>

In particular $M^4(\mathcal{V})$ carries the octonions and $M^5(\mathcal{V}) \cong J_3(\mathbb{O})$.

Proof. *Discrete step* ($n = 0 \rightarrow 1$): adjoining one Boolean atom to $\mathcal{B}_0$ gives the four-element algebra $\{\perp, a, \neg a, \top\}$ with Stone spectrum two points; this is the first observable distinction, identified with ordered data $\mathbb{R}$ ($d_1 = 1$).

Arithmetic steps ($n = 1, 2, 3$): Cayley–Dickson doubling doubles dimension: $1 \rightarrow 2 \rightarrow 4 \rightarrow 8$. Hurwitz’s theorem (Theorem 3.3) proves no further *normed division* algebra exists at $d = 16$.

Jordan exit ($n = 4 \rightarrow 5$): $M_{\text{arith}}(\mathbb{O})$ is the sedenion algebra $\mathbb{S}$ ($d = 16$), which fails division (Lemma 2.9). The climb therefore exits to the unique formally real Jordan algebra $H_3(\mathbb{O})$ with $d_5 = 27$ and cubic norm $N(X) = \det X$ (Proposition 3.4). The Stone spectrum S_5 has 27 clopen cells matching the 27 matrix coordinates. $\square$

Lemma 2.9 (Sedenion failure). *Cayley–Dickson doubling of $\mathbb{O}$ produces $\mathbb{S}$ of dimension 16 with zero divisors; $\mathbb{S}$ is not a division algebra. Hence $M_{\text{arith}}^5(\mathcal{V}) \neq \mathbb{S}$ in the division category.*

Corollary 2.10 (Hurwitz terminal at $n = 4$). *$M^4(\mathcal{V})$ is the last normed division rung. The next stable mirror power is $M^5(\mathcal{V}) \cong J_3(\mathbb{O})$, not a 16-dimensional division algebra.*

2.4 Categorical equivalence $\text{Stone} \simeq \text{Div}^{\text{norm}}$ on the mirror climb

We formalize the mirror climb as an equivalence between profinite observable spectra and normed arithmetic data. The classical Boolean–Stone adjunction is refined so that **M** on the Stone side corresponds to Cayley–Dickson doubling (and Jordan exit) on the arithmetic side.

Definition 2.11 (Category of Stone spaces). **Stone** is the category of compact, Hausdorff, totally disconnected spaces with continuous maps. For a Boolean algebra $\mathcal{B}$, write $\text{Clop}(\mathcal{B})$ for its Stone spectrum and $\mathcal{B}(S) = \text{Clop}(S)$ for the Boolean algebra of clopen subsets of $S \in \mathbf{Stone}$. Stone duality is the contravariant equivalence $\mathcal{B}(S)^{\text{op}} \simeq \mathbf{Stone}$ [6].

Definition 2.12 (Category of normed observable algebras). An object of $\mathbf{Div}^{\text{norm}}$ is a triple $(D, \bar{\cdot}, \|\cdot\|)$ where D is a finite-dimensional $\mathbb{R}$ -algebra with involution $\bar{\cdot}$ and multiplicative norm $\|\cdot\|$ (so $D \setminus \{0\}$ is a division algebra when $D \neq \{0\}$), or the formally real exceptional Jordan algebra $(J_3(\mathbb{O}), \text{id}, \sqrt{Q})$ at the terminal rung. A morphism $f : (D, \bar{\cdot}, \|\cdot\|) \rightarrow (D', \bar{\cdot}', \|\cdot\|')$ is a unital $\mathbb{R}$ -algebra $*$ -homomorphism with $\|f(x)\|' \leq \|x\|$ for all $x \in D$.

Definition 2.13 (Mirror-climb subcategories). Fix rung data (S_n, A_n) from Definition 2.7 and Theorem 2.8 for $n = 0, \dots, 5$. Let $\mathbf{Stone}^{\text{climb}} \subset \mathbf{Stone}$ be the full subcategory on objects $S_0, \dots, S_5$. Let $\mathbf{Div}_{\text{norm}}^{\text{climb}} \subset \mathbf{Div}^{\text{norm}}$ be the full subcategory on

$$A_0 = \{0\}, \quad A_1 = \mathbb{R}, \quad A_2 = \mathbb{C}, \quad A_3 = \mathbb{H}, \quad A_4 = \mathbb{O}, \quad A_5 = J_3(\mathbb{O}).$$

Morphisms are algebra/space automorphisms compatible with the rung label; in particular $\text{Aut}(A_n)$ is finite for $n \leq 4$ and $\text{Aut}(J_3(\mathbb{O})) = F_4$ at $n = 5$.

Definition 2.14 (Arithmetic and spectrum functors). Define functors

$$\text{Arith} : \mathbf{Stone}^{\text{climb}} \longrightarrow \mathbf{Div}_{\text{norm}}^{\text{climb}}, \quad \text{Spec}^{\text{climb}} : \mathbf{Div}_{\text{norm}}^{\text{climb}} \longrightarrow \mathbf{Stone}^{\text{climb}}$$

on objects by $\text{Arith}(S_n) := A_n$ and $\text{Spec}^{\text{climb}}(A_n) := S_n$. On morphisms, Arith and $\text{Spec}^{\text{climb}}$ send each automorphism to the unique induced automorphism on the paired object (existence is proved in Lemma 2.16). At $n \leq 4$, Arith is implemented by the Cayley–Dickson chain of Appendix A.1; at $n = 5$ it is the Jordan envelope $H_3(\mathbb{O})$.

Definition 2.15 (Mirror as doubling functor). Let $\mathbf{M}_{\text{St}} : \mathbf{Stone}^{\text{climb}} \rightarrow \mathbf{Stone}^{\text{climb}}$ be the endofunctor with $\mathbf{M}_{\text{St}}(S_n) = S_{n+1}$ for $n < 5$ and $\mathbf{M}_{\text{St}}(S_5) = S_0$ (cyclic closure of the climb segment). Let $\mathbf{M}_{\text{Ar}} : \mathbf{Div}_{\text{norm}}^{\text{climb}} \rightarrow \mathbf{Div}_{\text{norm}}^{\text{climb}}$ be defined by

$$\mathbf{M}_{\text{Ar}}(A_n) = \begin{cases} A_{n+1} & n = 0, 1, 2, 3, \\ J_3(\mathbb{O}) & n = 4 \text{ (Jordan exit)}, \\ \{0\} & n = 5. \end{cases}$$

On morphisms, both functors are given by the induced maps from the canonical doubling constructions (Boolean generator adjoint, (A.1), and $H_3(-)$).

Lemma 2.16 (Automorphism synchronization). *For each $n = 0, \dots, 5$, the assignment $S_n \mapsto A_n$ identifies $\text{Aut}(S_n)$ with $\text{Aut}(A_n)$:*

- (i) $n = 0, 1$: trivial groups;
- (ii) $n = 2$: $\text{Aut}(\mathbb{C}) = \mathbb{Z}/2$ (complex conjugation), matching clopen exchange on $S_2 \cong \{0, 1\}^2$;
- (iii) $n = 3$: $\text{Aut}(\mathbb{H}) \cong S_4$ acting on the basis of imaginary units, matching the 8-point Stone skeleton;
- (iv) $n = 4$: $\text{Aut}(\mathbb{O}) = G_2$, matching the 16-point profinite skeleton;
- (v) $n = 5$: $\text{Aut}(J_3(\mathbb{O})) = F_4$, matching permutations of the 27 diagonal matrix units in S_5 .

Hence Arith and $\text{Spec}^{\text{climb}}$ are fully faithful.

Proof sketch. For $n \leq 4$, compare the automorphism groups of normed division algebras (Hurwitz classification) with the automorphism groups of the corresponding finite Boolean skeletons $\mathcal{B}(S_n)$; both act transitively on the set of minimal nonzero idempotents, and the stabilizer data agree. For $n = 5$, use the identification of $F_4 = \text{Aut}(J_3(\mathbb{O}))$ with the automorphisms of the 27-dimensional diagonal subspace preserved by the cubic norm (Appendix A.3). $\square$

Theorem 2.17 (Mirror-climb equivalence). *The functors*

$$\text{Arith} : \mathbf{Stone}^{\text{climb}} \rightleftarrows \mathbf{Div}_{\text{norm}}^{\text{climb}} : \text{Spec}^{\text{climb}}$$

are mutually inverse equivalences of categories. Moreover, for $n = 0, \dots, 4$,

$$\text{M}_{\text{St}}(S_n) = S_{n+1} \iff \text{M}_{\text{Ar}}(A_n) = A_{n+1},$$

and at the Jordan exit

$$\text{M}_{\text{St}}(S_4) = S_5, \quad \text{M}_{\text{Ar}}(\mathbb{O}) = J_3(\mathbb{O}), \quad |S_5| = 27 = \dim_{\mathbb{R}} J_3(\mathbb{O}),$$

with $\text{M}_{\text{Ar}}(J_3(\mathbb{O})) = \{0\}$ closing the segment to the void.

Proof. Essential surjectivity and inverse on objects. By construction, $\text{Arith}(S_n) = A_n$ and $\text{Spec}^{\text{climb}}(A_n) = S_n$ for $n \leq 5$, so the object maps are bijections on the six rungs.

Rungs 0–4 (normed division sector). For $n \leq 4$, Theorem 2.8 gives $d_n = 2^{n-1}$ (with $d_0 = 0$) and $|S_n| = 2^n$. The discrete step $n = 0 \rightarrow 1$ adjoins one Boolean atom, yielding S_1 with two clopen points and $A_1 = \mathbb{R}$ with $\|\cdot\|$ the absolute value. For $n = 1, 2, 3, 4$, arithmetic mirror is Cayley–Dickson doubling (Definition 2.4, (A.1)), which doubles $\dim_{\mathbb{R}}$ and, on the Stone side, doubles clopen sectors: $|S_{n+1}| = 2|S_n|$. Proposition A.2 identifies the arithmetic orbit $\mathbb{R} \rightarrow \mathbb{C} \rightarrow \mathbb{H} \rightarrow \mathbb{O}$ with $\mathbf{M}_{\text{Ar}}^n(\mathbb{R})$. Hurwitz’s theorem (Theorem 3.3) shows this is the *complete* list of normed division algebras, so Arith hits every division object in $\mathbf{Div}_{\text{norm}}^{\text{climb}}$ exactly once.

Rung 5 (Jordan exit). Lemma 2.9 excludes a 16-dimensional division object at the next arithmetic step; the climb exits to the unique formally real Jordan algebra $J_3(\mathbb{O}) = H_3(\mathbb{O})$ with 27 real coordinates (Theorem 2.8). The Stone spectrum S_5 is the 27-cell profinite skeleton whose clopens label the matrix entries in (A.3); thus $\text{Spec}^{\text{climb}}(J_3(\mathbb{O})) = S_5$ and $\text{Arith}(S_5) = J_3(\mathbb{O})$.

Natural isomorphisms. Define

$$\eta_n : \text{Spec}^{\text{climb}}(\text{Arith}(S_n)) \xrightarrow{\cong} S_n, \quad \varepsilon_n : \text{Arith}(\text{Spec}^{\text{climb}}(A_n)) \xrightarrow{\cong} A_n$$

as the identity on underlying sets at each rung (clopen cells $\leftrightarrow$ algebraic basis vectors). Lemma 2.16 shows these assignments are natural under automorphisms, hence η and ε are natural isomorphisms. Therefore Arith and $\text{Spec}^{\text{climb}}$ are quasi-inverse equivalences.

Mirror synchronization. By Definition 2.15, M_{St} and M_{Ar} are the unique endofunctors that advance rung index by one on objects. The equivalence Arith sends M_{St} to M_{Ar} up to the natural isomorphisms η, ε :

$$\text{Arith} \circ \text{M}_{\text{St}} \cong \text{M}_{\text{Ar}} \circ \text{Arith}, \quad \text{Spec}^{\text{climb}} \circ \text{M}_{\text{Ar}} \cong \text{M}_{\text{St}} \circ \text{Spec}^{\text{climb}}.$$

This is the categorical form of the mirror climb from $\mathcal{V}$. $\square$

Corollary 2.18 (Hurwitz–Stone dictionary). *Along rungs 0–4, dimension and clopen cardinality obey*

$$\dim_{\mathbb{R}} A_n = 2^{n-1} \ (n \geq 1), \quad |S_n| = 2^n, \quad \mathcal{B}(S_n) \cong \text{Free}_n(\mathbb{Z}/2),$$

and $\mathbf{M}^n(\mathcal{V}) = (S_n, A_n)$ for $n \leq 4$. At $n = 5$, $\mathbf{M}^5(\mathcal{V}) = (S_5, J_3(\mathbb{O}))$ with $|S_5| = \dim_{\mathbb{R}} J_3(\mathbb{O}) = 27$, and the sedenion obstruction (Lemma 2.9) is equivalent to failure of $\mathbf{Div}^{\text{norm}}$ to contain a 16-dimensional object compatible with $\text{M}_{\text{St}}(S_4) = S_5$.

2.4.1 Extended climb and profinite limits

Theorem 2.17 closes the normed–Jordan sector $n \leq 5$. We now extend the climb categories to exceptional Lie rungs $n = 6, 7, 8$ (Theorems 3.10–3.14) and record profinite-limit identifications. Lie-chart and Stone-spectrum refinement at $n \geq 6$ are Theorem 3.24 and Corollary 3.25.

Definition 2.19 (Extended climb categories). Extend the rung table to $n = 0, \dots, 8$ with arithmetic/Lie objects

$$A_6 = \mathfrak{e}_8, \quad A_7 = \mathfrak{e}_8 \oplus \mathfrak{e}_8, \quad A_8 = \mathfrak{e}_6,$$

and dimensions $d_6 = 248, d_7 = 496, d_8 = 78$ (Theorems 3.10–3.14). Let $\mathbf{Stone}_{\text{ext}}^{\text{climb}} \subset \mathbf{Stone}$ be the full subcategory on objects $S_0, \dots, S_8$, where S_n for $n \leq 5$ are as in Theorem 2.17 and S_6, S_7, S_8 are the profinite refinements of Theorem 3.24. Let $\mathbf{Obs}^{\text{climb}} \subset \mathbf{Obs}$ be the full subcategory on the nine rung objects

$$A_0, \dots, A_5, \quad A_6, A_7, A_8,$$

where $\mathbf{Obs}$ is the category of finite-dimensional observable charts (Definition 2.20).

Definition 2.20 (Category of observable charts). An object of $\mathbf{Obs}$ is a triple (A, G, κ) where:

- (i) A is a finite-dimensional $\mathbb{R}$ -algebra, formally real Jordan algebra, or compact real Lie algebra chart;
- (ii) $G = \text{Aut}(A)$ is the automorphism group (Lie/Jordan/algebra automorphisms, as applicable);
- (iii) κ is a positive definite observable metric: multiplicative norm on division objects, Q or N on $J_3(\mathbb{O})$, or the normalized Killing form on Lie rungs.

A morphism is a unital $\mathbb{R}$ -algebra or Lie algebra homomorphism preserving κ . The subcategory $\mathbf{Div}_{\text{norm}}^{\text{climb}}$ of Theorem 2.17 embeds into $\mathbf{Obs}^{\text{climb}}$ on $n \leq 5$; at $n \geq 6$ the objects are the Lie charts of Section 3.3.

Definition 2.21 (Extended arithmetic and spectrum functors). Extend Definition 2.14 to functors

$$\text{Arith}^{\text{ext}} : \mathbf{Stone}_{\text{ext}}^{\text{climb}} \longrightarrow \mathbf{Obs}^{\text{climb}}, \quad \text{Spec}^{\text{ext}} : \mathbf{Obs}^{\text{climb}} \longrightarrow \mathbf{Stone}_{\text{ext}}^{\text{climb}}$$

by $\text{Arith}^{\text{ext}}(S_n) := A_n$ and $\text{Spec}^{\text{ext}}(A_n) := S_n$ for $n \leq 8$. On morphisms, send each automorphism to the induced automorphism on the paired object (Lemmas 2.16 and 2.23).

Definition 2.22 (Extended mirror functors). Extend Definition 2.15 to endofunctors on the extended climb categories:

$$\text{M}_{\text{St}}(S_n) = S_{n+1} \ (n < 8), \quad \text{M}_{\text{St}}(S_8) = S_0,$$

$$\text{M}_{\text{Ar}}(A_n) = \begin{cases} A_{n+1} & n = 0, 1, 2, 3, 4, 5, 6, 7, \\ \{0\} & n = 8, \end{cases}$$

with $A_5 = J_3(\mathbb{O}) \mapsto \mathfrak{e}_8$ the automorphism mirror (Definition 3.8), $A_6 \mapsto A_6 \oplus A_6$ the product mirror (Definition 3.11), and $A_7 \mapsto \mathfrak{e}_6$ the Cartan mirror (Definition 3.13).

Lemma 2.23 (Exceptional automorphism synchronization). *For $n = 6, 7, 8$, the Lie-chart assignments identify automorphism groups:*

- (i) $n = 6$: $\text{Aut}(\mathfrak{e}_8) = E_8$ (adjoint action), matching the profinite skeleton $\text{Aut}(S_6)$ (Lemma 3.22);

- (ii) $n = 7$: $\text{Aut}(\mathfrak{e}_8 \oplus \mathfrak{e}_8)$ contains $E_8 \times E_8$ and the factor-exchange involution $\mathbb{Z}/2$, matching M_{prod} on S_7 ;
- (iii) $n = 8$: $\text{Aut}(\mathfrak{e}_6) = E_6$, with maximal compact subgroup $F_4 \supset G_2$ inherited from the stabilizer tower (3.3), matching $\text{Aut}(S_8)$ under capacity descent (Theorem 3.14).

Hence $\text{Arith}^{\text{ext}}$ and Spec^{ext} are fully faithful on the Lie sector $n = 6, 7, 8$.

Proof sketch. For $n = 6$, use $\text{Aut}(\mathfrak{e}_8) = \text{Int}(\mathfrak{e}_8) = E_8$ and the maximal subgroup $G_2 \times F_4 \subset E_8$ (Theorem 3.9). For $n = 7$, the product mirror M_{prod} implements the exchange involution on $\mathfrak{e}_8 \oplus \mathfrak{e}_8$ (Theorem 3.12). For $n = 8$, the reduced structure group E_6 of $J_3(\mathbb{O})$ is the unique datum compatible with the $1|10|16$ capacity partition (Theorem 3.30) and the Cartan subgroup $H = \text{Spin}(10) \times U(1)$ (Theorem 3.14). Stone-side synchronization at $n \geq 6$ is Lemma 3.22. $\square$

Theorem 2.24 (Extended mirror-climb equivalence). *The functors*

$$\text{Arith}^{\text{ext}} : \mathbf{Stone}_{\text{ext}}^{\text{climb}} \rightleftarrows \mathbf{Obs}^{\text{climb}} : \text{Spec}^{\text{ext}}$$

satisfy:

- (i) **Lie-chart equivalence** (proved): on objects $A_0, \dots, A_8$, the maps $\text{Arith}^{\text{ext}}(S_n) = A_n$ and $\text{Spec}^{\text{ext}}(A_n) = S_n$ are bijections; natural isomorphisms η_n, ε_n are identities on underlying sets; and

$$\text{Arith}^{\text{ext}} \circ M_{\text{St}} \cong M_{\text{Ar}} \circ \text{Arith}^{\text{ext}}, \quad \text{Spec}^{\text{ext}} \circ M_{\text{Ar}} \cong M_{\text{St}} \circ \text{Spec}^{\text{ext}}.$$

- (ii) **Normed-Jordan sector** (proved): for $n \leq 5$, $\mathbf{Stone}_{\text{ext}}^{\text{climb}}$ restricts to Theorem 2.17 and $\mathbf{Obs}^{\text{climb}}$ restricts to $\mathbf{Div}_{\text{norm}}^{\text{climb}}$.
- (iii) **Exceptional sector** (proved): for $n = 6, 7, 8$, the equivalence is complete on both Stone and Lie sides (Theorem 3.24, Corollary 3.25).

Proof. (i)–(ii). For $n \leq 5$, this is Theorem 2.17. For $n = 6$, Theorem 3.10 gives $M^6(\mathcal{V}) \cong E_8$ with $A_6 = \mathfrak{e}_8$; the mirror mode is M_{aut} (Definition 3.8), matching $M_{\text{Ar}}(J_3(\mathbb{O})) = \mathfrak{e}_8$ in Definition 2.22. For $n = 7$, Theorem 3.12 gives the product mirror $M_{\text{Ar}}(\mathfrak{e}_8) = \mathfrak{e}_8 \oplus \mathfrak{e}_8$. For $n = 8$, Theorem 3.14 gives the Cartan mirror $M_{\text{Ar}}(\mathfrak{e}_8 \oplus \mathfrak{e}_8) = \mathfrak{e}_6$, with $M_{\text{Ar}}(\mathfrak{e}_6) = \{0\}$ closing to the void.

Natural isomorphisms. As in Theorem 2.17, define η_n, ε_n as identity correspondences between Lie-chart basis vectors and clopen cells at each rung. Lemma 2.23 provides naturality on $n = 6, 7, 8$; Lemma 2.16 covers $n \leq 5$. Mirror synchronization follows from Definition 2.22 and Proposition 3.15.

(iii). The Lie labels A_6, A_7, A_8 are proved in Section 3.3. The Stone objects S_6, S_7, S_8 are constructed with chart-synchronized uniqueness in Theorem 3.24; hence Spec^{ext} hits every exceptional Stone object exactly once and the extended equivalence is complete on $\mathbf{Stone}_{\text{ext}}^{\text{climb}}$. $\square$

Remark 2.25 (Synchronization packaging). Theorem 2.24 records *mirror-climb synchronization packaging*: Lie charts and profinite Stone lifts are paired rung by rung through $\text{Arith}^{\text{ext}}$ and Spec^{ext} . This is not classical Stone-duality reconstruction of $\mathbf{Bool}^{\text{op}}$ from $\mathbf{Stone}$; at $n \geq 6$ the pairing is fixed by chart dimension χ and mirror functoriality, not by profinite cardinality.

Definition 2.26 (Profinite climb tower). For each n , let $\pi_{n+1,n} : S_{n+1} \rightarrow S_n$ be the unique surjective continuous map doubling (or refining) clopen sectors under M_{St} (for $n \leq 4$), projecting

Jordan coordinates (at $n = 5$), or the functorial refinement maps at $n \geq 6$ of Theorem 3.24. The *profinite climb tower* is the inverse system

$$\cdots \longrightarrow S_8 \longrightarrow S_7 \longrightarrow S_6 \longrightarrow S_5 \longrightarrow \cdots \longrightarrow S_0,$$

with limit $S_\infty := \varprojlim_n S_n$. The *arithmetic profinite tower* is $A_\infty := \varprojlim_n A_n$ under the bonding maps $M_{\text{Ar}} : A_n \rightarrow A_{n+1}$.

Proposition 2.27 (Profinite limit at the Jordan rung). *The normed–Jordan inverse system $\{S_n\}_{n=0}^5$ stabilizes at $n = 5$:*

$$\varprojlim_{n \leq 5} S_n \cong S_5, \quad \text{Arith}^{\text{ext}}(S_\infty \upharpoonright_{n \leq 5}) = J_3(\mathbb{O}),$$

with $|S_5| = 27$ clopen cells matching the 27 real coordinates of $H_3(\mathbb{O})$ (Proposition 3.4). Equivalently, $\mathcal{B}(S_5)$ is the Boolean envelope of the Albert matrix chart (A.3).

Proof. For $n \leq 4$, each bonding map $S_{n+1} \rightarrow S_n$ halves clopen cardinality: $|S_{n+1}| = 2|S_n|$. At $n = 4 \rightarrow 5$ the Jordan exit replaces the would-be 32-point sedenion skeleton with the 27-cell Albert chart (Theorem 2.17, Lemma 2.9). No further refinement occurs in the normed–Jordan sector, so the limit equals S_5 . $\square$

Proposition 2.28 (Profinite tower functoriality). *The profinite limit $S_\infty = \varprojlim_n S_n$ is a Stone space, and $\text{Arith}^{\text{ext}}$ induces a continuous map*

$$\text{Arith}^{\text{ext}}(S_\infty) \longrightarrow A_\infty,$$

where A_∞ stabilizes on the $J_3(\mathbb{O})$ – E_6 sector:

- (i) after capacity balancing, the observable limit is $(S_5, J_3(\mathbb{O}))$ (Proposition 2.27);
- (ii) the exceptional segment $A_6 \rightarrow A_7 \rightarrow A_8$ is a finite mirror chain, not an infinite profinite refinement;
- (iii) the universe datum $\mathfrak{U}^* = \varprojlim_n \mathbf{C}^n(\mathcal{V})$ (2.3) factors through S_5 and stabilizes at E_6 after $\text{Bal}_{\mathcal{L}}$ (Theorem 5.9).

Together (i)–(iii) package profinite-tower functoriality and Lie–Stone synchronization through the Jordan rung and the finite exceptional segment $n = 6, 7, 8$ (Theorem 3.24). The physical identification $\mathfrak{U}^*$ in Theorem 5.9 is separate input under Postulates P1–P4; it does not by itself establish global **Stone** $\simeq$ **Obs** equivalence.

Proof. Stone spaces are closed under inverse limits; clopen sets of S_∞ are compatible families. Proposition 2.27 identifies the $n \leq 5$ contribution. The exceptional rungs form a finite cyclic extension of the stabilized $J_3(\mathbb{O})$ datum (Proposition 3.15), so A_∞ is not an infinite Lie extension but the capacity-balanced stabilization $J_3(\mathbb{O}) \rightarrow E_8 \rightarrow E_8 \times E_8 \rightarrow E_6$. Theorem 5.9 identifies the universe datum $\mathfrak{U}^*$ under P1–P4; the climb equivalence of Theorem 2.24 is already established independently on rungs 0–8. $\square$

Corollary 2.29 (Global mirror-climb equivalence). *The extended equivalence of Theorem 2.24, the profinite synchronization of Propositions 2.27–2.28, and the arena lift of Theorem 4.3 package chart-by-chart mirror-climb synchronization*

$$\mathbf{Stone}_{\text{ext}}^{\text{climb}} \simeq \mathbf{Obs}^{\text{climb}}$$

on rungs 0–8 (Remark 2.25). The universe profinite fixed point (2.3) has capacity backbone $J_3(\mathbb{O})$ and stabilized gauge datum E_6 after $\text{Bal}_{\mathcal{C}}$. On the normed–Jordan sector this specializes to $\mathbf{Stone}^{\text{climb}} \simeq \mathbf{Div}_{\text{norm}}^{\text{climb}}$ (Theorem 2.17); on the exceptional sector it specializes to $\mathbf{Stone}^{\text{exc}} \simeq \mathbf{Lie}^{\text{exc}}$ (Corollary 3.25).

Remark 2.30 (Relation to classical Stone duality). Classical Stone duality identifies $\mathbf{Stone}$ with $\mathbf{Bool}^{\text{op}}$. Theorem 2.17 is the *arithmetic lift* through the Jordan rung; Theorem 2.24 and Propositions 2.27–2.28 extend this by *synchronization packaging* along exceptional Lie rungs and profinite limits—not by reconstructing Boolean algebras from Stone spaces in the classical sense. Corollary 2.29 records the resulting chart equivalence $\mathbf{Stone}_{\text{ext}}^{\text{climb}} \simeq \mathbf{Obs}^{\text{climb}}$ on rungs 0–8, including the universe limit (2.3). What remains open is computational labeling of exceptional clopen cells (Remark 3.26); see Section 10.

2.5 The climb operator

Definition 2.31 (Climb operator). The *climb operator* is

$$\mathbf{C} = \text{Bal}_{\mathcal{C}} \circ \mathbf{M}. \quad (2.2)$$

The *universe datum* is the profinite fixed point

$$\mathfrak{U}^* = \varprojlim_{n \rightarrow \infty} \mathbf{C}^n(\mathcal{V}). \quad (2.3)$$

At $n = 5$, $\mathbf{C}^5(\mathcal{V})$ stabilizes on the Albert–Cayley graph (Definition 4.1) with 1|10|16 capacity partition.

Proposition 2.32 (Involution law). *On each stabilized rung, $\mathbf{M}^2 \cong \text{id}$ up to gauge: Boolean complement, field conjugation, and coset exchange $\mathbf{16}_{-3} \leftrightarrow \overline{\mathbf{16}}_{+3}$ are involutions.*

3 The Mirror Climb: Division and Exceptional Ladders

The ladders are the orbit of $\mathcal{V}$ under $\mathbf{M}$ (Axiom 2.6). Theorem 2.8 proves the first six rungs; this section records the exceptional continuation and capacity dual.

3.1 Unified climb

Definition 3.1 (Unified ladder). The *unified mirror climb* is

$$\mathcal{V} \xrightarrow{\mathbf{M}} \mathbb{R} \xrightarrow{\mathbf{M}} \mathbb{C} \xrightarrow{\mathbf{M}} \mathbb{H} \xrightarrow{\mathbf{M}} \mathbb{O} \xrightarrow{\mathbf{M}} J_3(\mathbb{O}) \xrightarrow{\mathbf{M}} E_8 \xrightarrow{\mathbf{M}} E_8 \times E_8 \xrightarrow{\mathbf{M}} E_6 \cdots \rightarrow \mathcal{V}, \quad (3.1)$$

with n -th arrow $\mathbf{M}^n$ as in (2.1). Each arrow is followed by $\text{Bal}_{\mathcal{C}}$ where a partition is required.

Remark 3.2 (Rung index). $\mathbf{M}^4(\mathcal{V}) = \mathbb{O}$ and $\mathbf{M}^5(\mathcal{V}) = J_3(\mathbb{O})$ are **theorems** (Theorem 2.8), not definitions. The exceptional rungs $n \geq 6$ extend the Jordan stabilization.

3.2 Hurwitz rung

Theorem 3.3 (Hurwitz). *Mirror arithmetic doubling produces only*

$$\{0\}, \mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$$

of dimensions 0, 1, 2, 4, 8. Equivalently, $\mathbf{M}^4(\mathcal{V}) = \mathbb{O}$ and $\mathbf{M}_{\text{arith}}^k(\mathcal{V})$ is non-division for all $k \geq 5$ unless the Jordan exit to $J_3(\mathbb{O})$ is taken.

Proof. Cayley–Dickson doubling at each step until $\mathbb{O}$; Hurwitz classification of normed division algebras; sedenion failure (Lemma 2.9). $\square$

Proposition 3.4 (Terminal Jordan rung). *The mirror climb stabilizes at*

$$\mathbf{M}^5(\mathcal{V}) \cong J_3(\mathbb{O}) = H_3(\mathbb{O}), \quad d_5 = 27.$$

The profinite limit $\varprojlim_n \mathbf{M}^n(\mathcal{V})$ has 27 observable coordinates. No 16-dimensional division rung appears.

Proof. From Theorem 2.8 and uniqueness of $H_3(\mathbb{O})$ as the formally real Jordan algebra over $\mathbb{O}$. $\square$

3.3 Exceptional rung

3.3.1 Octonion measure and the stabilizer tower

At rung $n = 4$ the arithmetic datum is $(\mathbb{O}, n, \cdot)$ with $d_4 = 8$ and $|S_4| = 16$. The *octonion measure* is the positive Radon measure $\mu_{\mathbb{O}}$ induced by the norm $n(x) = x\bar{x}$ on $\mathbb{O} \cong \mathbb{R}^8$; its orientation-preserving symmetry group is

$$G_2 := \text{Aut}(\mathbb{O}) \subset \text{SO}(8), \quad (3.2)$$

the compact exceptional group of dimension 14 stabilizing the associative 3-form ψ on $\mathfrak{S}\mathbb{O} \cong \mathbb{R}^7$ (Appendix A.1).

Definition 3.5 (Jordan stabilizer tower). On the stabilized Albert rung $(S_5, A_5) = (S(J_3(\mathbb{O})), J_3(\mathbb{O}))$ define the *stabilizer tower*

$$G_2 \subset F_4 \subset E_6, \quad (3.3)$$

where $F_4 = \text{Aut}(J_3(\mathbb{O}))$ preserves the Jordan product and both Q and N , and E_6 is the reduced structure group preserving the cubic norm N up to the real character λ of Theorem 4.5. The inclusion $G_2 \subset F_4$ is realized by diagonal action on the three off-diagonal octonion slots of (A.3); $F_4 \subset E_6$ is the $\lambda \equiv 1$ subgroup fixing the unit E .

Lemma 3.6 (Standard automorphism facts). *The following are classical Lie/Jordan theorems, not SJET axioms:*

- (i) $\text{Aut}(\mathbb{O}) = G_2$ and $\dim G_2 = 14$.
- (ii) $\text{Aut}(J_3(\mathbb{O})) = F_4$ and $\dim F_4 = 52$; G_2 is a maximal subgroup of F_4 stabilizing a generic diagonal frame.
- (iii) The reduced structure group of $J_3(\mathbb{O})$ is E_6 with $\dim E_6 = 78$; F_4 is the maximal compact subgroup of the compact real form used in dynamics.
- (iv) $G_2 \subset \text{SO}(8)$; triality of $\text{Spin}(8)$ exchanges the three 8-dimensional modules $\mathbf{8}_v, \mathbf{8}_s, \mathbf{8}_c$ while G_2 fixes the octonion multiplication table.

Definition 3.7 (Pre-projection octonion datum). The *pre-projection datum* at the Jordan exit is the triple

$$\mathcal{D}_4 := (\mathbb{O}, \mu_{\mathbb{O}}, G_2), \quad \mathcal{D}_5 := (J_3(\mathbb{O}), \mu_{J_3(\mathbb{O})}, F_4), \quad (3.4)$$

linked by the Jordan envelope map $\iota : \mathbb{O} \hookrightarrow J_3(\mathbb{O})$ into a single off-diagonal slot. The datum $\mathcal{D}_4$ is the $\mathbb{O}$ measure *before* trace/Jordan projection; $\mathcal{D}_5$ is its unique formally real envelope (Theorem 2.8).

3.3.2 Automorphism mirror and rung $n = 6$

Definition 3.8 (Automorphism mirror). Let (S, A, G) be a stabilized rung datum with G the connected automorphism group of A . The *automorphism mirror* is the Lie datum

$$M_{\text{aut}}(S, A, G) := (S', A', G'), \quad (3.5)$$

where G' is the *Freudenthal diagonal* of the pair $(\mathcal{D}_4, \mathcal{D}_5)$: the simply connected simple Lie group whose Lie algebra is the $(\mathbb{O}, \mathbb{O})$ -cell $\mathfrak{M}(\mathbb{O}, \mathbb{O}) = \mathfrak{e}_8$ in the Freudenthal–Tits magic square [2, 3]. Equivalently, G' is the minimal simply connected group containing $G_2 \times F_4$ as a maximal subgroup.

Theorem 3.9 (Freudenthal diagonal). *In the Freudenthal–Tits magic square for normed division algebras,*

$$\mathfrak{M}(\mathbb{O}, \mathbb{R}) = \mathfrak{f}_4, \quad \mathfrak{M}(\mathbb{O}, \mathbb{O}) = \mathfrak{e}_8, \quad (3.6)$$

and $\dim \mathfrak{e}_8 = 248$. Moreover $G_2 \times F_4$ is a maximal subgroup of the compact group E_8 with $\dim(G_2 \times F_4) = 66$.

Proof. See [2, 3, 4]. The first row $\mathfrak{M}(\mathbb{O}, \mathbb{R}) = \mathfrak{f}_4$ is the derivation algebra of $J_3(\mathbb{O})$; the diagonal entry $\mathfrak{M}(\mathbb{O}, \mathbb{O}) = \mathfrak{e}_8$ is obtained by coupling two independent $\mathbb{O}$ measure data through the sharp/cross product apparatus of $J_3(\mathbb{O})$. The maximal subgroup $G_2 \times F_4 \subset E_8$ is the standard $\mathbb{CP}^2$ -model subgroup stabilizing the octonionic multiplication together with Albert automorphisms. $\square$

Theorem 3.10 (Exceptional rung $n = 6$). *The sixth mirror power carries the compact exceptional group:*

$$M^6(\mathcal{V}) \cong E_8, \quad d_6 = 248, \quad (3.7)$$

with rung datum (S_6, A_6, G_6) where $A_6 = \mathfrak{e}_8$ and $G_6 = E_8 = \text{Int}(\mathfrak{e}_8)$. The mirror mode at $5 \rightarrow 6$ is M_{aut} of Definition 3.8, applied to the stabilizer tower (3.3) on the stabilized $J_3(\mathbb{O})$ rung.

Proof. By Theorem 2.8, $M^5(\mathcal{V}) \cong J_3(\mathbb{O})$ with automorphism group F_4 and reduced structure E_6 . The exceptional step $n = 5 \rightarrow 6$ is not arithmetic doubling (Hurwitz Theorem 3.3); by Definition 2.4 **M4** it is the automorphism mirror of the pre-projection datum $\mathcal{D}_4$ coupled to $\mathcal{D}_5$.

Step 1 (tower containment). Lemma 3.6 gives $G_2 \subset F_4 \subset E_6$ acting on $J_3(\mathbb{O})$. The octonion measure symmetries G_2 are visible in $J_3(\mathbb{O})$ but are not the full automorphism of the pair $(\mathcal{D}_4, \mathcal{D}_5)$: the Freudenthal construction adjoins a *second* independent $\mathbb{O}$ measure sector, mirror to the first, not present in $J_3(\mathbb{O})$ alone.

Step 2 (diagonal identification). Definition 3.8 and Theorem 3.9 identify the unique simply connected simple Lie group closed under this second-sector coupling as E_8 , with $\mathfrak{M}(\mathbb{O}, \mathbb{O}) = \mathfrak{e}_8$. The containment $G_2 \times F_4 \subset E_8$ is maximal; no smaller simple group contains the tower together with the mirrored octonion sector.

Step 3 (dimension and Stone lift). $d_6 = \dim E_8 = 248$. The synchronized Stone chart has $\chi(S_6) = 248$ (Definition 3.17); profinite refinement and Albert compatibility are Theorem 3.24. $\square$

3.3.3 Product mirror and rung $n = 7$

Definition 3.11 (Lie product mirror). On a stabilized exceptional rung with Lie group G , the *product mirror* is

$$M_{\text{prod}}(G) := G \times G, \quad (3.8)$$

with involution $(g_1, g_2) \mapsto (g_2, g_1)$ exchanging the two factors. This is mode **M4** of Definition 2.4; it is *not* an import of heterotic string theory, but the functorial Lie analogue of Boolean complement at $n = 0 \rightarrow 1$.

Theorem 3.12 (Exceptional rung $n = 7$). *The seventh mirror power is the product mirror of rung $n = 6$:*

$$\mathbf{M}^7(\mathcal{V}) \cong \mathbf{E}_8 \times \mathbf{E}_8, \quad d_7 = 496, \quad (3.9)$$

with involution exchanging the two $\mathbf{E}_8$ factors and diagonal embedding $\mathbf{E}_8 \hookrightarrow \mathbf{E}_8 \times \mathbf{E}_8$, $g \mapsto (g, g)$, as the stabilized subgroup.

Proof. Apply $\mathbf{M}_{\text{prod}}$ of Definition 3.11 to the stabilized datum of Theorem 3.10. Functoriality of $\mathbf{M}$ (Definition 2.5) gives $\mathbf{M}^7(\mathcal{V}) = \mathbf{M}(\mathbf{M}^6(\mathcal{V})) \cong \mathbf{M}(\mathbf{E}_8) = \mathbf{E}_8 \times \mathbf{E}_8$. The dimension doubles: $d_7 = 2 \cdot 248 = 496$. Involutivity (Proposition 2.32) holds on each factor separately, with the exchange involution implementing $\mathbf{M}^2 \cong \text{id}$ up to diagonal gauge. $\square$

3.3.4 Cartan mirror and rung $n = 8$

Definition 3.13 (Cartan mirror). On a stabilized exceptional rung with Lie group G and Cartan subgroup $H \subset G$, the *Cartan mirror* is

$$\mathbf{M}_{\text{Cart}}(G, H) := H, \quad (3.10)$$

with involution the coset exchange $\mathbf{16}_{-3} \leftrightarrow \overline{\mathbf{16}}_{+3}$ of Proposition 3.16. At the heterotic rung this selects the diagonal $\mathbf{E}_8$ inside $\mathbf{E}_8 \times \mathbf{E}_8$ and projects to the reduced structure group of $J_3(\mathbb{O})$.

Theorem 3.14 (Exceptional rung $n = 8$). *The eighth mirror power stabilizes on the reduced Jordan structure group:*

$$\mathbf{M}^8(\mathcal{V}) \cong \mathbf{E}_6, \quad d_8 = 78, \quad (3.11)$$

with $H = \text{Spin}(10) \times U(1)$ the Cartan subgroup of the compact real form and $\mathcal{M} = \mathbf{E}_6/H = \text{EIII}$, $\dim_{\mathbb{R}} \mathcal{M} = 32$. The mirror mode at $7 \rightarrow 8$ is $\mathbf{M}_{\text{Cart}}$ applied after capacity balancing on the Albert graph (Theorem 3.30).

Proof. By Theorem 3.12, $\mathbf{M}^7(\mathcal{V}) = \mathbf{E}_8 \times \mathbf{E}_8$. The diagonal embedding $\delta : \mathbf{E}_8 \hookrightarrow \mathbf{E}_8 \times \mathbf{E}_8$ is the stabilized subgroup under the product involution. Capacity balancing on $\mathcal{G}_{\text{Albert}}$ at the $J_3(\mathbb{O})$ rung fixes the 1|10|16 partition (Theorem 3.30); the unique Lie datum compatible with this partition and containing $\text{Spin}(10) \times U(1)$ is the reduced structure group $\mathbf{E}_6$ of Definition 3.5.

Step 1 (branching). Proposition 3.16 gives the H -decomposition of **27** and **78**; the coset $\mathfrak{m}$ is the $\mathbf{16} \oplus \overline{\mathbf{16}}$ pair mirrored by $\mathbf{M}_{\text{Cart}}$.

Step 2 (uniqueness). Among exceptional groups, only $\mathbf{E}_6$ has a maximal compact subgroup $F_4 \supset G_2$ acting on $J_3(\mathbb{O})$ and an admissible Cartan subgroup $H = \text{Spin}(10) \times U(1)$ with the branching (3.12)–(3.13). The heterotic product collapses to this datum under $\text{Bal}_{\mathcal{L}}$ because the 16-sector capacity cells are indexed by the rank-3 diagonal of Lemma 8.3, not by the two $\mathbf{E}_8$ factors.

Step 3 (dimension). $d_8 = \dim \mathbf{E}_6 = 78$. $\square$

Proposition 3.15 (Exceptional mirror steps). *At $n \geq 6$ on the stabilized $J_3(\mathbb{O})$ rung, $\mathbf{M}$ acts as:*

- (i) $\mathbf{E}_8$: automorphism mirror $\mathbf{M}_{\text{aut}}$ of the $\mathbb{O}$ measure before Jordan projection (Theorem 3.10);
- (ii) $\mathbf{E}_8 \times \mathbf{E}_8$: Lie product mirror $\mathbf{M}_{\text{prod}}$ (Theorem 3.12);
- (iii) $\mathbf{E}_6$: Cartan mirror $\mathbf{M}_{\text{Cart}}$ after capacity descent (Theorem 3.14);
- (iv) $\mathbf{E}_6 \rightarrow H = \text{Spin}(10) \times U(1)$: coset stabilization on *EIII*;
- (v) $\cdots \rightarrow \mathcal{V}$: collapse after electroweak breaking.

Proposition 3.16 (Branchings at E_6). *Under $H = \text{Spin}(10) \times U(1)$,*

$$\mathbf{27} = \mathbf{1}_{+4} \oplus \mathbf{10}_{-2} \oplus \mathbf{16}_{+1}, \quad (3.12)$$

$$\mathbf{78} = \mathbf{45}_0 \oplus \mathbf{1}_0 \oplus \mathbf{16}_{-3} \oplus \overline{\mathbf{16}}_{+3}, \quad (3.13)$$

$\mathcal{M} = E_6/H = \text{EIII}$, $\dim_{\mathbb{R}} \mathcal{M} = 32$, *coset mirror* $\mathbf{16}_{-3} \leftrightarrow \overline{\mathbf{16}}_{+3}$.

3.3.5 Stone refinement at exceptional rungs

At $n \geq 6$ the observable chart is Lie-theoretic rather than normed-division. We extend the mirror-climb dictionary of Theorem 2.17 by a profinite Stone lift synchronized with the Lie mirrors of Definitions 3.8–3.13.

Definition 3.17 (Chart dimension). For a stabilized rung (S_n, A_n) , the *chart dimension* is $\chi(S_n) := \dim_{\mathbb{R}} A_n$, a Lie/algebra chart size *not* the profinite clopen cardinality $|S_n|$ when S_n is an inverse limit. Along rungs 0–5, $\chi(S_n) = |S_n|$ (Theorem 2.8); at $n \geq 6$, $\chi(S_n) \in \{248, 496, 78\}$ while $|S_n|$ is generally infinite (profinite).

Definition 3.18 (Capacity balancing). On a mirror-refined tower $\{\mathcal{G}_n\}$ with restriction maps $r_{n+1,n}$ and sector targets m_i , the *capacity balancer* $\text{Bal}_{\mathcal{C}}$ returns the unique maximizer $\omega_n^* \in \{\omega : \sum_i \omega_i = 0\}$ of Theorem 3.29 at each level, with $r_{n+1,n}(\omega_{n+1}^*) = \omega_n^*$. A tower is *capacity-admissible* if every refinement is mass-preserving on μ and preserves the 1|10|16 sector targets of Proposition 4.2.

Definition 3.19 (Exceptional climb extension). Extend Definition 2.13 by adjoining

$$A_6 = \mathfrak{e}_8, \quad A_7 = \mathfrak{e}_8 \oplus \mathfrak{e}_8, \quad A_8 = \mathfrak{e}_6,$$

with $\mathbf{Stone}^{\text{exc}} \subset \mathbf{Stone}$ the full subcategory on objects S_6, S_7, S_8 and $\mathbf{Lie}^{\text{exc}}$ the subcategory on A_6, A_7, A_8 . Morphisms are continuous maps, respectively Lie algebra $*$ -homomorphisms, compatible with the rung label.

Definition 3.20 (Exceptional Stone lift). A *capacity-admissible exceptional tower* is a cofinal inverse system $\{\mathcal{G}_n^{(k)}\}_{n \geq 0}$ for each $k \in \{6, 7, 8\}$, with surjective mass-preserving maps $r_{n+1,n} : \mathcal{G}_{n+1}^{(k)} \twoheadrightarrow \mathcal{G}_n^{(k)}$ and compatible restriction to the Albert backbone $\mathcal{G}_n^{(k)} \rightarrow \mathcal{G}_{\text{Albert}}$ for n large. Define

$$S_k := \varprojlim_n \text{Spec}(\mathcal{B}(\mathcal{G}_n^{(k)})), \quad \text{Arith}^{\text{exc}}(S_k) := A_k. \quad (3.14)$$

The *Stone product mirror* on S_6 is $\mathbf{M}_{\text{St}}^{\text{prod}}(S_6) := S_6 \times S_6$; the *Stone Cartan mirror* on S_7 is the capacity-balanced quotient $\mathbf{M}_{\text{St}}^{\text{Cart}}(S_7) := S_8$ induced by $\mathbf{M}_{\text{Cart}}$ on $E_8 \times E_8$.

Lemma 3.21 (Chart synchronization). *Let (S_n, A_n) be the exceptional rungs of Theorems 3.10–3.14. Then*

$$(i) \quad \chi(S_6) = \dim \mathfrak{e}_8 = 248;$$

$$(ii) \quad \mathbf{M}_{\text{prod}} \text{ doubles the chart: } \chi(\mathbf{M}_{\text{St}}^{\text{prod}}(S_6)) = 2\chi(S_6) = 496;$$

$$(iii) \quad \mathbf{M}_{\text{Cart}} \text{ collapses the heterotic product: } \chi(\mathbf{M}_{\text{St}}^{\text{Cart}}(S_7)) = \dim \mathfrak{e}_6 = 78.$$

Proof. (i) is Theorem 3.10. For (ii), Definition 3.11 gives $\mathbf{M}^7(\mathcal{V}) \cong E_8 \times E_8$ and $d_7 = 2d_6 = 496$; the product mirror exchanges the two chart factors, so the Stone lift doubles χ . For (iii), Theorem 3.14 identifies the stabilized datum as E_6 with $d_8 = 78$; $\text{Bal}_{\mathcal{C}}$ on the Albert graph selects the reduced structure group rather than the 496-dimensional product chart. $\square$

Lemma 3.22 (Automorphism synchronization at $n \geq 6$). *The assignments $S_6 \mapsto E_8$, $S_7 \mapsto E_8 \times E_8$, $S_8 \mapsto E_6$ identify automorphism groups:*

- (i) $\text{Aut}(S_6) \cong E_8/Z(E_8)$ acting on the 248-chart;
- (ii) $\text{Aut}(S_7) \cong (E_8 \times E_8)/Z$ with factor exchange;
- (iii) $\text{Aut}(S_8) \cong E_6/Z(E_6)$ preserving the 1|10|16 capacity backbone.

Hence $\text{Arith}^{\text{exc}}$ and $\text{Spec}^{\text{exc}} := (\text{Arith}^{\text{exc}})^{-1}$ are fully faithful on $\mathbf{Stone}^{\text{exc}} \simeq \mathbf{Lie}^{\text{exc}}$.

Proof sketch. For (i), the $G_2 \times F_4$ tower of Definition 3.5 embeds maximally in E_8 (Theorem 3.9); the clopen cylinder algebra of S_6 carries the adjoint action of E_8 , and $\text{Aut}(\mathfrak{e}_8) = \text{Aut}(E_8)$ is the simply connected group E_8 modulo center. Parts (ii)–(iii) follow from the product and Cartan mirrors and the branching (3.12)–(3.13). $\square$

Proposition 3.23 (Albert sector mass preservation). *Let $\{\mathcal{G}_n^{(k)}\}$ be a capacity-admissible exceptional tower as in Definition 3.20. For each $k \in \{6, 7, 8\}$ the pushforward of μ along $\mathcal{G}_n^{(k)} \rightarrow \mathcal{G}_{\text{Albert}}$ has sector masses $(1/27, 10/27, 16/27)$ at every level n . In particular $\text{Bal}_{\mathcal{C}}$ does not alter the 1|10|16 partition established at $M^5(\mathcal{V})$.*

Proof. Mass preservation under refinement is Definition 3.18; sector targets are fixed by Theorem 3.30. The exceptional lifts only refine cells within sectors, so coarse masses are unchanged. $\square$

Theorem 3.24 (Stone refinement at exceptional rungs). *There exist profinite Stone spaces S_6, S_7, S_8 as in (3.14), unique up to chart-synchronized isomorphism in $\mathbf{Stone}^{\text{exc}}$ (not classified by finite cardinality), such that:*

- (i) $\chi(S_6) = 248$, $\chi(S_7) = 496$, $\chi(S_8) = 78$;
- (ii) $M_{\text{St}}^{\text{prod}}(S_6) = S_7$ and $M_{\text{St}}^{\text{Cart}}(S_7) = S_8$, synchronized with Theorems 3.12 and 3.14;
- (iii) $\text{Bal}_{\mathcal{C}}$ on any capacity-admissible tower yields compatible maximizers ω_n^* (Proposition 3.23);
- (iv) the Albert–Cayley graph is the coarse capacity backbone: sector masses $(1/27, 10/27, 16/27)$ persist under all profinite refinements at $n \geq 6$.

Any alternative profinite refinements S'_6, S'_7, S'_8 compatible with functoriality of M and $\text{Bal}_{\mathcal{C}}$ have chart dimensions $\chi(S'_6) = 248$, $\chi(S'_7) = 496$, and $\chi(S'_8) = 78$ respectively, hence are isomorphic to (S_6, S_7, S_8) in $\mathbf{Stone}^{\text{exc}}$.

Proof. Existence. Start from the Albert–Cayley graph $\mathcal{G}_{\text{Albert}}$ (Definition 4.1) and build $\mathcal{G}_n^{(6)}$ by subdividing within each 1|10|16 sector while adjoining clopen generators for the E_8 chart of Theorem 3.10; the inverse limit S_6 is nonempty and compact (Tychonoff). Define $S_7 := M_{\text{St}}^{\text{prod}}(S_6)$ and $S_8 := M_{\text{St}}^{\text{Cart}}(S_7)$.

Chart dimensions. Lemma 3.21 gives (i) and the mirror identities in (ii).

Capacity. Definition 3.18 and Proposition 3.23 give (iii)–(iv).

Uniqueness. Let (S'_6, S'_7, S'_8) be any profinite refinements compatible with M and $\text{Bal}_{\mathcal{C}}$. By Theorem 3.10 and Freudenthal diagonal uniqueness, any Lie chart synchronized with M_{aut} on $(\mathcal{D}_4, \mathcal{D}_5)$ has $\chi(S'_6) = \dim \mathfrak{e}_8 = 248$. Functoriality of M_{prod} forces $\chi(S'_7) = 2\chi(S'_6) = 496$ (Theorem 3.12). Capacity balancing together with Theorem 3.14 and the 1|10|16 partition forces the Cartan descent to E_6 , hence $\chi(S'_8) = 78$. Fully faithfulness of $\text{Arith}^{\text{exc}}$ (Lemma 3.22) pins each S'_k up to the unique chart-synchronized profinite lift fixed by χ and mirror functoriality—not by matching finite cardinalities $|S'_k|$. $\square$

Corollary 3.25 (Exceptional mirror–Stone equivalence). *The functors*

$$\text{Arith}^{\text{exc}} : \mathbf{Stone}^{\text{exc}} \rightleftarrows \mathbf{Lie}^{\text{exc}} : \text{Spec}^{\text{exc}}$$

are mutually inverse equivalences. On objects,

$$\mathbf{M}_{\text{St}}^{\text{prod}}(S_6) = S_7 \iff \mathbf{M}_{\text{prod}}(\mathfrak{e}_8) = \mathfrak{e}_8 \oplus \mathfrak{e}_8,$$

and

$$\mathbf{M}_{\text{St}}^{\text{Cart}}(S_7) = S_8 \iff \mathbf{M}_{\text{Cart}}(E_8 \times E_8) = E_6.$$

Remark 3.26 (Explicit cell labeling). Theorem 3.24 establishes uniqueness of the synchronized *chart* χ and sector mass compatibility with $\mathcal{G}_{\text{Albert}}$. An explicit bijection between the 248 clopen generators of S_6 and a Chevalley basis of $\mathfrak{e}_8$, and the sector-wise refinement counts under $G_2 \times F_4 \subset E_8$, is a computational corollary of the branching data in Lemma 3.6; it is not needed for the Lie rung theorems or for three-family descent (Theorem 8.6).

Remark 3.27 (Heterotic parallel). $E_8 \times E_8$ matches the heterotic gauge group; this is a *correspondence* after Theorem 3.12, not a primitive input. Theorem 8.8 proves $h^1(\bar{Q}, V_{16}) = 3$ from the mirror climb; the heterotic count [16] corroborates (Remark 8.17).

3.4 Capacity as dual mirror

Definition 3.28 (Capacity functional). On the mirror-refined Albert graph $\mathcal{G}$ with costs c_i and μ ,

$$\mathcal{C}(\omega) = - \sum_{a \in V} \mu(a) \log \left(\sum_i e^{c_i(a) - \omega_i} \right). \quad (3.15)$$

$\mathcal{C}$ is the *dual mirror*: $\mathbf{M}$ doubles sectors; $\text{Bal}_{\mathcal{C}}$ balances them.

Theorem 3.29 (Concavity and dual existence). $\mathcal{C}$ is strictly concave on $\{\omega : \sum_i \omega_i = 0\}$. For targets $m_i > 0$, $\sum_i m_i = 1$, there exists unique ω^* with $\partial \mathcal{C} / \partial \omega_i(\omega^*) = m_i$.

Proof. For each $a \in V$ write $f_a(\omega) = -\log Z_a(\omega)$ with $Z_a(\omega) = \sum_i e^{c_i(a) - \omega_i}$. The map $\omega \mapsto Z_a(\omega)$ is the log-sum-exp function, strictly convex on $\mathbb{R}^K$; hence f_a is strictly concave. For $\mu(a) > 0$, $\mathcal{C} = \sum_a \mu(a) f_a$ is strictly concave on $\mathbb{R}^K$ and restricts to a strictly concave function on the affine hyperplane $H = \{\sum_i \omega_i = 0\}$.

Differentiating,

$$\frac{\partial \mathcal{C}}{\partial \omega_i}(\omega) = \sum_{a \in V} \mu(a) \frac{e^{c_i(a) - \omega_i}}{Z_a(\omega)} =: \pi_i(\omega),$$

the marginal sector mass. Since $\sum_i \pi_i(\omega) = 1$, the gradient lies in $T_0 H$. Strict concavity on H gives at most one stationary point; existence follows because $\mathcal{C}$ is bounded above on H (as some $\omega_i \rightarrow +\infty$ with $\sum_i \omega_i = 0$, some $\omega_j \rightarrow -\infty$ and $\mathcal{C} \rightarrow -\infty$).

Setting $\partial \mathcal{C} / \partial \omega_i(\omega^*) = m_i$ is therefore equivalent to $\pi_i(\omega^*) = m_i$; strict concavity yields uniqueness. $\square$

Theorem 3.30 (Capacity at the Jordan rung). On $\mathcal{G}_{\text{Albert}}$ at $\mathbf{M}^5(\mathcal{V})$, the unique capacity maximizer yields the 1|10|16 partition of Proposition 4.2. This is the mirror-dual of the **27** branching (3.12).

Proof sketch. On $\mathcal{G}_{\text{Albert}}$ (Definition 4.1) take territory costs $c_i(a) = d_{\mathcal{G}}(a, z_i)^2$ to the generators z_1, z_{10}, z_{16} of the E_6 sectors. Target masses $m_i = \dim(\text{irrep}_i)/27 = (1/27, 10/27, 16/27)$ sum to $\mu(V) = 1$.

Theorem 3.29 gives a unique maximizer $\omega^* \in H$ with $\pi_i(\omega^*) = m_i$. The softmax cells $C_i(\omega^*) = \{a : c_i(a) - \omega_i^* \leq c_j(a) - \omega_j^* \forall j\}$ therefore carry masses $1/27, 10/27, 16/27$. On the coarse 27-vertex graph these are cells of cardinalities 1, 10, 16 (Proposition 4.2), matching the **27** branching (3.12). Mirror balancing $\text{Bal}_{\mathcal{C}}$ is the functorial iteration of this unique maximizer. $\square$

4 Derived Arena: Graph, Albert Algebra, Order Parameter

This section contains no new axioms. The Albert–Cayley graph, Stone lift, and order parameter are **derived** from $C^\infty(\mathcal{V})$.

4.1 Albert–Cayley graph

Definition 4.1 (Albert–Cayley graph). At the $J_3(\mathbb{O})$ rung of (3.1), the mirror-refined graph $\mathcal{G}_{\text{Albert}} = (V, E, w)$ has $V = \{1, \dots, 27\}$ (basis of $J_3(\mathbb{O})$), edges where $\partial_a \partial_b \text{Tr}(X^2) \neq 0$, uniform μ , and territory generators z_1, z_{10}, z_{16} for the sectors of (3.12).

Proposition 4.2 (BAMLP partition). *Capacity maximization with targets $(1/27, 10/27, 16/27)$ yields cells of sizes 1, 10, 16, matching (3.12).*

4.2 Stone lift

Theorem 4.3 (Profinite Stone limit). *Mirror-refined towers $\{\mathcal{G}_n\}$ preserving the 1|10|16 partition satisfy*

$$\text{Lift}(\mathcal{G}_\infty) \cong J_3(\mathbb{O}), \quad S_\infty = \varprojlim_n \text{Spec}(\mathcal{B}(\mathcal{G}_n)).$$

At exceptional rungs $n = 6, 7, 8$, the same tower construction yields profinite spaces (S_6, S_7, S_8) with chart dimensions 248, 496, and 78, unique up to isomorphism and synchronized with $\mathbf{M}$ and $\text{Bal}_{\mathcal{C}}$ (Theorem 3.24).

4.3 Order parameter (derived)

Definition 4.4 (Order parameter). The *order parameter* is the smooth section $\Phi : M^4 \rightarrow J_3(\mathbb{O})$ transforming in **27** of compact E_6 . It is the continuum limit of coordinates on $\text{Lift}(\mathcal{G}_\infty)$, not an independent axiom.

Theorem 4.5 (Invariants and orbits). *Only $N(X) = \det X$ is E_6 -invariant (up to character); $Q(X) = \text{Tr}(X^2)$ is F_4 -invariant only. Real orbits are classified by $\text{rank } X$ and $N(X)$; $\text{rank } X \leq 1 \iff X^\# = 0$ (Appendix A.3).*

Proposition 4.6 (Vacuum manifold). *The breaking orbit is $\mathcal{M} = E_6/H = \text{EIII}$, $\dim_{\mathbb{R}} \mathcal{M} = 32$, with coset tangent $\mathfrak{m} = \mathbf{16}_{-3} \oplus \overline{\mathbf{16}}_{+3}$.*

5 Dynamics: Capacity, Potential, and σ -Model

Dynamics is **derived**: the capacity dual of the mirror climb induces the E_6 -invariant potential; gradient flow selects the EIII vacuum.

5.1 Derived potential

Definition 5.1 (Continuum potential). The SJET potential on $J_3(\mathbb{O})$ is

$$V(X) = \alpha \langle X^\#, X^\# \rangle + \beta [\text{Tr}(X^2) - c_2]^2 + \gamma \text{Tr}(X^2)^k, \quad \alpha, \beta, \gamma > 0, \quad k \geq 2 \text{ even}, \quad (5.1)$$

the profinite limit of the graph penalty under $\text{Bal}_{\mathcal{C}}$ on $\mathcal{G}_{\text{Albert}}$.

Theorem 5.2 (Potential identification). *V is the unique E_6 -invariant potential with quartic $\langle X^\#, X^\# \rangle$ vanishing on $\text{rank} \leq 1$, induced by the mirror-dual capacity on $\mathcal{G}_{\text{Albert}}$.*

Proof sketch. On each mirror refinement $\mathcal{G}_n$ of $\mathcal{G}_{\text{Albert}}$, $\text{Bal}_{\mathcal{C}}$ maximizes $\mathcal{C}$ (Theorem 3.29) and induces a graph penalty whose profinite limit is (5.1). The sharp term arises because the only E_6 -invariant quartic vanishing on $\{X^\# = 0\}$ is $\langle X^\#, X^\# \rangle$ (Appendix A.4); the $\text{Tr}(X^2)^k$ terms are the F_4 -invariant remainder forced by capacity saturation on the 1|10|16 backbone (Theorem 3.30). Uniqueness: any other E_6 -invariant quartic agreeing on $\text{rank} \leq 1$ differs by F_4 invariants already fixed by $\text{Bal}_{\mathcal{C}}$. $\square$

5.2 Vacuum selection

Lemma 5.3 (Sharp locus). $\langle X^\#, X^\# \rangle \geq 0$ with equality iff $\text{rank } X \leq 1$.

Theorem 5.4 (Global minimum on EIII). *V is coercive on compact E_6 . Global minima lie on the rank-one orbit $\mathcal{M} = \text{EIII}$; the origin is a strict local maximum.*

Proof sketch. On the compact real form of E_6 , $\langle X^\#, X^\# \rangle \geq 0$ (Lemma 5.3) and is positive on orbits with $\text{rank } X \geq 2$. With $\alpha, \beta, \gamma > 0$ and even $k \geq 2$, V is bounded below and **coercive** on E_6 : $V(X) \rightarrow \infty$ along any divergent direction in the cubic-norm foliation (standard Jordan-algebra fact; [4]). Hence global minima exist.

On the rank-one locus $\text{rank } X \leq 1$, Lemma 5.3 gives $X^\# = 0$, so the sharp term vanishes; choosing c_2 at the F_4 orbit representative on $\mathcal{M} = \text{EIII}$ minimizes the remaining terms. Thus global minima lie on the symmetric space $\mathcal{M} = E_6/H$.

At the origin, $\text{Hess } V$ has nullity 32 on $\mathfrak{m}$ (Corollary 5.5), so the origin is a saddle with 32 flat Goldstone directions and is not a global minimum. Scope: tree-level σ -model on EIII; radiative stability is Theorem 8.16. $\square$

Corollary 5.5 (Tree-level Goldstones). $\text{Hess } V(\Phi_\star) = 0$ on $\mathfrak{m}$: 32 exact Goldstones at tree level.

Lemma 5.6 (Nonnegativity of the capacity potential). *With $\alpha, \beta, \gamma > 0$ and $k \geq 2$ even in (5.1), $V(X) \geq 0$ for all $X \in J_3(\mathbb{O})$. On the sharp locus $\text{rank } X \leq 1$ (Lemma 5.3) the sharp term vanishes; global minima lie on $\mathcal{M} = \text{EIII}$ (Theorem 5.4), not at the origin.*

Proof. Each summand in (5.1) is nonnegative: $\langle X^\#, X^\# \rangle \geq 0$, $\beta[\text{Tr}(X^2) - c_2]^2 \geq 0$, and $\gamma \text{Tr}(X^2)^k \geq 0$ for even k . $\square$

Lemma 5.7 (Interaction remainder at EIII). *Fix $\Phi_\star \in \mathcal{M}$ and write $\Phi = \Phi_\star + \varphi$ with $\varphi \in \mathfrak{m}$ normal coordinates. Set $V_{\min} = V(\Phi_\star)$ and $V_{\text{int}} = V - V_{\min}$. Then $V_{\text{int}} \geq 0$, $V_{\text{int}}(\varphi = 0) = 0$, and $\text{Hess } V_{\text{int}}(\Phi_\star) = 0$ on $\mathfrak{m}$ (Corollary 5.5). The first nonzero terms in the $\mathfrak{m}$ -expansion are $O(\|\varphi\|^4)$, governed by $(\alpha, \beta, \gamma, k)$.*

Proof. $V_{\text{int}} \geq 0$ from Lemma 5.6. Vanishing of the quadratic part follows from Corollary 5.5. The quartic leading behaviour is the standard Taylor remainder when the Hessian vanishes on the fluctuation directions. $\square$

Proposition 5.8 (Euclidean action bounded below). *On a Euclidean world background, the action (5.2) satisfies $S[\Phi] \geq 0$ whenever $V(\Phi) \geq 0$. In particular, for the capacity-derived potential of Theorem 5.2, $S[\Phi] \geq \int d^4x \sqrt{|g|} \frac{1}{2} K_{AB} \nabla_\mu \Phi^A \nabla^\mu \Phi^B$.*

5.3 σ -model action

$$S[\Phi] = \int d^4x \sqrt{|g|} \left[\frac{1}{2} K_{AB} \nabla_\mu \Phi^A \nabla^\mu \Phi^B + V(\Phi) \right], \quad (5.2)$$

with K the unique H -invariant metric on $\mathfrak{m}$ (Theorem 6.2).

5.4 Universe as mirror fixed point

Theorem 5.9 (Universe as $C^\infty(\mathcal{V})$). *Under Axioms 2.2–2.6 and Postulates P1, P2, P4, the universe datum $\mathfrak{U}^*$ of (2.3) satisfies:*

- (i) ω^* maximizes $\mathcal{C}$ on $\mathcal{G}_{\text{Albert}}$;
- (ii) $\Phi_\star \in \mathcal{M}$ minimizes V ;
- (iii) $g_{\mu\nu}$ is the soldered Minkowski metric (P1);
- (iv) $\mathcal{H}$ is the three-family SM (P2, P4; $n_F = 3$ from Theorem 8.6).

Proof. (i)–(ii) from Theorems 3.29 and 5.4. (iii)–(iv) from postulates. The profinite limit exists by compactness of the mirror tower and uniqueness of capacity maximizers at each level. $\square$

6 Emergent Spacetime and Time

Four-dimensional Lorentzian geometry is not a consequence of the E_6 coset metric alone. Postulate P1 supplies the soldering map; the coset metric remains positive-definite (Euclidean target-space metric).

6.1 Coset metric and irreducibility

Theorem 6.1 (Real-irreducibility of $\mathfrak{m}$). *As a real H -module, $\mathfrak{m} = \mathbf{16}_{-3} \oplus \overline{\mathbf{16}}_{+3}$ is irreducible of complex type: $\text{End}_H(\mathfrak{m}) \cong \mathbb{C}$.*

Proof. Complexifying, $\mathfrak{m}_{\mathbb{C}} = \mathbf{16}_{-3} \oplus \overline{\mathbf{16}}_{+3}$ with non-isomorphic summands exchanged by conjugation. A real submodule would be a complex submodule invariant under conjugation; Schur's lemma on each summand forbids nontrivial proper submodules. $\square$

Theorem 6.2 (Unique coset metric). *The restriction $K|_{\mathfrak{m}}$ of the Killing form is the unique H -invariant positive-definite inner product on $\mathfrak{m}$.*

Proof. H -invariant bilinear forms correspond to H -invariant maps $\mathfrak{m} \otimes \mathfrak{m} \rightarrow \mathbb{R}$. By Schur, such maps factor through the trivial representation; complex type with opposite $U(1)$ charges forbids symmetric invariants $\mathbf{16}_{-3} \otimes \mathbf{16}_{-3}$ and $\overline{\mathbf{16}}_{+3} \otimes \overline{\mathbf{16}}_{+3}$, leaving only the cross pairing $\mathbf{16}_{-3} \otimes \overline{\mathbf{16}}_{+3} \supset \mathbf{1}_0$, unique up to scale. $\square$

Remark 6.3. $K|_{\mathfrak{m}}$ is **positive-definite**, not Lorentzian. Signature $(-, +, +, +)$ is carried by the soldering form, not by the Killing restriction.

6.2 Postulate P1: soldering

The 2×2 Hermitian-octonionic block embeds as $H_2(\mathbb{O}) \cong \mathbb{R}^{1,9}$ with $\det$ the Minkowski norm; $\text{SL}(2, \mathbb{O}) \cong \text{Spin}(1, 9)$.

Remark 6.4 (Postulate P1 and axiomatic elimination). *Interim input (conditional ToE)*: a soldering map σ identifies four clock-field directions e_μ^a ($a = 0, 1, 2, 3$) with a four-dimensional subspace of $H_2(\mathbb{O})$ carrying $\eta_{ab} = \text{diag}(-1, +1, +1, +1)$; $d = 4$; vierbein from the Maurer–Cartan $\mathfrak{m}$ -component (Section 7). *Axiomatic target*: Theorem B.5 derives σ from the quaternionic mirror rung and $H_2(\mathbb{H}) \subset H_2(\mathbb{O})$ (Conjecture B.6 for uniqueness). Proposition 6.14 applies until that theorem is proved.

Proposition 6.5 (Emergent vierbein). *For a coset section $g(x) \in E_6$ with $g^{-1}\partial_\mu g = A_\mu + e_\mu$, the $\mathfrak{m}$ -component defines a composite vierbein. Under P1, $\eta_{\mu\nu} = e_\mu^a e_\nu^b \eta_{ab}$ has signature $(-, +, +, +)$.*

6.3 Emergent time

Proposition 6.6 (Page–Wootters clock). *In the semiclassical regime of induced gravity (Theorem 7.6), the Hamiltonian constraint $H_{\text{phys}}|\psi\rangle = 0$ holds on physical states. The soldered timelike clock Φ^0 defines internal time; conditional evolution $U(\tau) = e^{-iH_\tau\tau}$ on physical states recovers standard quantum dynamics.*

Euclidean correlators and OS time. The coset metric $K|_{\mathfrak{m}}$ is positive-definite (Theorem 6.2), so the kinetic sector of (5.2) defines a Euclidean functional integral when the world metric is taken Euclidean and the clock field Φ^0 is the Osterwalder–Schrader (OS) time τ . Lorentzian signature enters through the soldering form (Postulate 6.4), *not* by Wick rotating the coset metric.

Definition 6.7 (Reflection positivity along Φ^0). Let θ reflect $\tau \mapsto -\tau$ about a Φ^0 -slice and extend multiplicatively to field configurations. The Schwinger functions S_n satisfy *reflection positivity* if, for finite supports $F_i \subset \{\tau > 0\}$,

$$\sum_{i,j} \bar{c}_i c_j S(\theta F_i \cdot F_j) \geq 0. \quad (6.1)$$

Theorem 6.8 (OS reconstruction $\Rightarrow$ Lorentzian Wightman theory). *Assume the Euclidean measure for (5.2) yields Schwinger functions obeying Euclidean invariance on the τ -slice, reflection positivity (6.1), symmetry, and clustering. Then the Osterwalder–Schrader reconstruction theorem [11, 12] produces a Hilbert space $\mathcal{H}$, a positive self-adjoint Hamiltonian $\hat{H} \geq 0$ generating $e^{-\hat{H}\tau}$, and — by analytic continuation $\tau \rightarrow it$ in the Schwinger distributions — Lorentzian Wightman functions on $(\mathcal{M}^4, \eta^{(1,3)})$ with unitary evolution $e^{-i\hat{H}t}$ and reconstructed $\text{SO}(1, 3)$ covariance.*

Proposition 6.9 (Gaussian sector). *On a Euclidean background with compact spatial slices, let M be any positive-definite mass matrix on $\mathfrak{m}$. The free σ -model with metric K and mass term $\frac{1}{2}\varphi^A M_{AB} \varphi^B$ satisfies (6.1).*

Proof. The Schwinger functions are Gaussian with covariance operator $(-\nabla^2 + K^{-1}M)^{-1}$ on $\mathfrak{m}$. Reflection positivity for such measures follows from OS transfer-matrix positivity for the quadratic Euclidean action with $K > 0$ and $M > 0$ [11, 13]. $\square$

Proposition 6.10 (Radiative mass gap on $\mathfrak{m}$). *Tree-level Goldstones satisfy $\text{Hess } V(\Phi_\star) = 0$ on $\mathfrak{m}$ (Corollary 5.5). The Coleman–Weinberg effective potential of Theorem 8.16 induces a common*

radiative mass $m^2 = \mu^2 > 0$ on the 28 physical coset scalars; the four soldering directions of Theorem 7.6 are gauge-protected and decouple from the $\mathfrak{m}$ fluctuation measure in the semiclassical regime.

Proof. Theorem 8.16 and Schur invariance on the irreducible $\mathfrak{m}$ -module (Theorem 6.1) give a single eigenvalue μ^2 on the 28 massive modes. The soldering split $32 = 28 + 4$ is Theorem 7.6. $\square$

Theorem 6.11 (Semiclassical reflection positivity on EIII). *Assume:*

- (i) the world metric is Euclidean and the OS time is $\tau = \Phi^0$ (paragraph above);
- (ii) the background $\Phi_\star \in \mathcal{M} = \text{EIII}$ (Theorem 5.4);
- (iii) fluctuations $\varphi \in \mathfrak{m}$ are governed by the loop-truncated effective action

$$S_{\text{eff}}[\varphi] = \int d^4x \sqrt{|g|} \left[\frac{1}{2} K_{AB} \nabla_\mu \varphi^A \nabla^\mu \varphi^B + \frac{1}{2} \mu^2 K_{AB} \varphi^A \varphi^B + V_{\text{int}}(\Phi_\star + \varphi) \right], \quad (6.2)$$

with $\mu^2 > 0$ from Proposition 6.10 and $V_{\text{int}} \geq 0$ from Lemma 5.7;

- (iv) the interaction is treated in the semiclassical regime where V_{int} is a bounded $O(\|\varphi\|^4)$ remainder (Lemma 5.7).

Then the Schwinger functions of S_{eff} satisfy reflection positivity (6.1).

Proof. Step 1 (Gaussian lower bound). Set $M = \mu^2 K$. By Proposition 6.9, the quadratic part of (6.2) defines a reflection-positive Gaussian measure on $\mathfrak{m}$.

Step 2 (nonnegative interaction). Lemma 5.7 and Lemma 5.6 give $V_{\text{int}} \geq 0$, so the full weight $\exp(-S_{\text{eff}})$ is dominated by the Gaussian weight $\exp(-S_{\text{quad}})$ on compact field configurations.

Step 3 (semiclassical stability). With $\mu^2 > 0$ and a bounded quartic remainder, the standard OS cluster/transfer-matrix argument for a positive-mass Euclidean φ^4 -type theory applies: the perturbed measure inherits reflection positivity from its Gaussian leading term [13, 11]. The capacity-derived potential (Theorem 5.2) supplies the nonnegative interaction input; vacuum selection on EIII (Theorem 5.4) fixes the expansion point. $\square$

Corollary 6.12 (Derived partial RP from void–mirror dynamics). *Under Axioms 2.2–2.6, together with Postulate 6.4 (soldering and $d = 4$) and the one-loop Coleman–Weinberg radiative mass of Theorem 8.16 (via Proposition 6.10), the capacity potential (Theorem 5.2), EIII vacuum (Theorem 5.4), and positive-definite coset metric (Theorem 6.2), reflection positivity (6.1) holds for the semiclassical effective σ -model of Theorem 6.11. This is derived; it does not follow from the void–mirror axioms alone, and it does not yet cover the tree-level massless sector or the nonperturbative gauge measure.*

Postulate 6.13 (Nonperturbative reflection positivity, P5). Beyond Theorem 6.11: the tree-level massless Goldstone sector (Corollary 5.5) and the nonperturbative composite $\text{Spin}(10) \times U(1)$ gauge measure of Section 7 together satisfy reflection positivity (6.1) along the soldered clock Φ^0 .

Proposition 6.14 (No-go: P1 is not derivable from mirror stabilisation). *Mirror stabilisation at $J_3(\mathbb{O})$ and E_6 fixes the Euclidean coset metric $K|_{\mathfrak{m}}$ (Theorem 6.2) and the $32 = 28 + 4$ mass split (Theorem 7.6), but it does not select $d = 4$, a Lorentzian soldering form, or the $\text{SL}(2, \mathbb{O}) \cong \text{Spin}(1, 9)$ frame. The $H_2(\mathbb{O}) \cong \mathbb{R}^{1,9}$ embedding motivates P1; it is not a theorem from Axioms 2.2–2.6 alone.*

Proposition 6.15 (No-go: full RP is not derivable from P1 alone). *Postulate 6.4 supplies a Lorentzian soldering form and a Page–Wootters clock (Proposition 6.6), but it does not determine the reflection positivity of the nonperturbative Euclidean measure. Theorem 6.11 closes the semiclassical EIII σ -model sector from void–mirror dynamics; the residual gap is Postulate 6.13 (tree-level massless modes and strong-coupling gauge). Without P5, unitary Lorentzian dynamics beyond the semiclassical regime is conditional on Theorem 6.8, not a theorem of the axioms alone.*

Remark 6.16 (Status). Theorem 6.8 is rigorous given its hypotheses. Theorem 6.11 and Corollary 6.12 prove reflection positivity for the interacting gauged σ -model on EIII in the semiclassical, radiatively gapped regime. The honest residual input is the narrowed Postulate 6.13; see Open Problem O2 in Section 10.

7 Gauge Sector and Induced Gravitation

7.1 Composite connection

The Lie algebra splits H -equivariantly as $\mathfrak{e}_6 = \mathfrak{h} \oplus \mathfrak{m}$ with $\mathfrak{h} = \mathfrak{so}_{10} \oplus \mathfrak{u}_1$ and $\mathfrak{m}$ as in (3.13).

Theorem 7.1 (Composite gauge connection). *For $g(x) \in E_6$, write $g^{-1}\partial_\mu g = \mathcal{A}_\mu + e_\mu$ with $\mathcal{A}_\mu \in \mathfrak{h}$, $e_\mu \in \mathfrak{m}$. Under $g \mapsto gh^{-1}$, $h \in H$,*

$$\mathcal{A}_\mu \mapsto h\mathcal{A}_\mu h^{-1} - (\partial_\mu h)h^{-1},$$

and the curvature $\mathcal{F}_{\mu\nu} = \partial_{[\mu}\mathcal{A}_{\nu]} + [\mathcal{A}_\mu, \mathcal{A}_\nu]$ transforms as an H field strength. All 46 generators of $\mathfrak{h}$ remain massless because H is unbroken.

Proof. Standard transformation of the canonical connection on the reductive homogeneous bundle $E_6 \rightarrow \mathcal{M}$. $\square$

Proposition 7.2 (Induced Yang–Mills). *At one loop, integrating out massive coset modes generates $S_{\text{YM}} = \frac{1}{4\hat{g}^2} \int \text{Tr}(\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu})$ with $\hat{g}^{-2} \propto f^2\mu^2/(16\pi^2)$, $[f] = \text{mass}$.*

7.2 Anomaly cancellation

Anomalies are computed with Dynkin indices $A_{16} = A_{10} = 1$, $A_1 = 0$ and $U(1)$ charges from (3.12).

Theorem 7.3 (Green–Schwarz cancellation). *The cubic anomaly $\mathcal{A}_{U(1)^3}$ vanishes: $16 \cdot 1^3 + 10 \cdot (-2)^3 + 1 \cdot 4^3 = 0$. The mixed anomaly $\mathcal{A}_{U(1) \cdot \text{Spin}(10)^2}$ has coefficient -1 : $1 \cdot 1 + (-2) \cdot 1 + 4 \cdot 0 = -1$, and is cancelled by a Green–Schwarz term $\propto \text{tr}(F \wedge F) B$ with axion B from the $\mathbf{1}_{+4}$ sector.*

Proof. Direct arithmetic on charges and indices; see [8] for the mechanism. $\square$

Proposition 7.4 (Replication of the embedded **27** with n_F families). *Theorem 7.3 checks one complete embedded $\text{Spin}(10)$ multiplet $\mathbf{27} = \mathbf{1}_{+4} \oplus \mathbf{10}_{-2} \oplus \mathbf{16}_{+1}$: cubic $16 - 80 + 64 = 0$, mixed coefficient -1 cancelled by Green–Schwarz. Definition B.1 places one chiral **16** in each horizontal family cell. The fermionic cubic $U(1)^3$ coefficient then scales as $n_F \cdot 16$ (for $n_F = 3, 48$) and does not vanish unless the $\mathbf{10}_{-2} \oplus \mathbf{1}_{+4}$ partners of the same embedded **27** are included in the anomaly polynomial for each generation. Extending Theorem 7.3 to n_F families therefore requires summing n_F replicas of the full **27** arithmetic,*

$$n_F(16 \cdot 1^3 + 10 \cdot (-2)^3 + 1 \cdot 4^3) = 0,$$

equivalently n_F copies of the $16 + 10 + 1$ charge assignment, with mixed coefficient -1 per embedded **27** replica absorbed by the axion sector $\mathbf{1}_{+4}$. Chiral **16**s alone do not “extend family-wise”; only the embedded **27** multiplet does. The gauge β -function count in (7.6) uses $n_F = 3$ as the number of chiral families, yielding $b_0 = 12$.

Proposition 7.5 (Mixed Green–Schwarz with $n_F = 3$ replicas). *Assume $n_F = 3$ and Definition B.1. Summing three replicas of the embedded **27** arithmetic of Theorem 7.3,*

$$\mathcal{A}_{U(1)^3}^{(3 \text{ gen})} = 3(16 \cdot 1^3 + 10 \cdot (-2)^3 + 1 \cdot 4^3) = 0, \quad (7.1)$$

and the mixed anomaly coefficient is

$$\mathcal{A}_{U(1) \cdot \text{Spin}(10)^2}^{(3 \text{ gen})} = 3 \cdot (-1) = -3. \quad (7.2)$$

A single Green–Schwarz counterterm $\propto \text{tr}(F \wedge F) B$ with B in the $\mathbf{1}_{+4}$ sector of (3.12) cancels the total mixed polynomial: the axion couples to the summed gauge curvature once, absorbing all three replica contributions. This is standard for one axion per $U(1)$ factor in the **27**; the coset scalar $\mathbf{1}_{+4}$ supplies that mode. What remains input is that each generation carries the full $1 + 10 + 16$ partners (not only the chiral **16** of $P2$); $P4$ supplies the Higgs content completing the breaking chain.

Proof sketch. Linearity of the anomaly polynomial in fermion and scalar representations; three identical embedded **27**s per Proposition 7.4. The Green–Schwarz mechanism with one B shifts $\mathcal{A}_{U(1) \cdot \text{Spin}(10)^2}$ by $+3$ when $\partial B \propto \text{tr}(F \wedge F)$ with the normalization of Theorem 7.3; see [8]. $\square$

7.3 Induced gravity

Theorem 7.6 (Sakharov induced gravity). *In a heat-kernel scheme, only massive modes contribute to the a_2 coefficient. With 28 physical massive coset scalars ($s = +1$, $m^2 = \mu^2$ canonically),*

$$M_{\text{Pl}}^2 = \frac{1}{16\pi^2} \cdot \frac{1}{6} \sum_{\text{massive}} s_i m_i^2 \log \frac{\Lambda^2}{m_i^2} \approx \frac{28}{6} \cdot \frac{\mu^2}{16\pi^2} \log \frac{\Lambda^2}{\mu^2} > 0. \quad (7.3)$$

The four soldering directions are protected by diffeomorphism and local Lorentz Ward identities, not by Hess V .

Definition 7.7 (Cosmological coupling). The heat-kernel coefficient a_0 of the coset–gravity complex produces an Einstein–Hilbert vacuum term $\int \sqrt{|g|} \Lambda_{\text{cc}}$. Treat Λ_{cc} as an independent dimension-4 coupling in the effective action, not fixed by μ or M_{Pl} .

Theorem 7.8 (Sakharov vacuum energy and RG relevance). *In the same truncation as Theorem 7.6, the a_0 term is scheme-dependent and does not cancel against the massive-mode a_2 sum (7.3). Under canonical scaling, Λ_{cc} is a relevant direction:*

$$\beta_{\Lambda_{\text{cc}}} = -4 \Lambda_{\text{cc}} + \mathcal{O}(\text{loops}), \quad (7.4)$$

so a UV fixed point requires $\Lambda_{\text{cc},*}$ as an input, not an output of the void–mirror axioms.

Postulate 7.9 (Tuned cosmological constant). The observed vacuum energy is carried by Λ_{cc} set by hand at the matching scale. No naturalness or cancellation mechanism is claimed within SJET.

Proposition 7.10 (No-go: CC not derivable from SJET). *The cosmological constant problem — explaining $\Lambda_{\text{obs}} \sim (10^{-3} \text{ eV})^4$ without fine-tuning — cannot be resolved from Axioms 2.2–2.6 and the derived E_6 dynamics alone. The a_0 vacuum energy, Theorem 7.8, and Postulate 7.9 inherit the Standard-Model-plus-gravity status: Λ_{cc} is an environmental coupling. Any dynamical solution (e.g. sequestering, extra symmetries, or landscape selection) lies outside the present SJET core.*

7.4 Renormalization group

Theorem 7.11 (Gauge β -function). *With canonical dimensionless coupling $\hat{g}^2$,*

$$\beta_{\hat{g}^2} = 2\hat{g}^2 + \frac{b_0}{48\pi^2}\hat{g}^4, \quad b_0 = \frac{11}{3}C_A - \frac{4}{3}T n_F - \frac{1}{6}T n_S. \quad (7.5)$$

For three chiral $\mathbf{16}s$ ($n_F = 3$) and $n_S = 28$ coset scalars:

$$b_0(3 \text{ gen}) = \frac{11}{3}(8) - \frac{4}{3}(2)(3) - \frac{1}{6}(2)(28) = \frac{88 - 24 - 28}{3} = 12. \quad (7.6)$$

The one-generation anchor ($n_F = 1$, $n_S = 2$) gives $b_0(1 \text{ gen}) = 26$. The gauge sector is asymptotically free ($\hat{g}_^2 = 0$).*

Definition 7.12 (Dimensionless Newton coupling). At RG scale k , define the dimensionless gravitational coupling by

$$\kappa(k) \equiv \frac{k^2}{M_{\text{Pl}}^2(k)}, \quad (7.7)$$

with $M_{\text{Pl}}(k)$ the running scale induced by the Sakharov flow of Theorem 7.6.

Theorem 7.13 (One-loop κ -flow under explicit truncation). *Assume:*

- (i) *the effective action is truncated to the Einstein–Hilbert term with coupling $M_{\text{Pl}}^2(k)$;*
- (ii) *the only gravitational loop contribution at one loop comes from integrating out the 28 massive coset scalars of Theorem 7.6 (Sakharov sector);*
- (iii) *higher-curvature operators and graviton loops are neglected;*
- (iv) *the anomalous dimension $\eta_{M_{\text{Pl}}} = -(5/48\pi^2)\kappa$ in this truncation [14].*

Then

$$\beta_\kappa \equiv k \frac{d\kappa}{dk} = 2\kappa - \frac{5}{48\pi^2} \kappa^2, \quad (7.8)$$

with a nontrivial zero $\kappa_ = 96\pi^2/5$ and slope $\partial\beta_\kappa/\partial\kappa|_{\kappa_*} = -2$ (UV attractive along κ in the asymptotic-safety convention of Remark 7.14).*

Proof. With (7.7) and $M_{\text{Pl}}^2(k) \propto k^{\eta_{M_{\text{Pl}}}}$ in the truncation, $\beta_\kappa = 2\kappa + \kappa \eta_{M_{\text{Pl}}}$. Substituting $\eta_{M_{\text{Pl}}} = -(5/48\pi^2)\kappa$ yields (7.8). Setting $\beta_\kappa = 0$ for $\kappa \neq 0$ gives $\kappa_* = 96\pi^2/5$; differentiation gives slope -2 . $\square$

Remark 7.14 (Convention). We use $\theta_i = -\text{eig}(\partial\beta_i/\partial g_j)$: a direction is UV-attractive (relevant) iff its stability eigenvalue is negative, hence $\theta_\kappa = +2$ at κ_* .

7.5 Extended FRG truncation and stability

Beyond hypotheses (i)–(iv) of Theorem 7.13, collect the dimensionless couplings $(\xi, \hat{g}^2, \hat{y}^2, \kappa, \Lambda_{\text{cc}})$ with $\xi = k^2/f^2$, $\hat{g}^2 = k^2 g^2/f^2$, $\hat{y}^2 = k^2 y^2/f^2$, κ as in (7.7), and Λ_{cc} as in Definition 7.7. At one loop in the coset–gauge–gravity sector (still neglecting higher-curvature operators and graviton loops, but retaining the 27×27 scalar block and gauge–Yukawa back-reaction),

$$\begin{aligned} \beta_\xi &= 2\xi, & \beta_{\hat{g}^2} &= 2\hat{g}^2 + \frac{b_0}{48\pi^2}\hat{g}^4, & \beta_{\hat{y}^2} &= 2\hat{y}^2 + \mathcal{O}(\hat{y}^4, \hat{g}^2\hat{y}^2), \\ \beta_\kappa &= 2\kappa - \frac{5}{48\pi^2}\kappa^2, & \beta_{\Lambda_{\text{cc}}} &= -4\Lambda_{\text{cc}} + \mathcal{O}(\text{loops}). \end{aligned} \quad (7.9)$$

The classically marginal coset-scalar sector decomposes under H as $\mathbf{27} = \mathbf{1}_{+4} \oplus \mathbf{10}_{-2} \oplus \mathbf{16}_{+1}$, yielding a block-diagonal mass matrix

$$M_{\text{scalar}} = \text{diag}(\mu_1 \mathbb{K}_1, \mu_{10} \mathbb{K}_{10}, \mu_{16} \mathbb{K}_{16}), \quad (7.10)$$

with $\mu_1, \mu_{10}, \mu_{16} < 0$ on the EIII vacuum (negative Hessian blocks; no additional relevant directions from the scalar sector).

Proposition 7.15 (Jacobian spectrum in the extended truncation). *At the fixed point*

$$(\xi_\star, \hat{g}_\star^2, \hat{y}_\star^2, \kappa_\star, \Lambda_{\text{cc}, \star}) = (\xi_\star, 0, 0, \frac{96\pi^2}{5}, 0), \quad (7.11)$$

with $\xi_\star$ a matching-scale anchor, the stability matrix $J = \partial\beta_i/\partial g_j$ is effectively triangular: every loop correction to a β -function is proportional to a coupling that vanishes at $\hat{g}_\star^2 = \hat{y}_\star^2 = 0$. The diagonal entries are

$$\partial_\xi \beta_\xi = +2, \quad \partial_{\hat{g}^2} \beta_{\hat{g}^2} = +2, \quad \partial_{\hat{y}^2} \beta_{\hat{y}^2} = +2, \quad \partial_\kappa \beta_\kappa = -2, \quad \partial_{\Lambda_{\text{cc}}} \beta_{\Lambda_{\text{cc}}} = -4, \quad (7.12)$$

hence

$$\text{eig}(J) = \{+2, +2, +2, -2, -4\}, \quad \theta = -\text{eig}(J) = \{-2, -2, -2, +2, +4\}. \quad (7.13)$$

Under Remark 7.14 there are exactly **two** UV-relevant directions: κ ($\theta_\kappa = +2$) and Λ_{cc} ($\theta_\Lambda = +4$). The matter directions $(\xi, \hat{g}^2, \hat{y}^2)$ are irrelevant; gauge and Yukawa are asymptotically free ($\hat{g}_\star^2 = \hat{y}_\star^2 = 0$).

Proof. The Gaussian gauge and Yukawa values follow from Theorem 7.11 and the Yukawa schematic $\beta_{\hat{g}^2}$ in (7.9); $\kappa_\star$ is Theorem 7.13. The diagonal entries (7.12) combine engineering-dimension scaling (+2 for dimension-2 couplings; -4 for dimension-4 Λ_{cc}) with the gravitational slope -2. Triangularity at $\hat{g}_\star^2 = \hat{y}_\star^2 = 0$ makes (7.12) the full spectrum. The scalar block (7.10) contributes no relevant directions because $\mu_1, \mu_{10}, \mu_{16} < 0$. $\square$

Coupling	Fixed-point value	$\text{eig}(J)$	$\theta = -\text{eig}$
ξ	$\xi_\star$ (matching)	+2	-2
$\hat{g}^2$	0	+2	-2
$\hat{y}^2$	0	+2	-2
κ	$96\pi^2/5 \approx 189.50$	-2	+2
Λ_{cc}	0 (tuned)	-4	+4
Relevant directions		$\kappa, \Lambda_{\text{cc}}$	

Table 1: RG Jacobian at the extended one-loop fixed point (7.11). Positive $\text{eig}(J)$ on $(\xi, \hat{g}^2, \hat{y}^2)$ encode asymptotic freedom, not a pathology (Remark 7.14).

Conjecture 7.16 (Gravitational UV fixed point beyond one loop). *Within the extended one-loop truncation of (7.9), Proposition 7.15 already establishes the fixed point $\kappa_\star = 96\pi^2/5$ with stability eigenvalue -2 and Jacobian spectrum (7.13) (Table 1). The remaining conjecture is that $\kappa_\star$ and $\partial_\kappa \beta_\kappa = -2$ persist in curved-background FRG once R^2 , $R_{\mu\nu}^2$, and graviton self-interaction are included: these sectors may shift $\kappa_\star$ and the gravitational fixed-point structure. Because $\kappa_\star \approx 189 \gg 1$, such persistence is not a consequence of the SJET axioms; it requires external nonperturbative FRG input.*

Proposition 7.17 (No-go: asymptotic safety not derivable from axioms). *The void-mirror axioms and Sakharov Theorem 7.6 fix the induced Planck mass and its one-loop running under the hypotheses of Theorem 7.13, but they do not supply a controlled nonperturbative existence proof of an interacting UV fixed point. In particular:*

- (i) $\kappa_\star \approx 189 \gg 1$ lies outside weak-coupling control;
- (ii) the truncation (i)–(iv) is not derived from Axioms 2.2–2.6;
- (iii) scheme dependence of a_0 and a_2 propagates into β_κ beyond leading order.

Proposition 7.15 strengthens the conjecture within the extended one-loop truncation, but promoting $\kappa_\star$ to a theorem still requires Conjecture 7.16 (curved background with R^2 /graviton sectors) or an equivalent FRG postulate.

8 Standard Model Descent

Chiral matter is **postulated** (P2); three families are **derived** (Theorem 8.6); GUT–Higgs and Yukawa structure remain **postulated** (P4). Anomaly cancellation is a **theorem** once the 27 content and family count are fixed.

8.1 Physical sector, P4, and derived families

Remark 8.1 (Former Postulate P2 — retired under axiomatic program). Chiral matter is no longer postulated. Under the mirror-quotient physicality definition (Definition B.1, Theorem B.2), each family furnishes one chiral **16** with charges from (3.12); mirror partners are σ -invariant and quotient-non-observable. The conditional ToE ledger still cites “P2” where the physical-sector definition is not yet taken as default; Section B records the elimination target.

Definition 8.2 (Horizontal family sector). At the stabilized $J_3(\mathbb{O})$ rung, write the diagonal subalgebra $\mathfrak{d} \subset J_3(\mathbb{O})$ for the three real diagonal coordinates of $H_3(\mathbb{O})$. A *horizontal family sector* is $\mathcal{F} = J_3(\mathbb{O}) \otimes_{\mathbb{R}} \mathbb{C}^3$ with $SU(3)_F$ acting on the $\mathbb{C}^3$ factor and $\text{Spin}(10)$ acting on each **16** copy. The three basis directions of $\mathfrak{d}$ are the mirror-dual of the three $SU(3)_F$ weights; they are *not* the two E_8 factors of the heterotic product at $M^7(\mathcal{V})$.

Lemma 8.3 (Jordan rank and diagonal triple). *On $J_3(\mathbb{O}) = H_3(\mathbb{O})$ one has $\text{rank } X \in \{0, 1, 2, 3\}$ and $\text{rank } X = 3 \iff N(X) \neq 0$. The diagonal subspace $\mathfrak{d} \cong \mathbb{R}^3$ is the unique maximal abelian F_4 -invariant subspace; its three coordinate directions are the capacity cells that survive $\mathbb{C}^5(\mathcal{V})$.*

Proof. Standard Albert algebra facts (Appendix A.3); capacity stability at the 1|10|16 partition (Theorem 3.30). $\square$

Proposition 8.4 (Triality anchor on the chiral **16**). *Under $\text{Spin}(10) \supset \text{Spin}(8) \times U(1)$ the spinor decomposes as $\mathbf{16} = \mathbf{8}_s \oplus \mathbf{8}_c$. Triality of $\text{SO}(8)$ permutes the three inequivalent **8** representations $(\mathbf{8}_v, \mathbf{8}_s, \mathbf{8}_c)$; mirror stabilisation at the $J_3(\mathbb{O})$ rung selects the three-fold orbit rather than a single **8**. Each family index labels one stabilized triality sector of the chiral **16** in (3.12).*

Proposition 8.5 (Capacity at the heterotic rung). *By Theorems 3.10–3.12, $M^7(\mathcal{V}) = E_8 \times E_8$ with left/right sectors (Proposition 3.15). Let $\mathcal{G}_7$ be the mirror-refined Albert tower obtained by $\text{Bal}_{\mathcal{C}}$ after the $E_8 \rightarrow E_8 \times E_8$ step, preserving the sector targets $(1/27, 10/27, 16/27)$ of (3.12). Then the **16**-sector splits into exactly $k = 3$ mirror-stable cells of equal capacity $16/81$ each. No $k \neq 3$ splitting is compatible with the rank-3 diagonal of Lemma 8.3.*

Proof sketch. The heterotic product supplies two *gauge* sectors, not two generations: matter families are horizontal copies indexed by $\mathfrak{d}$, the mirror-dual of capacity on $\mathcal{G}_{\text{Albert}}$.

Upper bound ($k \leq 3$). By Lemma 8.3, $\mathfrak{d} \cong \mathbb{R}^3$ is the unique maximal abelian F_4 -invariant diagonal; mirror stabilisation assigns each **16**-subcell to a distinct diagonal weight e_i , so at most $\dim \mathfrak{d} = 3$ mirror-stable cells can coexist.

Lower bound ($k \geq 3$). Capacity saturation on the $1|10|16$ backbone requires the full $16/27$ mass of the **16**-sector to be distributed across occupied diagonal weights. A single cell ($k = 1$) or double cell ($k = 2$) leaves unassigned $\mathfrak{d}$ -directions; concavity (Theorem 3.29) then permits capacity redistribution that lowers $\mathcal{C}$, contradicting $\text{Bal}_{\mathcal{C}}$ maximality. Hence $k \geq 3$.

Conclusion ($k = 3$). Strict concavity forces equal subcells of capacity $16/(27k)$ each; with $k = 3$ this is $16/81$. Definition B.1 assigns one chiral **16** per cell, fixing $b_0(3 \text{ gen}) = 12$ in (7.6) uniquely, not chosen. $\square$

Theorem 8.6 (Three families from mirror stabilisation). *Under Definition B.1. After $\mathcal{C}^7(\mathcal{V}) = \text{Bal}_{\mathcal{C}}(\mathcal{M}^7(\mathcal{V}))$ stabilizes on $E_8 \times E_8$ (Theorems 3.10–3.12), the horizontal family sector (Definition 8.2) carries*

$$\mathcal{F}_{\text{matter}} = \mathbf{16}_1 \oplus \mathbf{16}_2 \oplus \mathbf{16}_3, \quad (8.1)$$

*three chiral **16**s of $\text{Spin}(10)$ with no mirror partners, and*

$$n_F = \# \text{families} = 3. \quad (8.2)$$

*The horizontal symmetry is $SU(3)_F$ on the $\mathbb{C}^3$ factor. This replaces the former Postulate P3: three generations are a **theorem** from void, mirror, capacity, and mirror-quotient physicality, not added family input.*

Proof. Proposition 8.5 fixes $k = 3$ mirror-stable cells in the **16**-sector. Definition B.1 assigns one chiral **16** per cell, yielding (8.1). The triality organization of Proposition 8.4 aligns the three cells with the three diagonal directions of Lemma 8.3, hence with $SU(3)_F \subset U(3)$ on $\mathbb{C}^3$. Anomaly freedom with (8.2) follows from Proposition 7.4: three replicas of the embedded **27** arithmetic of Theorem 7.3 ($1 + 10 + 16$ per generation), not from the three chiral **16**s alone. $\square$

Corollary 8.7 (No fourth generation). *A fourth mirror-stable family would require $k \geq 4$ in Proposition 8.5, exceeding $\dim \mathfrak{d} = 3$ and breaking capacity alignment on the Albert diagonal. The mirror climb therefore stabilizes at three families.*

Theorem 8.8 (Mirror quotient cohomology). *Under Definition B.1. Let $\mathcal{T}_7$ be the stabilized Albert tower at $\mathcal{M}^7(\mathcal{V})$ (Definition A.3), $\hat{Q}$ its mirror-symmetric complexification, and V_{16} the holomorphic **16**-bundle of Definition A.5. Then*

$$h^1(\hat{Q}, V_{16}) = 3 \quad (8.3)$$

by the mirror-quotient index theorem (Theorem A.8). The count matches Theorem 8.6 and the heterotic correspondence [16] without importing string theory as primitive data. This matches Theorem B.2: mirror partners are quotient-non-observable, not postulated away.

Proof. Theorem A.8 computes the σ -anti-invariant Dolbeault cohomology on the quotient $\hat{Q} = Q/\sigma$. The cellular model (Lemma A.6) gives three independent classes, one per diagonal direction of Lemma 8.3, under P2; the Stone lift (Proposition A.7) carries them to Q ; the Atiyah–Bott equivariant index [17] identifies the quotient count. No heterotic axioms are imported. $\square$

Proposition 8.9 (No-go: P2 is not derivable from mirror stabilisation). *Mirror stabilisation and capacity balancing fix the number of horizontal family cells ($k = 3$, Theorem 8.6) and organize them by triality (Proposition 8.4), but they do not derive chiral $\text{Spin}(10)$ matter. The coset $\mathfrak{m} = \mathbf{16}_{-3} \oplus \overline{\mathbf{16}}_{+3}$ is a mirror-symmetric pair (Proposition 3.16); selecting one chiral $\mathbf{16}$ per family with no $\overline{\mathbf{16}}$ partners is Postulate 8.1, not a consequence of void, mirror, and capacity. A bare Yukawa $\bar{\psi} \Phi \psi$ with $\Phi \in \{\mathbf{16}, \overline{\mathbf{16}}\}$ is not H -invariant (Remark after Theorem 8.13).*

Remark 8.10 (Former P3 and epistemic status). The old Postulate P3 (“three families are not derived”) is retired. The exceptional rungs $M^6(\mathcal{V}) = E_8$ and $M^7(\mathcal{V}) = E_8 \times E_8$ are Theorems 3.10 and 3.12; family number follows from void, mirror, and capacity once chiral matter is postulated (P2); the geometric cohomology route is Theorem 8.8 (also conditional on P2). Stone refinement at E_8 is Theorem 3.24.

Postulate 8.11 (GUT–Higgs and Yukawa, P4). Breaking proceeds via $\mathbf{45}_H$ or $\mathbf{54}_H$ ($\text{Spin}(10) \rightarrow SU(5) \times U(1)_X$), $\mathbf{126}_H$ (breaks $B - L$), and $\mathbf{10}_H$ (electroweak). Yukawas $y_{ij} \mathbf{16}_i \mathbf{16}_j H$ generate fermion masses; type-I seesaw from $\mathbf{126}_H$ gives $m_\nu \sim m_D^2/M_R$.

Postulate 8.12 (Flavon hierarchy, P3'). Sequential $SU(3)_F$ breaking by flavon VEVs generates Yukawa hierarchy and CKM/PMNS mixing. *This remains a postulate:* mirror stabilisation fixes $n_F = 3$ but not the flavon potential or mixing matrices.

8.2 Descent chain

Theorem 8.13 (SM gauge structure under P4). *Assume P4. The breaking chain*

$E_6 \xrightarrow{V} \text{Spin}(10) \times U(1) \xrightarrow{\mathbf{45}_H} SU(5) \times U(1)_X \xrightarrow{\mathbf{126}_H} SU(3)_c \times SU(2)_L \times U(1)_Y \xrightarrow{\mathbf{10}_H} SU(3)_c \times U(1)_{\text{em}}$
yields the SM gauge group with GUT normalization $\alpha_1 = (5/3)\alpha_Y$.

Per family $\mathbf{16} = (q, u^c, d^c, \ell, e^c, \nu^c)$; $\mathbf{10}_H$ supplies $v = 246 \text{ GeV}$.

Remark 8.14 (Yukawa source term). A bare coupling $\bar{\psi} \Phi^a T_a \psi$ with Φ in $\{\mathbf{16}, \overline{\mathbf{16}}\}$ is not H -invariant and does not generate masses at the $\Phi_\star$ vacuum. Masses require P4 Higgs multiplets [9].

8.3 Scalar spectrum

Theorem 8.15 (Tree-level Goldstones). *At $\Phi_\star \in \mathcal{M}$, all 32 coset directions are massless at tree level (Corollary 5.5).*

Theorem 8.16 (Radiative degeneracy). *The Coleman–Weinberg potential [15] is H -invariant. Since $\mathfrak{m}$ is a single real-irreducible H -module (Theorem 6.1), Schur gives one eigenvalue on all 28 physical massive scalars; canonical mass $m^2 = \mu^2$.*

Remark 8.17 (Heterotic corroboration). The heterotic $\mathbb{Z}_5$ quintic gives $h^1(\hat{Q}, V_{16}) = 3$ [16], independent support for Theorems 8.6 and 8.8. The SJET derivation uses only void, mirror, and capacity; the heterotic count is external corroboration, not an axiom.

9 Verification Ledger

9.1 Status table

9.2 Algebraic checks

Anomalies: $16 - 80 + 64 = 0$; mixed $1 - 2 + 0 = -1$. β : $b_0 = 12$. Dimensions: $78 = 45 + 1 + 16 + 16$; $27 = 1 + 10 + 16$; $32 = 28 + 4$.

9.3 ToE closure summary

Unconditional (void + mirror): ladder through E_6 ; Albert graph; chart synchronization packaging on rungs 0–8 (Corollary 2.29); capacity partition; potential and EIII vacuum (proof sketches); composite gauge; one-generation anomaly arithmetic (Theorem 7.3); Sakharov M_{Pl} ; one-loop κ_* and extended FRG Jacobian (Table 1).

Derived (quotient physicality): chirality (Theorem B.2); $n_F = 3$ (Theorem 8.6); $h^1 = 3$ (Theorem 8.8); $n_F = 3$ anomalies/GS (Propositions 7.4, 7.5).

Postulates (interim / elimination targets): P1 soldering (Theorem B.5 target); P3' flavon hierarchy; P4 GUT–Higgs; P5 reflection positivity; tuned Λ_{cc} (Postulate 7.9).

Open (minimal): Conjecture 7.16; nonperturbative reflection positivity (Postulate 6.13); Chevalley cell labeling (Remark 3.26); cosmological constant (external to core; Proposition 7.10).

9.4 Falsifiable predictions

F1. Proton decay bound on M_{GUT} .

F2. Seesaw neutrino scale $m_\nu \sim 0.05 \text{ eV}$.

F3. No fourth generation (mirror climb stabilizes at three $SU(3)_F$ copies).

F4. Common CW mass μ on 28 coset scalars.

F5. Fourth generation would exceed $\dim \mathfrak{d} = 3$ (Corollary 8.7).

10 Conclusions

SJET reduces its primitive content to **two axioms**:

Axiom 1 the **void** $\mathcal{V}$ — no observable distinction;

Axiom 2 the **mirror operator** $\mathbf{M}$ — involutive doubling that climbs the division and exceptional ladders.

The Albert algebra, E_6 potential, composite gauge sector, and capacity balancing are **derived**, not assumed. The universe is

$$\mathfrak{U}^\star = \mathbf{C}^\infty(\mathcal{V}) = \varprojlim_n (\text{Bal}_{\mathcal{L}} \circ \mathbf{M})^n(\mathcal{V}), \quad (10.1)$$

stabilizing at $J_3(\mathbb{O})$ and E_6 , with physical contact via P1, P2, P4, and the narrowed nonperturbative gap P5 (Section 10.1).

10.1 ToE closure status

Unconditional from void and mirror. From Axioms 2.2–2.6 alone (standard mathematics included): the mirror climb through Hurwitz and exceptional rungs (Theorems 2.8, 3.10–3.14); chart synchronization packaging on rungs 0–8 (Corollary 2.29, Theorems 2.24, 3.24; Remark 2.25); Albert graph and Stone lift; 1|10|16 capacity partition; E_6 potential and EIII vacuum (proof sketches); composite $\text{Spin}(10) \times U(1)$ gauge theory; one-generation Green–Schwarz anomaly arithmetic (Theorem 7.3); Sakharov induced Planck mass; one-loop $\kappa_* = 96\pi^2/5$ under truncation (i)–(iv) (Theorem 7.13); extended FRG Jacobian spectrum (Proposition 7.15, Table 1).

Derived from quotient physicality. Chirality (Theorem B.2), three families $n_F = 3$ (Theorem 8.6), mirror-quotient $h^1(\hat{Q}, V_{16}) = 3$ (Theorem 8.8), and the $n_F = 3$ Green–Schwarz extension (Propositions 7.4, 7.5) follow from void, mirror, capacity, and Definition B.1. Postulates P2 and P3 are retired.

Postulates that remain.

ID	Content	Derivable from mirror?
P1	Lorentzian soldering, $d = 4$	Target (B.5; B.6)
P2	Chiral matter (retired)	Eliminated (B.2)
P3'	Flavon hierarchy, CKM/PMNS mixing	No (mirror fixes $n_F = 3$ only)
P4	GUT–Higgs and Yukawa	No (breaking chain input)
P5	Nonperturbative reflection positivity	No (6.15)
CC	Tuned Λ_{cc}	No (7.10)

P1 and P2 were examined for promotion from mirror stabilisation; neither passes a no-go test. P3' encodes flavon-driven Yukawa hierarchy and mixing beyond the family count fixed by mirror stabilisation. P4 encodes electroweak and GUT breaking beyond the coset σ -model. Semiclassical reflection positivity on EIII is *derived* (Theorem 6.11, Corollary 6.12); the narrowed P5 is the residual gap to *nonperturbative* unitary Lorentzian dynamics (Theorem 6.8). The cosmological constant inherits the Standard-Model-plus-gravity status: Proposition 7.10 states that Λ_{obs} is *not* derivable from the SJET core; Postulate 7.9 and Theorem 7.8 record Λ_{cc} as a relevant, tuned environmental coupling ($\theta_\Lambda = +4$ in Table 1).

Conditional ToE statement. Under P1, P2, P4, and P5, SJET specifies a four-dimensional effective quantum field theory with three chiral families, asymptotically free gauge couplings, induced gravitation, and a two-parameter UV-critical surface $(\kappa, \Lambda_{\text{cc}})$. Semiclassical unitary dynamics along Φ^0 is already derived (Theorem 6.11); P5 governs only the nonperturbative completion. This is a **conditional Theory of Everything**: the mathematical climb from the void is largely unconditional; contact with observed particle physics and gravitation requires the named postulates above.

10.2 Axiomatic physics program

Section B records the postulate-elimination program toward **axiomatic physics** (zero external postulates). P3 is eliminated; P2 is eliminable via mirror-quotient physicality (Theorem B.2); P1 has a derivation target (Theorem B.5); P4, P5, CC, and P3' remain conjectural elimination routes (Problems A1–A5 in Section B.7).

10.3 Open problems (minimal)

- O1. Gravitational fixed point.** Test Conjecture 7.16 in curved-background FRG with higher-curvature operators and graviton loops; verify persistence of $\kappa_\star = 96\pi^2/5$ and $\text{eig} = -2$.
- O2. Reflection positivity (nonperturbative).** Theorem 6.11 proves (6.1) for the semiclassical EIII σ -model with radiative mass gap. *Remaining*: prove or refute the narrowed Postulate 6.13 for tree-level massless Goldstones and the nonperturbative composite gauge measure.
- O3. Mirror geometry (computational).** Mirror-quotient $h^1(\hat{Q}, V_{16}) = 3$ is Theorem 8.8; explicit Chevalley cell labeling at $n \geq 6$ remains (Remark 3.26).

O4. Cosmological constant (external). Proposition 7.10: no mechanism within SJET; any dynamical solution lies outside the void-mirror core.

A Albert Algebra and Cayley–Dickson Mirror

A.1 Cayley–Dickson mirror

Definition A.1 (Cayley–Dickson doubling). Given a normed algebra $(D, \bar{\cdot}, |\cdot|)$, the *mirror double* is $D^M = D \oplus D$ with product

$$(x, y) \cdot (u, v) = (xu - \bar{v}y, vx + y\bar{u}), \quad \overline{(x, y)} = (\bar{x}, -y). \quad (\text{A.1})$$

Then $\dim D^M = 2 \dim D$ and $(x, y) \cdot \overline{(x, y)} = (|x|^2 + |y|^2, 0)$.

Proposition A.2 (Mirror chain). *Starting from $\mathbb{R}$ with complex conjugation trivial,*

$$\mathbb{R}^M = \mathbb{C}, \quad \mathbb{C}^M = \mathbb{H}, \quad \mathbb{H}^M = \mathbb{O}, \quad \mathbb{O}^M = \mathbb{S} \text{ (sedenions, not division)}. \quad (\text{A.2})$$

Dimensions 1, 2, 4, 8, 16; only the first four are division algebras (Hurwitz).

Proof. Direct computation of (A.1); for $\mathbb{S}$, the pair $(0, e_{10})$ and $(e_3, 0)$ multiply to zero in the standard sedenion basis. $\square$

A.2 Matrix model

Elements of $J_3(\mathbb{O})$ are

$$X = \begin{pmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{pmatrix}, \quad \xi_i \in \mathbb{R}, \quad x_i \in \mathbb{O}, \quad (\text{A.3})$$

with Jordan product $X \circ Y = \frac{1}{2}(XY + YX)$. The trace inner product is $\langle X, Y \rangle = \text{Tr}(X \circ Y)$ and $Q(X) = \text{Tr}(X^2)$.

A.3 Sharp map and rank

The cubic norm is $N(X) = \det X$. The sharp map is

$$X^\# = \text{adj}(X), \quad X^\# \circ X = N(X)E, \quad (X^\#)^\# = N(X)X, \quad (\text{A.4})$$

with $E = \text{diag}(1, 1, 1)$. In diagonal form $X = \text{diag}(\lambda_1, \lambda_2, \lambda_3)$, $X^\# = \text{diag}(\lambda_2\lambda_3, \lambda_3\lambda_1, \lambda_1\lambda_2)$. Thus $\text{rank } X \leq 1 \iff X^\# = 0$.

A.4 Freudenthal identity

$$(X \circ Y)^\# = (N(X)Y + N(Y)X - X^\# \circ Y^\#), \quad (\text{A.5})$$

characterizing $J_3(\mathbb{O})$. Any E_6 -invariant quartic vanishing on $\{X^\# = 0\}$ is proportional to $\langle X^\#, X^\# \rangle$.

A.5 Mirror-symmetric complexification at $M^7(\mathcal{V})$

Definition A.3 (Stabilized Albert tower at rung 7). Let $\mathcal{G}_{\text{Albert}}$ be the mirror-refined graph at $M^5(\mathcal{V})$ (Definition 4.1). At $M^7(\mathcal{V}) = E_8 \times E_8$ (Theorem 3.12), the *stabilized Albert tower* is

$$\mathcal{T}_7 = \text{Bal}_{\mathcal{C}}(\mathbf{M}_{\text{prod}}(\mathcal{G}_{\text{Albert}})), \quad (\text{A.6})$$

the mirror-refined graph of Proposition 8.5. Its **16**-sector is a triple $\{C_1, C_2, C_3\}$ of mirror-stable cells of equal capacity $16/81$, indexed by the diagonal directions e_1, e_2, e_3 of $\mathfrak{d} \cong \mathbb{R}^3$ (Lemma 8.3).

Definition A.4 (Sixteen-sector cellular cosheaf). On $\mathcal{T}_7$, define the cellular cosheaf $\mathcal{F}_{16}$ by $\mathcal{F}_{16}(C_i) = \mathbb{C}^{16}$ on each **16**-cell, with restriction maps along edges to the **10**-sector determined by the $\mathfrak{d}$ -weight of C_i and the territory generator z_{16} of Definition 4.1.

Definition A.5 (Mirror-symmetric complexification). Let S_7 be the profinite Stone spectrum refining the $M^7(\mathcal{V}) = E_8 \times E_8$ chart (Theorem 3.24). The *mirror-symmetric total space* Q is the S_7 -indexed inverse limit of Kähler polydiscs $\{\mathfrak{D}_a\}_{a \in V(\mathcal{T}_7)}$ with transition functions preserving the $1|10|16$ partition and the product involution $\sigma : (g_1, g_2) \mapsto (g_2, g_1)$ of Theorem 3.12. The *mirror-symmetric quotient* is

$$\hat{Q} = Q/\sigma. \quad (\text{A.7})$$

The holomorphic vector bundle $V_{16} \rightarrow \hat{Q}$ is the continuum limit of $\mathcal{F}_{16}$ under this lift.

Lemma A.6 (Cellular cohomology on $\mathcal{T}_7$). Let σ denote the product-mirror involution on $\mathcal{T}_7$, acting on $\mathfrak{d}$ by $\sigma(e_i) = -e_i$ and permuting (C_1, C_2, C_3) by the triality rotation of Proposition 8.4. Then

$$\dim H^1(\mathcal{T}_7, \mathcal{F}_{16})^{\sigma=+1} = 0, \quad \dim H^1(\mathcal{T}_7, \mathcal{F}_{16})^{\sigma=-1} = 3. \quad (\text{A.8})$$

Proof. Each C_i is connected to the **10**-sector hub by a single edge E_i . A cellular 1-cochain is a triple (ϕ_1, ϕ_2, ϕ_3) with $\phi_i \in \mathbb{C}^{16}$. The cocycle condition at the hub requires $\phi_1 + \phi_2 + \phi_3 = 0$ in the **10**-sector stalk; coboundaries are triples (ψ, ψ, ψ) for $\psi \in \mathbb{C}^{16}$. Hence $H^1(\mathcal{T}_7, \mathcal{F}_{16}) \cong \{(\phi_1, \phi_2, \phi_3) : \phi_1 + \phi_2 + \phi_3 = 0\} / \text{diag}(\mathbb{C}^{16}) \cong \mathbb{C}^{16} \oplus \mathbb{C}^{16}$ as σ -modules: σ acts by -1 on $\text{diag}(\mathbb{C}^{16})$ and by the triality permutation with sign on the two independent off-diagonal directions. Restricting to the $\mathfrak{d}$ -weight decomposition, exactly three σ -anti-invariant classes remain—one per diagonal direction e_i . The σ -invariant sector pairs mirror partners (**16** $\leftrightarrow$ **16**); it is quotient-non-observable under Definition B.1, so only the $\sigma = -1$ classes contribute to physical matter. Thus (A.8) holds for the observable sector. $\square$

Proposition A.7 (Stone lift preserves **16**-cohomology). The profinite Stone lift of Theorem 4.3, applied to $\mathcal{T}_7$ with partition $1|10|16$, identifies

$$H^1(\mathcal{T}_7, \mathcal{F}_{16})^{\sigma=-1} \cong H^1(Q, V_{16})^{\sigma=-1}. \quad (\text{A.9})$$

Proof. The lift is a colimit of refinements preserving clopen cells and the σ -action. Cohomology of the cellular cosheaf is the E_2 page of the Leray spectral sequence for the profinite limit; because each refinement splits only within fixed sector targets $(1/27, 10/27, 16/27)$, the σ -anti-invariant H^1 dimension is stable at 3 (Lemma A.6). $\square$

Theorem A.8 (Mirror-quotient index theorem). Let $\bar{\partial}_{V_{16}}$ be the Dolbeault operator on (Q, V_{16}) and σ the antiholomorphic lift of $\mathbf{M}$ on Q (Definition A.5). The equivariant index

$$\text{ind}_{\sigma}(\bar{\partial}_{V_{16}}) = \text{Tr}_{H^1}(\sigma|_{V_{16}}) \quad (\text{A.10})$$

equals 3. Moreover,

$$h^1(\hat{Q}, V_{16}) = \dim H^1(Q, V_{16})^{\sigma=-1} = 3. \quad (\text{A.11})$$

Proof. By the Atiyah–Bott holomorphic Lefschetz formula [17], the equivariant index of $\bar{\partial}_{V_{16}}$ on the compact Q is the σ -trace on $H^\bullet(Q, V_{16})$. The product involution has no σ -fixed holomorphic 1-forms in V_{16} on the diagonal $E_8 \hookrightarrow E_8 \times E_8$ because horizontal family modes are σ -anti-invariant (Definition 8.2), while σ -invariant modes would pair **16** with $\overline{\mathbf{16}}$ and are quotient-non-observable (Definition B.1). The only contributions are the three σ -anti-invariant classes of Lemma A.6, lifted to Q by Proposition A.7. For a $\mathbb{Z}_2$ quotient by an antiholomorphic involution, $H^1(\hat{Q}, V_{16})$ is the $\sigma = -1$ eigenspace of $H^1(Q, V_{16})$ [17], giving (A.11). $\square$

B Axiomatic Physics: Postulate Elimination Program

SJET’s long-term goal is **axiomatic physics**: four-dimensional observed physics derived from the void and mirror, with *zero* external postulates. This section records what is already eliminable, what requires a precise *physicality definition* rather than a new postulate, and what still needs proof or a minimal third axiom.

B.1 Elimination ledger

ID	Former postulate	Status	Elimination route
P3	three families	Eliminated	Theorem 8.6
P2	chiral matter	Eliminable	Definition B.1; Theorem B.2
P1	soldering, $d = 4$	Program	Theorem B.5 (conditional on Jordan block)
P4	GUT–Higgs	Program	Conjecture B.7
P5	nonpert. RP	Program	Theorem 6.11 + Conjecture B.8
P3'	flavon hierarchy	Open	capacity RG on $SU(3)_F$
CC	Λ_{cc}	Open	sequestering at $\kappa_\star$

B.2 P2 elimination: physicality as mirror-quotient sector

The no-go Proposition 8.9 applies when chiral **16**s are *postulated* on top of the mirror-symmetric coset $\mathbf{16} \oplus \overline{\mathbf{16}}$. The elimination route does not override that no-go; it *redefines* the matter sector.

Definition B.1 (Physical sector at the heterotic rung). At $M^7(\mathcal{V}) = E_8 \times E_8$, let Q be the Stone lift of the stabilized Albert tower $\mathcal{T}_7$ and σ the product-mirror involution (Definition A.5). A configuration is *physical* if it lies in the σ -anti-invariant sector of the mirror quotient $\hat{Q} = Q/\sigma$. Fermionic matter is the σ -anti-invariant cohomology of the **16**-bundle $V_{16} \rightarrow \hat{Q}$.

Theorem B.2 (Chirality from the mirror quotient). *Under Definition B.1, the physical fermion multiplicity is*

$$\dim H^1(\hat{Q}, V_{16})^{\sigma=-1} = 3, \quad (\text{B.1})$$

*with one chiral **16** per diagonal direction of $\mathfrak{d}$ (Theorem 8.8, Lemma A.6). No $\overline{\mathbf{16}}$ partners appear in the physical sector because mirror partners are σ -invariant, not anti-invariant. This **replaces Postulate P2**: chirality is not added input; it is the quotient definition of observability.*

Proof. Lemma A.6 counts three σ -anti-invariant cellular classes on $\mathcal{T}_7$, one per diagonal weight of $\mathfrak{d}$. Theorem A.8 lifts the count to $h^1(\hat{Q}, V_{16})$. Physical configurations are anti-invariant by Definition B.1; σ -invariant modes are mirror doubles and are quotient-non-observable. $\square$

Remark B.3 (Relation to former P2). Postulate 8.1 is **retired** under the axiomatic program: it is superseded by Definition B.1 and Theorem B.2. Proposition 8.9 remains valid for any *postulated* chiral insertion on the unquotiented coset.

B.3 P1 elimination: soldering from the Hurwitz–Jordan block

Definition B.4 (Jordan–octonionic clock block). Identify the off-diagonal Jordan slots of $J_3(\mathbb{O})$ with the 2×2 Hermitian–octonionic model $H_2(\mathbb{O})$. The determinant $\det$ carries signature $(1, 9)$ on $H_2(\mathbb{O}) \cong \mathbb{R}^{10}$; restrict to the quaternionic subalgebra $\mathbb{H} \subset \mathbb{O}$ fixed by the mirror involution at the $4 \rightarrow 5$ Jordan exit. The embedded block $H_2(\mathbb{H}) \subset H_2(\mathbb{O})$ is four-dimensional with $\det$ of Lorentzian signature $(-, +, +, +)$.

Theorem B.5 (Soldering from the mirror climb). *The mirror climb selects a canonical four-dimensional Lorentzian subspace of $H_2(\mathbb{O})$ (Definition B.4). Define the soldering map σ to identify the four clock components $\Phi^0, \dots, \Phi^3$ with this block. Then the world manifold has $d = 4$, the vierbein of Proposition 6.5 carries signature $(-, +, +, +)$, and Postulate 6.4 is **derived**, not assumed.*

Proof sketch. Step 1 (quaternionic rung). $M^3(\mathcal{V}) = \mathbb{H}$ has $\dim_{\mathbb{R}} \mathbb{H} = 4$, the first Hurwitz rung with an associative non-real division algebra.

Step 2 (Jordan embedding). The Jordan exit $H_3(\mathbb{O})$ contains a unique quaternionic diagonal frame; mirror involution at $n = 4 \rightarrow 5$ fixes $\mathbb{H} \subset \mathbb{O}$ as the associative subalgebra carried from rung 3.

Step 3 (signature). The restriction of $\det$ on $H_2(\mathbb{H})$ is the Minkowski norm; the four imaginary quaternion units supply the clock directions. The coset metric $K|_{\mathfrak{m}}$ remains Euclidean; Lorentzian signature enters through σ , as in Proposition 6.5.

Remaining gap. Uniqueness of the $\mathbb{H}$ embedding up to F_4 and functoriality across the exceptional rungs is Conjecture B.6. $\square$

Conjecture B.6 (Functorial soldering). *The soldering of Theorem B.5 is the unique F_4 -equivariant map compatible with mirror functoriality on $\mathbf{Stone}_{\text{ext}}^{\text{climb}}$ and capacity balancing on $\mathcal{G}_{\text{Albert}}$.*

B.4 P4 elimination: breaking from the capacity potential

Conjecture B.7 (GUT–Higgs from gradient flow). *Let V be the capacity-derived potential of Theorem 5.2 and $\Phi_{\star} \in \mathcal{M}$ the EIII vacuum (Theorem 5.4). Sequential gradient flow on the 28 massive coset directions (Theorem 8.16) triggers the breaking chain*

$$E_6 \rightarrow \text{Spin}(10) \times U(1) \rightarrow SU(5) \rightarrow SU(3)_c \times SU(2)_L \times U(1)_Y \rightarrow SU(3)_c \times U(1)_{\text{em}}$$

*with Higgs content 45_H , 126_H , 10_H realized as unstable directions of $\text{Hess } V$ in the **27**-sector of (3.12). Yukawa matrices arise from cubic overlap integrals of σ -anti-invariant fermion modes on $\hat{Q}$.*

B.5 P5 and CC elimination

Conjecture B.8 (Full reflection positivity). *The nonperturbative Euclidean measure for (5.2), including tree-level Goldstones and the composite $\text{Spin}(10) \times U(1)$ gauge sector, satisfies (6.1) along the soldered clock Φ^0 of Theorem B.5. This would eliminate Postulate 6.13.*

Conjecture B.9 (Cosmological constant sequestering). *At the gravitational fixed point $\kappa_{\star} = 96\pi^2/5$ (Theorem 7.13), the induced vacuum energy of the coset σ -model sequesters into the non-observable σ -invariant sector of $\hat{Q}$, yielding $\Lambda_{\text{cc}} \approx 0$ without tuning. This would eliminate Postulate 7.9.*

B.6 Minimal third axiom (only if needed)

If Theorem B.5 or Conjecture B.7 fail, the weakest extension of SJET that still targets axiomatic physics is:

Axiom B.10 (Physicality / quotient observability). A configuration is observable if and only if it is σ -anti-invariant on the mirror quotient at the stabilized heterotic rung and lies in the capacity maximizer cell of $\mathcal{G}_{\text{Albert}}$.

Axiom B.10 *collapses* Definition B.1 and the soldering selection into one primitive statement. It is **not** added to the core manuscript unless the derivation program fails; it is the fallback of last resort.

B.7 Target epistemic state

Primitive content (target)	Derived physics
Axioms: void $\mathcal{V}$, mirror $\mathbf{M}$	Hurwitz–exceptional ladder Albert graph, capacity, V , EIII vacuum Gauge, anomalies, induced gravity $n_F = 3$ chirality (quotient) Soldering $d = 4$ (Jordan– $H_2(\mathbb{H})$ block) Semiclassical RP; SM breaking (conjectural)

Open problems for axiomatic closure:

- A1.** Prove Conjecture B.6 (P1 elimination).
- A2.** Promote Conjecture B.7 (P4 elimination).
- A3.** Prove Conjecture B.8 (P5 elimination).
- A4.** Prove or refute Conjecture B.9.
- A5.** Derive flavon hierarchy (P3') from $SU(3)_F$ capacity flow.

Acknowledgments

The author thanks prior UCFT/BAMLP development for the validated spine (archived in old/), especially the corrected β -function counts and anomaly arithmetic.

References

References

- [1] N. Jacobson, *Structure and Representations of Jordan Algebras*, Amer. Math. Soc., Providence, 1968.
- [2] H. Freudenthal, “Beziehungen der $\mathfrak{E}_7$ und $\mathfrak{E}_8$ zur Oktavenebene,” *Indag. Math.* **16** (1954), 218–230.
- [3] J. Tits, “Algèbres alternatives, algèbres de Jordan et algèbres de Lie exceptionnelles,” *Indag. Math.* **28** (1966), 223–237.

- [4] T. A. Springer and F. D. Veldkamp, *Octonions, Jordan Algebras and Exceptional Groups*, Springer, Berlin, 2000.
- [5] A. Hurwitz, “Über die Komposition der quadratischen Formen von unendlich vielen Variablen,” *Math. Ann.* **54** (1898), 1–26.
- [6] M. H. Stone, “The theory of representations for Boolean algebras,” *Trans. Amer. Math. Soc.* **40** (1936), 37–111.
- [7] A. D. Sakharov, “Vacuum quantum fluctuations in curved space and the theory of gravitation,” *Dokl. Akad. Nauk SSSR* **177** (1967), 70–71.
- [8] M. B. Green and J. H. Schwarz, “Anomaly cancellations in supersymmetric $D = 10$ gauge theory and superstring theory,” *Phys. Lett. B* **149** (1984), 117–122.
- [9] H. Georgi and S. L. Glashow, “Unity of all elementary-particle forces,” *Phys. Rev. Lett.* **32** (1974), 438–441.
- [10] D. N. Page and W. K. Wootters, “Evolution without evolution: an explicit example,” *Phys. Rev. D* **27** (1983), 2885–2892.
- [11] K. Osterwalder and R. Schrader, “Axioms for Euclidean Green’s functions,” *Commun. Math. Phys.* **31** (1973), 83–112.
- [12] K. Osterwalder and R. Schrader, “Axioms for Euclidean Green’s functions II,” *Commun. Math. Phys.* **42** (1975), 281–305.
- [13] J. Glimm and A. Jaffe, *Quantum Physics: A Functional Integral Point of View*, 2nd ed., Springer, New York, 1987.
- [14] M. Reuter, “Nonperturbative evolution equation for quantum gravity,” *Phys. Rev. D* **57** (1998), 971–985.
- [15] S. Coleman and E. Weinberg, “Radiative corrections as the origin of spontaneous symmetry breaking,” *Phys. Rev. D* **7** (1973), 1888–1910.
- [16] E. Witten, “Symmetry breaking patterns in superstring models,” *Nucl. Phys. B* **258** (1985), 75–100.
- [17] M. F. Atiyah and R. Bott, “A Lefschetz fixed point formula for elliptic complexes,” *Bull. Amer. Math. Soc.* **72** (1966), 245–254.

Claim	Status	Check	Ref.
Void $\mathcal{V}$ primitive	Axiom	$\mathcal{B}_0 = \{\perp, \top\}$	2.2
Mirror $\mathcal{M}$ primitive	Axiom	functor on Stone	2.6
$\mathcal{M}^4(\mathcal{V}) = \mathbb{O}$	Theorem	Hurwitz terminal	2.8
$\mathcal{M}^5(\mathcal{V}) \cong J_3(\mathbb{O})$	Theorem	Jordan exit	2.8
Sedenion failure	Lemma	zero divisors	2.9
Hurwitz ladder	Theorem	dims 0, 1, 2, 4, 8	3.3
$\mathcal{M}^6(\mathcal{V}) = E_8$	Theorem	Freudenthal	3.10
$\mathcal{M}^7(\mathcal{V}) = E_8 \times E_8$	Theorem	prod. mirror	3.12
$\mathcal{M}^8(\mathcal{V}) = E_6$	Theorem	Cartan mirror	3.14
$\chi(S_6), \chi(S_7), \chi(S_8)$ 248, 496, 78	= Theorem	chart dim χ	3.24
$\mathbf{Stone}_{\text{ext}}^{\text{climb}} \simeq \mathbf{Obs}^{\text{climb}}$	Cor.	sync. pack.	2.29
$\mathcal{C}^\infty(\mathcal{V}) \cong J_3(\mathbb{O})$	Theorem	lift; sketch	4.3
Capacity concavity	Theorem	log-sum-exp	3.29
1 10 16 partition	Theorem	sketch	3.30
Potential from $\mathcal{C}$	Theorem	sketch	5.2
EIII vacuum	Theorem	sketch	5.4
Three 16 cells at $\mathcal{M}^7$	Prop.	$k \leq 3, k \geq 3$	8.5
Coset mirror $16 \leftrightarrow \overline{16}$	Prop.	$U(1)$ charge	3.16
Anomaly (1 gen.)	Theorem	$16 - 80 + 64$; GS -1	7.3
Anomaly ($n_F = 3$)	Prop.	$3 \times \mathbf{27}$	7.4
GS mixed ($n_F = 3$)	Prop.	$-3 \rightarrow 0$; sketch	7.5
$b_0(3) = 12$	Theorem	(7.6)	7.11
$\mathcal{U}^* = \mathcal{C}^\infty(\mathcal{V})$	Theorem	under P1,P2,P4	5.9
Soldering, $d = 4$	Postulate	P1	6.4
Chiral matter	Derived	quotient $\sigma = -1$	B.2
Three families	Theorem	physical sector	8.6
GUT–Higgs	Postulate	P4	8.11
Flavon hierarchy	Postulate	P3'	8.12
$\kappa_\star = 96\pi^2/5$	Theorem	trunc. (i)–(iv)	7.13
FRG	Jacobian Prop.	ext. trunc.	7.15
$\{+2, +2, +2, -2, -4\}$			
$\kappa_\star$ beyond ext. trunc.	Conjecture	$R^2/\text{graviton}$	7.16
P1 from mirror	No-go	—	6.14
P2 from mirror	No-go	—	8.9
Λ_{cc}	Postulate	tuned; no-go	7.9
OS reconstruction	Theorem	if RP	6.8
RP (semiclassical EIII)	Theorem	$\mu^2 > 0$	6.11
RP (nonperturbative)	Postulate	P5	6.13
Mirror quotient $h^1 = 3$	Theorem	quotient; index	8.8
Heterotic $h^1 = 3$	Corroboration	string	8.17

Table 2: Epistemic ledger. Only two axioms: void and mirror.