

Lab 10

Definition: The quantile function of a random variable X is the inverse of its cumulative distribution function:

$$Q_X(p) = \min\{x : F_X(x) \geq p\}, \quad 0 < p < 1.$$

Definition: The empirical cumulative distribution function F_n (written ECDF) is the probability distribution that places probability mass $\frac{1}{n}$ on each of the values x_1, x_2, \dots, x_n . The empirical PMF takes the form

$$f_X(x) = \frac{1}{n}, \quad x \in (x_1, x_2, \dots, x_n).$$

If the value x_i is repeated k times, the mass at x_i is accumulated to $\frac{k}{n}$.


More using `distr` and `distrEx`

Defining continuous distribution manually


```
library(distr)
library(distrEx)

# simple discrete distribution
D2 <- DiscreteDistribution(supp = c(1:5), prob = c(0.1, 0.2, 0.3, 0.2, 0.2))
D2

## Distribution Object of Class: DiscreteDistribution
plot(D2)
```



```
f <- function(x) 3 * x^2
X <- AbscontDistribution(d = f, low1 = 0, up1 = 1)
plot(X)
```



```
p(X)(Inf)
```

```
## [1] 1
```

```
p(X)(0)
```


```
## [1] 0
```

```
E(X)
```

```
## [1] 0.7496337
```

Emperical CDF

```
X <- rnorm(100) # X is a sample of 100 normally distributed random variables
ECDF <- ecdf(X) # ECDF is a function giving the empirical CDF of X
plot(ECDF)
```



Multivariate Distributions

```
S <- rolldie(2, makespace = TRUE)
S <- addrv(S, FUN = sum, invars = c("X1", "X2"),
           name = "U")
S <- addrv(S, FUN = min, invars = c("X1", "X2"),
           name = "V")
head(S)
```

```
##   X1 X2 U V      probs
## 1  1  1 2 1 0.02777778
## 2  2  1 3 1 0.02777778
## 3  3  1 4 1 0.02777778
## 4  4  1 5 1 0.02777778
## 5  5  1 6 1 0.02777778
## 6  6  1 7 1 0.02777778
```

```
JointD <- marginal(S, vars = c("U", "V"))
head(JointD)
```

```
##   U V      probs
## 1 2 1 0.02777778
## 2 3 1 0.05555556
## 3 4 1 0.05555556
## 4 5 1 0.05555556
## 5 6 1 0.05555556
## 6 7 1 0.05555556
```

```
xtabs(round(probs, 3) ~ U + V, data = JointD)
```


```
##      V
## U      1      2      3      4      5      6
## 2 0.028 0.000 0.000 0.000 0.000 0.000
## 3 0.056 0.000 0.000 0.000 0.000 0.000
## 4 0.056 0.028 0.000 0.000 0.000 0.000
## 5 0.056 0.056 0.000 0.000 0.000 0.000
## 6 0.056 0.056 0.028 0.000 0.000 0.000
## 7 0.056 0.056 0.056 0.000 0.000 0.000
## 8 0.000 0.056 0.056 0.028 0.000 0.000
## 9 0.000 0.000 0.056 0.056 0.000 0.000
## 10 0.000 0.000 0.000 0.056 0.028 0.000
## 11 0.000 0.000 0.000 0.000 0.056 0.000
## 12 0.000 0.000 0.000 0.000 0.000 0.028
```

```
marginal(JointD, vars = "U")
```

```
##      U      probs
## 1  2 0.02777778
## 2  3 0.05555556
## 3  4 0.08333333
## 4  5 0.11111111
## 5  6 0.13888889
## 6  7 0.16666667
## 7  8 0.13888889
## 8  9 0.11111111
## 9 10 0.08333333
## 10 11 0.05555556
## 11 12 0.02777778
```

Bivariate normal distribution

```
library(mvtnorm)
x <- y <- seq(from = -3, to = 3, length.out = 30)
f <- function(x, y) dmvnorm(cbind(x, y), mean = c(0, 0),
                             sigma = diag(2))
z <- outer(x, y, FUN = f)
persp(x, y, z, theta = -30, phi = 30, ticktype = "detailed")
```



Differentiation

```
f<-expression(x^3+2*x+3)
d<-D(f,'x')
d
```

```
## 3 * x^2 + 2
```

```
d1<-D(D(f,'x'),'x')
d1
```

```
## 3 * (2 * x)
```

Differentiation in two variables

```
f<-expression(x^2+y^2+2*x*y-3*x+4*y+4)
D1=D(f,'x')
D1
```

```
## 2 * x + 2 * y - 3
```

```
f<-expression(x^2+y^2+2*x*y-3*x+4*y+4)
D2=D(f,'y')
D2
```

```
## 2 * y + 2 * x + 4
```

Other differentiation

```
exp<- expression(sin(cos(x + y^2)))
dx<-D(exp,'x');dx
```

```
## -(cos(cos(x + y^2)) * sin(x + y^2))
```

```
dy<-D(exp,'y');dy
```

```
## -(cos(cos(x + y^2)) * (sin(x + y^2) * (2 * y)))
```

Integration

```
func<-function(x)(x^3+2*x^2+3*x+9)
I<-integrate(func,lower=0,upper=3)
I
```

78.75 with absolute error < 8.7e-13

Other Integration

```
integrate(dnorm, -1.96, 1.96)
```

0.9500042 with absolute error < 1e-11

```
integrate(dnorm, -Inf, Inf)
```

1 with absolute error < 9.4e-05

How to evaluate tripple integral?

Evaluate

$$\int_0^{0.5} \int_0^{0.5} \int_0^{0.5} \frac{2}{3} (x^2 + y^2 + z^2) \, dx \, dy \, dz.$$

```
library(cubature)

f <- function(x) { 2/3 * (x[1] + x[2] + x[3]) }
# "x" is vector x[1], x[2], x[3] are referring to x1, x2 and x3 respectively.
adaptIntegrate(f, lowerLimit = c(0, 0, 0), upperLimit = c(0.5, 0.5, 0.5))

## $integral
## [1] 0.0625
##
## $error
## [1] 1.387779e-17
##
## $functionEvaluations
## [1] 33
##
## $returnCode
## [1] 0

## a slowly-convergent integral

integrand <- function(x) {1/((x+1)*sqrt(x))}
integrate(integrand, lower = 0, upper = Inf)
```

3.141593 with absolute error < 2.7e-05

Another package to integrate

Evaluate

$$\int_0^1 \int_0^1 \cos x \cos y \, dx \, dy.$$

```
library('pracma')
fun <- function(x, y) cos(x) * cos(y)
integral2(fun, 0, 1, 0, 1, reltol = 1e-10)
```

```
## $Q
## [1] 0.7080734
##
## $error
## [1] 0
```

Compute the volume of a sphere

Evaluate

$$\int_{x=0}^1 \int_{y=0}^{\sqrt{1-x^2}} \sqrt{1-x^2-y^2} \, dy \, dx.$$

```
f <- function(x, y) sqrt(1 - x^2 - y^2)
xmin <- 0; xmax <- 1
ymin <- 0; ymax <- function(x) sqrt(1 - x^2)
I <- integral2(f, xmin, xmax, ymin, ymax)
I$Q
```

```
## [1] 0.5236076
```

Compute the volume over a sector

```
I <- integral2(f, 0, pi/2, 0, 1, sector = TRUE)
I$Q
```

```
## [1] 0.5236226
```

Integrate $\frac{1}{\sqrt{x+y}(1+x+y)^2}$ over the triangle $0 \leq x \leq 1$, $0 \leq y \leq 1-x$. Note that the integrand is infinite at $(0,0)$.

```
f <- function(x,y) 1/( sqrt(x + y) * (1 + x + y)^2 )
ymax <- function(x) 1 - x
I <- integral2(f, 0, 1, 0, ymax)
I$Q + 1/2 - pi/4
```

```
## [1] -3.247091e-08
```

```
## Compute this integral as a sector
rmax <- function(theta) 1/(sin(theta) + cos(theta))
I <- integral2(f, 0, pi/2, 0, rmax, sector = TRUE, singular = TRUE)
I$Q + 1/2 - pi/4
```

```
## [1] -4.998646e-11
```

Evaluate

$$\int_{x=2}^4 \int_{x-1}^{x+6} \int_{-2}^{4+y^2} (x^2 + y^2 + z) \, dz \, dy \, dx.$$

```
integrand3 <- function(x, y, z) x^2 + y^2 + z
xmin <- 2; xmax <- 4
ymin <- function(x) x - 1
ymax <- function(x) x + 6
```

```
zmin <- -2
zmax <- function(x, y) 4 + y^2
integral3(integrand3, xmin, xmax, ymin, ymax, zmin, zmax)

## [1] 47416.76
```