


## Summary


Pathogens can evolve rapidly in response to changing conditions (e.g., epidemic stage or public health interventions). Models of **eco-evolutionary dynamics** often neglect important epidemiological processes, such as the dynamics of sexual partnerships. We compared **models with a range of complexity** of partnership dynamics and extra-partnership contact.

## Tradeoff theory



- **tradeoff theory**: virulence evolution mediated by transmission-vs-clearance tradeoff
- still debated [1, 2]
- HIV [3]: **set-point viral load** correlated with transmission probability, rate of progression to AIDS (data from Rakai, Uganda)
- eco-evolutionary virulence dynamics: [4]

## Epidemiological structure




Champreon *et al.* 2013 [5]

Spectrum of model complexity:


- **pair formation**: instantaneous or delayed?
- **extra-pair contact (epc)**: present or absent?
- **implicit model**: no explicit partnerships, force of infection expression derived from  $\mathcal{R}_0$  of pair-formation model (without epc)
- **random-mixing model**: standard SIR model  
Simplified disease model (single stage only)

## Sampling strategy




- Latin hypercube sampling: parameters from [5]
- parameters calibrated across models to the same **initial epidemic growth rate ( $r$ )**

## Basic eco-evolutionary dynamics




- significant variation across model structures
- least (random) and most (pairform+epc) models **most similar**: single individuals and extra-pair contact wash out effects of structure
- implicit model is **most different**
- random-mixing model underestimates (?) variability

## Univariate summary



## Bivariate summary



## Conclusions and open questions

- Random-mixing models gave the closest match to the most realistic models; extra-pair contact washed out the effects of epidemiological structure
- Variation among models (model structure)  $\approx$  variation within models (parameter uncertainty)
- Large differences in evolutionary dynamics among different epidemiological models suggest caution in predicting evolutionary responses

## References

- [1] Ebert D, Bull JJ. Challenging the trade-off model for the evolution of virulence: is virulence management feasible? *Trends Microbiol.* 2003;11(1):15–20.
- [2] Alizon S, Michalakis Y. Adaptive virulence evolution: the good old fitness-based approach. *Trends in Ecology & Evolution.* 2015 Jan;30(5):248–254. Available from: <http://www.cell.com/article/S016953471500049X/abstract>.
- [3] Fraser C, Lythgoe K, Leventhal GE, Shirreff G, Hollingsworth TD, Alizon S, et al. Virulence and Pathogenesis of HIV-1 Infection: An Evolutionary Perspective. *Science.* 2014 Mar;343(6177):1243727. Available from: <http://www.sciencemag.org/content/343/6177/1243727>.
- [4] Shirreff G, Pellis L, Laeyendecker O, Fraser C. Transmission Selects for HIV-1 Strains of Intermediate Virulence: A Modelling Approach. *PLoS Computational Biology.* 2011 Oct;7(10):e1002185. WOS:000297262700019.
- [5] Champreon D, Bellan S, Dushoff J. HIV Sexual Transmission Is Predominantly Driven by Single Individuals Rather than Discordant Couples: A Model-Based Approach. *PLoS ONE.* 2013 12;8(12):e82906.

## Acknowledgements

We thank Christophe Fraser for providing access to code, and the Natural Sciences and Engineering Research Council of Canada (NSERC) for funding.

## Sensitivity

