


Summary


Pathogens can evolve rapidly in response to changing conditions (e.g., epidemic stage or public health interventions). Models of **eco-evolutionary dynamics** often neglect important epidemiological processes, such as the dynamics of sexual partnerships. We compared **models with a range of complexity** of partnership dynamics and extra-partnership contact.

Tradeoff theory



- ▶ virulence evolution mediated by transmission-vs-clearance tradeoff
- ▶ still debated [1, 2]
- ▶ HIV [3]: **set-point viral load** correlated with transmission probability, rate of progression to AIDS (data from Rakai, Uganda)
- ▶ eco-evolutionary virulence dynamics: [4]

Epidemiological structure




Champredon *et al.* 2013 [5]


Spectrum of model complexity:

- ▶ **pair formation**: instantaneous or delayed?
 - ▶ **extra-pair contact (epc)**: present or absent?
 - ▶ **implicit model**: no explicit partnerships, force of infection expression derived from R_0 of pair-formation model (without epc)
 - ▶ **random-mixing model**: standard SIR model
- Simplified disease model (single stage only)

Sampling strategy




Eco-evolutionary dynamics




- ▶ significant variation across model structures
- ▶ least (random) and most (pairform+epc) models **most similar**: single individuals and extra-pair contact wash out effects of structure
- ▶ implicit model is **most different**
- ▶ random-mixing model underestimates (?) variability

Univariate summary



Sensitivity



Conclusions and open questions

- ▶ Random-mixing models gave the closest match to the most realistic models; extra-pair contact washed out the effects of epidemiological structure
- ▶ Variation among models (model structure) \approx variation within models (parameter uncertainty)
- ▶ Large differences in evolutionary dynamics among different epidemiological models suggest caution in predicting evolutionary responses

References

- [1] Ebert D, Bull JJ. Challenging the trade-off model for the evolution of virulence: is virulence management feasible? *Trends Microbiol.* 2003;11(1):15–20.
- [2] Alizon S, Michalakos Y. Adaptive virulence evolution: the good old fitness-based approach. *Trends in Ecology & Evolution.* 2015 Jan;30(5):248–254.
- [3] Fraser C, Lythgoe K, Leventhal GE, Shirreff G, Hollingsworth TD, Alizon S, et al. Virulence and Pathogenesis of HIV-1 Infection: An Evolutionary Perspective. *Science.* 2014 Mar;343(6177):1243727.
- [4] Shirreff G, Pellis L, Laeyendecker O, Fraser C. Transmission Selects for HIV-1 Strains of Intermediate Virulence: A Modelling Approach. *PLoS Computational Biology.* 2011 Oct;7(10):e1002185. WOS:000297262700019.
- [5] Champredon D, Bellan S, Dushoff J. HIV Sexual Transmission Is Predominantly Driven by Single Individuals Rather than Discordant Couples: A Model-Based Approach. *PLoS ONE.* 2013 12;8(12):e82906.

Acknowledgements

We thank Christophe Fraser for providing access to code, and the Natural Sciences and Engineering Research Council of Canada (NSERC) for funding.