

Solutions to lab 3 exercise 6

Bob Douma

13 November 2017

Below the solutions to Exercise 6 of Lab 3

1. The first step is to read in the data. The data is stored in different sheet of the file `shapes.xlsx`. Select a sheet, go to **file** and 'save as' and save the sheet as a comma-separated-file (.csv). Beware that if you are using a computer with dutch settings, you may run into problems for two reasons: First, the decimal character is a comma , in Dutch while a dot . in english. Second the default list separator is a semicolon ; in Dutch while a , in English. Check the file that you saved to check the settings of your computer. If this is not correct; go to control panel; to region and language settings; to advanced settings and change the decimal character into a . and the list separator (lijstschijdingsteken) in to a ,.

```
setwd("D://...//...//...") # fill in location
shapes1 = read.csv(shapes1.csv)
```

2. After you have read in the data, you can make a plot through `plot(shapes1$y~ shapes1$x)`. Multiple plots can be made through specifying `par(mfrow=c(3,2))`. This will setup the grid, after using `plot` six times, the grid will be filled with plots.
3. Choosing appropriate deterministic functions

dataset 1 light response curve. There are a number of options of functions to choose from, depending on the level of sophistication: $\frac{ax}{(b+x)}$, $a(1 - e^{(-bx)})$, $\frac{1}{2\theta}(\alpha I + p_{max} - \sqrt{(\alpha I + p_{max})^2 - 4\theta I p_{max}})$ see page 98.

dataset 2 The dataset describes a functional responses. Bolker mentions four of those $\min(ax, s)$
 $\frac{ax}{(b+x)}$, $\frac{ax^2}{(b^2+x^2)}$, $\frac{ax^2}{(b+cx+x^2)}$

dataset 3 Allometric relationship have the form ax^b

dataset 4 This could be logistic growth $n(t) = \frac{K}{1+(\frac{K}{n_0})e^{-rt}}$ or the gompertz function $f(x) = e^{-ae^{-bx}}$

dataset 5 What about a negative exponential? ae^{-bx} or a power function ax^b

dataset 6 Species reponse curves are curves that describe the probability of presence as a function of some factor. A good candidate good be a unimodal response curve. You could take the equation of the normal distribution without the scaling constant: e.g. $ae^{-\frac{(x-\mu)^2}{2\sigma^2}}$

4. For example: `curve(2x+2x,from=0,to=20)` Reasonable values will be supplied in later tutorials