

Model complexity and model choice for animal movement models

Ben Bolker, McMaster University

Departments of Mathematics & Statistics and Biology

Guelph Biomathematics & Biostatistics Symposium

9 June 2016

Outline

- 1 Animal movement
- 2 Florida panthers
- 3 Hidden Markov models
- 4 Basic analysis (van de Kerk et al., 2015)
- 5 Incorporating diurnal variation
- 6 Broader issues/outlook

Acknowledgements

People Michael Li, Madelon van de Kerk, Dave Onorato,
Madan Oli

Agencies US Fish and Wildlife, US Geological Survey, US
National Park Service

Funding NSERC Discovery grant, NSF IGERT program

To do

- pix!
- general REFS: Turchin, Morales et al, Langrock ... ?

Animal movement: data

- observations:
e.g. mass mark-recapture, longitudinal density, direct observation, telemetry (VHF, GPS)
- most methods provide a sequence of times and locations for each individual
- summaries:
 - home range
(convex hull, kernel density estimate, etc.)
 - root-mean-squared displacement
 - step length and turning angle
- covariates:
e.g. habitat map,
individual characteristics (sex, age, weight ...)

Animal movement: questions

- simple description
- how do animals' movements change as a function of their (internal or external) environment?
what does that tell us about their biology?
- how might animals' distributions, etc. change when conditions (density, habitat, ...) change?

Biological/conservation issues

- Florida panther: *Felix ...*
- severely reduced habitat, endangered
- small, isolated population
- currently recovering


(cute panther pictures here)

Panther movement questions


- how does panther movement vary by sex and life history stage
(juvenile, adult, mom with kittens ...)
- how does movement mediate exposure to threats (more cute pictures here ... or roadkill?)
(intraspecific aggression, roadkill) ?
- can we predict the effects of future changes in population density / population structure / habitat?

Panther movement data

- panthers tracked, captured
- GPS collars
- 18 males (13 male, 5 female, 1-15 years old)
- 3200 panther days, hourly/bihourly; 49000 locations
- ?? per panther




example movement tracks



Hidden Markov models

- finite mixture model with temporal dependence
- discrete time steps
- discrete latent state; *transition matrix*
- observations from *emission distributions*
(continuous or discrete, univariate or multivariate)
- **multiphasic movement** (Fryxell et al., 2008; Langrock et al., 2012)



Hidden Markov models (cont.)


notation, underbraces

$$P(\mathbf{Y}_{1:T}, \mathbf{Z}_{1:T} | \boldsymbol{\theta}, \mathbf{X}_{1:T}) = P(z_1 | \mathbf{x}_1) P(\mathbf{y}_1 | z_1, \mathbf{x}_1) \cdot \prod_{k=2}^T P(z_k | z_{k-1}, \mathbf{x}_k) P(\mathbf{y}_k | z_k, \mathbf{x}_k) \quad (1)$$


Hidden Markov models (part 3)

- *forward-backward algorithm* for estimating parameters
- *Viterbi algorithm* for estimating most probable state sequences
- depmixS4 package (Visser and Speekenbrink, 2010) (also moveHMM (Michelot et al., 2016))
- hidden *semi-Markov* models: allow for non-geometric *dwell distributions* (Langrock, 2011; Augustine, 2016): move.HMM


State distributions



Parameter estimates




Tracks with Viterbi estimates



Transition parameters

picture/table here of transition parameters (network diagram??)

Diurnal variation



what can we conclude so far?

good news

- basic biology: males move faster, farther
- three states are identifiable, sensible
- ...

bad news

- diurnal variation in Viterbi results - but it's not in the model!
- estimates of model complexity are too high

what can we conclude so far?


good news

- basic biology: males move faster, farther
- three states are identifiable, sensible
- ...

bad news

- diurnal variation in Viterbi results - but it's not in the model!
- estimates of model complexity are too high

Model complexity



Broadening the model

Attempting to fix these problems:

- extend the model to allow covariates
- specifically, allow for diurnal variation
 - simplify model (log-Normal step length only)
 - *fixed* state-specific emissions parameters (step length mean and std dev)
 - time-varying transition parameters
 - also try *finite mixture models* (independent occupancy)
- how much does this help?

Temporal models

figure showing alternative temporal models

Animal movement
○○

Panthers
○○○○

HMM
○○○

Basic analysis
○○○○○○○

Diurnal model
○○●○○○

Broader issues/outlook
○○○

References

Comparison

Figure 1 from paper

Goodness of fit

Figure 2 from paper

Temporal patterns

Figures 3 (and 4?) from paper

Model complexity

Figure showing BIC plots for all cats tried

Animal movement
○○

Panthers
○○○○

HMM
○○○

Basic analysis
○○○○○○○

Diurnal model
○○○○○○●

Broader issues/outlook
○○○

References

Model complexity

Simulation results

Big data and small models

- simple model families +
model misspecification →
overparameterization

Animal movement
○○

Panthers
○○○○

HMM
○○○

Basic analysis
○○○○○○○

Diurnal model
○○○○○○○

Broader issues/outlook
○○●

References

an aside on AIC vs BIC

Tools needed

- cross-validation (Wenger and Olden, 2012)
- diagnostic plots
- score tests?

References

- Augustine, B., 2016. Flexible, user-friendly hidden (semi) Markov models for animal movement data.
- Fryxell, J.M., Hazell, M., et al., 2008. *Proceedings of the National Academy of Sciences*, 105(49):19114–19119. ISSN 0027-8424, 1091-6490. doi:10.1073/pnas.0801737105.
- Langrock, R., 2011. *Computational Statistics and Data Analysis*, 55(1):715–724. ISSN 01679473.
- Langrock, R., King, R., et al., 2012. *Ecology*, 93(11):2336–2342. ISSN 0012-9658. doi:10.1890/11-2241.1.
- Michelot, T., Langrock, R., and Patterson, T.A., 2016. *Methods in Ecology and Evolution*, in press. doi:10.1111/2041-210X.12578.
- van de Kerk, M., Onorato, D.P., et al., 2015. *Journal of Animal Ecology*, 84(2):576–585.
- Visser, I. and Speekenbrink, M., 2010. *Journal of Statistical Software*, 36(7):1–21.
- Wenger, S.J. and Olden, J.D., 2012. *Methods in Ecology and Evolution*, 3(2):260–267. ISSN 2041210X. doi:10.1111/j.2041-210X.2011.00170.x.