

# Estimation of parameters for stochastic dynamic models

Ben Bolker

McMaster University  
Departments of Mathematics & Statistics and Biology

20 June 2017

# Outline

- 1 Overview
- 2 Stochastic simulation
  - Discrete time
  - Continuous time
- 3 Fitting: simple approaches
  - Trajectory matching
  - Gradient matching
  - Comparison

# Outline

- 1 Overview
- 2 Stochastic simulation
  - Discrete time
  - Continuous time
- 3 Fitting: simple approaches
  - Trajectory matching
  - Gradient matching
  - Comparison

# Modeling

Typical stats	Typical math
stochastic	deterministic
static	dynamic
phenomenological	mechanistic

- Time-series models: mostly **phenomenological** and **linear** (e.g. ARIMA, spectral/wavelet analyses)
- Biomath models: mostly **mechanistic** and **nonlinear** (e.g. Lotka-Volterra, SIR, Fitzhugh-Nagumo)

# Modeling

- **time**: continuous or discrete
- **state**: continuous (e.g. quantitative genetics) or discrete (e.g. Mendelian)
- **evolution**: deterministic or stochastic

e.g.

- ODEs: continuous-time, continuous-state, deterministic
- branching processes: continuous-time, discrete-state, stochastic

# Process and measurement error

- For stochastic models need to define both a **process model** and an **observation model** (= measurement model)

Process model  $Y(t+1) \sim F(Y(t))$

Measurement model  $Y_{\text{obs}}(t) \sim Y(t)$

- Only **process** error affects the future dynamics of the process (usually)
- Might decompose process model into a deterministic model for the expectation and (additive?) noise around the expectation:  
e.g.  $Y(t) = \mu + \epsilon$ ,  $Y(t) \sim \text{Poisson}(\exp(\eta))$

# Process and measurement error

- For stochastic models need to define both a **process model** and an **observation model** (= measurement model)

Process model  $Y(t+1) \sim F(Y(t))$

Measurement model  $Y_{\text{obs}}(t) \sim Y(t)$

- Only **process** error affects the future dynamics of the process (usually)
- Might decompose process model into a deterministic model for the expectation and (additive?) noise around the expectation:  
e.g.  $Y(t) = \mu + \epsilon$ ,  $Y(t) \sim \text{Poisson}(\exp(\eta))$

# Process and measurement error

- For stochastic models need to define both a **process model** and an **observation model** (= measurement model)

Process model  $Y(t+1) \sim F(Y(t))$

Measurement model  $Y_{\text{obs}}(t) \sim Y(t)$

- Only **process** error affects the future dynamics of the process (usually)
- Might decompose process model into a deterministic model for the expectation and (additive?) noise around the expectation:  
e.g.  $Y(t) = \mu + \epsilon$ ,  $Y(t) \sim \text{Poisson}(\exp(\eta))$



# Consequences

- Process error induces dynamic **changes in variance**
- Process+observation error induce **correlations** between subsequent observations
- Observation at next time step depends on **unobserved** value at current time step
- Simple statistical methods (i.e. uncorrelated, equal variance) are incorrect

# Consequences

- Process error induces dynamic **changes in variance**
- Process+observation error induce **correlations** between subsequent observations
- Observation at next time step depends on **unobserved** value at current time step
- Simple statistical methods (i.e. uncorrelated, equal variance) are incorrect



# Consequences

- Process error induces dynamic **changes in variance**
- Process+observation error induce **correlations** between subsequent observations
- Observation at next time step depends on **unobserved** value at current time step
- Simple statistical methods  
(i.e. uncorrelated, equal variance)  
are incorrect

# Consequences

- Process error induces dynamic **changes in variance**
- Process+observation error induce **correlations** between subsequent observations
- Observation at next time step depends on **unobserved** value at current time step
- Simple statistical methods  
(i.e. uncorrelated, equal variance)  
are incorrect

# Linear example



How should we interpret this single realization?


# Outline

- 1 Overview
- 2 Stochastic simulation
  - Discrete time
  - Continuous time
- 3 Fitting: simple approaches
  - Trajectory matching
  - Gradient matching
  - Comparison

# Why simulate?

- understand dynamics
- test methods in best-case scenario
- explore precision/power
- quantify properties of statistical estimators
- evaluate robustness

# Linear model



$$N(1) = a$$

$$N(t+1) \sim \text{Normal}(N(t) + b, \sigma_{\text{proc}}^2)$$

$$N_{\text{obs}}(t) \sim \text{Normal}(N(t), \sigma_{\text{obs}}^2)$$



# R code (version 1)

```
## set up parameters etc.  
nt <- 20; a <- 6; b <- 1  
sd_proc <- sqrt(2)  
sd_obs <- sqrt(2)  
N <- Nobs <- numeric(nt)  
set.seed(101)  ## for reproducibility  
## actual model  
N[1] <- a  
Nobs[1] <- rnorm(1, N[1], sd_obs)  
for (i in 1:nt) {  
  N[i+1] <- rnorm(1, N[i]+b, sd_proc)  
  Nobs[i+1] <- rnorm(1, N[i+1], sd_obs)  
}
```

# R code (version 2)

```
library(deSolve)
linfun <- function(t,y,parms) {
  ## with() is magic to use param names directly
  g <- with(as.list(c(y,parms)), {
    N_new <- rnorm(1,mean=N+b,sd=sd_proc)
    c(N=N_new,Nobs=rnorm(1,mean=N_new,sd=sd_obs))
  })
  return(list(g)) ## deSolve needs this format
}


set.seed(101)
N0 <- c(N=a,Nobs=rnorm(1,a,sd_obs))
linparms <- c(a=6,b=1,sd_proc=sd_proc,sd_obs=sd_obs)
ode(N0,1:nt,linfun,linparms,method="iteration")
```

# R code (version 3)

For this particular example, we can cheat because the process error doesn't affect the future dynamics — it just accumulates:

```
N_det <- a+b*(0:(nt-1))  
set.seed(101)  ## for reproducibility  
proc_noise <- rnorm(nt-1,mean=0,sd=sd_proc)  
N <- N_det+cumsum(c(0,proc_noise))  
N_obs <- rnorm(nt,mean=N,sd=sd_obs)
```

## Hyperbolic nonlinear model



$$N(1) = N_0$$

$$N(t+1) \sim \text{Poisson} \left( \frac{aN(t)}{b+N(t)} \right)$$

$$N_{\text{obs}}(t) \sim \text{Binomial}(N(t), p)$$

# R code

```
hypfun <- function(t,y,parms) {  
  g <- with(as.list(c(y,parms)), {  
    N_det <- a*N/(b+N)  
    N <- rpois(1,lambda=N_det)  
    N_obs <- rbinom(1,size=N,prob=prob_obs)  
    c(N=N,Nobs=N_obs)  
  })  
  return(list(as.numeric(g))) ## deSolve needs numeric() (  
}  
  
set.seed(101)  
NO <- c(N=4,N_obs=4)  
hypparms <- c(a=6,b=1,prob_obs=0.9)  
ode(NO,times=1:nt,func=hypfun,  
    parms=hypparms,method="iteration")
```

# Stochastic ODEs

- continuous-time, continuous-state
- ordinary differential equations plus  $dW$   
(= derivative of a Brownian motion/Wiener process)
- delicate analysis (For biologists: Turelli (1977); Roughgarden (1995). For mathematicians: Øksendal (2003))
- More common for cellular/physiological than population models
- Solve via **Euler-Maruyama**  
(= Euler + appropriately scaled Gaussian noise)

# (continuous-time) Markov processes

- continuous-time, discrete-state
- specify (limits of) probabilities of transitions per unit time, e.g.  $P(N \rightarrow N + 1)$  in the interval  $(t, t + dt)$  is  $rN(t) dt$
- Even harder than SDEs to analyze rigorously ...
- But computationally straightforward: **Gillespie algorithm** and variations (Gillespie, 2007): exponentially distributed time between transitions

# Outline

- 1 Overview
- 2 Stochastic simulation
  - Discrete time
  - Continuous time
- 3 Fitting: simple approaches**
  - Trajectory matching
  - Gradient matching
  - Comparison



# Trajectory matching

- Easiest: simulate the deterministic version of the model (i.e., with neither observation nor process error) and compare
- Because measurement/observation error is (typically) independent at each observation, overall log-likelihood is sum of log-likelihood
- for Normally distributed, equal-variance error, maximum likelihood estimation equivalent to least-squares
- Very common for ODE models, e.g. Gani and Leach (2001); van Veen et al. (2005)
- Brute force can be slow/unstable: use **sensitivity equations** (Raue et al., 2013)

# Pseudo-code

```
## deterministic dynamics:  
## function of parameters, possibly including ICs  
determ_fun <- function(determ_params) {  
  ## code...  
}  
  
## objective function (neg. log-likelihood, SSQ, ...)  
## 'params' includes process and observation parameters  
obj_fun <- function(params,data) {  
  estimate <- determ_fun(params[determ_params])  
  obj <- likfun(estimate,data,params[obs_params])  
  return(obj)  
}  
  
find_minimum(obj_fun,starting_params,...)
```

# Real code #1 (for loops)

```
determ_fun <- function(p,nt) {  
  with(as.list(p),a+b*(1:nt))  
}  
  
obj_fun <- function(p,nt,Nobs) {  
  estimate <- determ_fun(p[c("a","b")],nt)  
  ## negative log-lik. of Normal  
  obj <- -sum(dnorm(Nobs,estimate,p["sd"],log=TRUE))  
  return(obj)  
}  
  
optim(fn=obj_fun,par=c(a=5,b=2,sd=1),nt=20,Nobs=linN)
```

## Real code #2 (using mle2())

```
library(bbmle)
obj_fun <- function(a,b,sd,nt,Nobs) {
  estimate <- determ_fun(a,b,nt)
  ## negative log-lik. of Normal
  obj <- -sum(dnorm(Nobs,estimate,sd,log=TRUE))
  return(obj)
}
determ_fun <- function(a,b,nt) a+b*(1:nt)
mle2(obj_fun,
      data=list(Nobs=linN,nt=nt),
      start=list(a=5,b=2,sd=1.01),
      method="Nelder-Mead")
```

mle2() simplifies computation of confidence intervals, likelihood profiles, etc..

## Real code #3 (mle2() formula interface)



```
library(bbmle)
determ_fun <- function(a,b,nt) a+b*(1:nt)
mle2(Nobs~dnorm(determ_fun(a,b,nt),sd=sd),
     data=list(Nobs=linN,nt=nt),
     start=list(a=5,b=2,sd=1.01),
     method="Nelder-Mead")
```

Formula interface further simplifies getting predicted values, etc.  
(but may make debugging harder!)

# mle2 notes

- wrapper for `optim`
- **assumes** objective function is negative log-likelihood
- uses `method="BFGS"` by default (maybe switch to Nelder-Mead)
- unlike `optim`, obj. function takes parameters separately: `objfun(alpha,beta)` instead of `objfun(params)`
- use `trace=TRUE` to track parameters and obj fun value
- nicer accessors (`coef()`, `logLik()`, etc.: see `methods(class="mle2")`)

# Logistic model fit



# Optimization tips/trouble-shooting

- use sensible starting values
  - for GA/MCMC, use values that are **different** (allow exploration) but **not crazy** (crash/get stuck)
- Nelder-Mead is slower and more robust than BFGS
- test objective/mean function externally
- use `cat()` to print parameter values, see where you're running into trouble
- use constraints (`method="L-BFGS-B"`, or BOBYQA from `nloptr` package) or transform parameters (e.g. fit  $\log(\beta)$  rather than  $\beta$ )





# Gradient matching

- Next-easiest approach: assume **only** process error (no measurement error)
- $N(t+1)$  depends only on  $N(t)$  (which we know exactly):  
**conditional independence**
- **One-step-ahead prediction**
- Simple for discrete-time models  
(we need to specify  $N(t+1) \sim N(t)$  anyway)
- More complicated for continuous-time models (Ellner et al., 2002): fit a smooth curve to data, then fit to derivatives of the curve

# Pseudo-code

```
## deterministic dynamics:
## function of parameters and previous values
onestep_fun <- function(determ_params,Nt) { ... }
## objective function (neg. log-likelihood, SSQ, ...)
obj_fun <- function(params,data) {
  obj <- ... ## numeric vector of length (nt-1)
  for (i in 1:(nt-1)) {
    estimate <- onestep_fun(N[i],params[determ_params])
    obj[i] <- fun(estimate,N[i+1],params[obs_params])
  }
  return(sum(obj))
}
find_minimum(obj_fun,starting_params,...)
```

# Logistic growth fit




# Comparison

How can we use these?

- Try both and hope the answers are not importantly different  
...
- Use biological knowledge of whether process  $\gg$  observation error or vice versa

# Logistic fit comparisons



# References

- Ellner, S.P., Seifu, Y., and Smith, R.H., 2002. *Ecology*, 83(8):2256–2270.
- Gani, R. and Leach, S., 2001. *Nature*, 414(6865):748–751.
- Gillespie, D.T., 2007. *Annual Review of Physical Chemistry*, 58:35–55. ISSN 0066-426X. doi:10.1146/annurev.physchem.58.032806.104637. PMID: 17037977.
- Melbourne, B.A. and Chesson, P., 2006. *Ecology*, 87:1478–1488.
- Raue, A., Schilling, M., et al., 2013. *PLOS ONE*, 8(9):e74335. ISSN 1932-6203. doi:10.1371/journal.pone.0074335.
- Roughgarden, J., 1995. *Theory of Population Genetics and Evolutionary Ecology: An Introduction*. Benjamin Cummings, facsimile edition. ISBN 0134419650.
- Schnute, J.T., 1994. *Canadian Journal of Fisheries and Aquatic Sciences*, 51:1676–1688.
- Turelli, M., 1977. *Theoretical Population Biology*, 12(2):140–178. ISSN 00405809.
- van Veen, F.J.F., van Holland, P.D., and Godfray, H.C.J., 2005. *Ecology*, 86(12):1382–1389.
- Øksendal, B.K., 2003. *Stochastic differential equations: an introduction with applications*. Springer, Berlin; New York. ISBN 3540047581 9783540047582.