


Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.01$, $\theta = 0$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.01$, $\theta = 0.5$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates (epsilon = 0.01) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.01$, $\theta = 1$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates (epsilon = 0.01) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.02$, $\theta = 0$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.02$, $\theta = 0.5$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability

Full 2D ODE; $\psi = 1$, $\rho = 0.02$, $\theta = 1$; $I_0 = 1$




— Numerical finite-horizon Kendall

Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.05$, $\theta = 0$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.05$, $\theta = 0.5$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.05$, $\theta = 1$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.10$, $\theta = 0$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability


Full 2D ODE; $\psi = 1$, $\rho = 0.10$, $\theta = 0.5$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.

Conditional persistence probability

Full 2D ODE; $\psi = 1$, $\rho = 0.10$, $\theta = 1$; $I_0 = 1$



Blue points and grey bars: new adaptive tau-leaping estimates ($\epsilon = 0.01$) and 95% Wilson intervals.
Green: numerical finite-horizon Kendall. Curves stop where the finite-horizon interval does not exist.