

1 description

Let p denote fraction of infected patches and N denote population density of infected patches.




Figure 1: patch state transition

spread: Hosts move around and simultaneously bring pathogens to susceptible patches (combining movement and infection in one step to avoid calculating average density again)

$$f_s : \binom{p}{N} \rightarrow \binom{p + (1 - e^{-Bp})(1 - p)}{N + c(1 - p)(S - N)}$$

B is infection rate (currently setting B as constant since we've already set R_0 as constant too), c is moving probability of individual rats, and S is the population density of susceptible patches (assumed constant).

epidemic: Infected hosts die and so some infected patches burnout.

$$f_e : \binom{p}{N} \rightarrow \binom{P_1 p}{(1 - zD)N}$$

$P_1 = \text{P1_prob}(N)$ is the persistence probability, z is final size of this epidemic wave, D is the death probability of infected hosts.

growth: Assume a constant birth rate r .

$$f_g : \binom{p}{N} \rightarrow \binom{p}{(1 + r)N}$$

The overall iteration process is

$$f = f_g \circ f_e \circ f_s : \binom{p}{N} \rightarrow \binom{P_1(p + (1 - e^{-Bp})(1 - p))}{(1 + r)(1 - zD)(N + c(1 - p)(S - N))}$$

where $P_1 = \text{P1_prob}(N + c(1 - p)(S - N))$

2 parameters

We want to make sure that $f([0, 1]) \subset [0, 1]$. The function attains its maximum at $x^* = \frac{k+1}{2k}$. Note that $f(0) = 0$ and $f(1) = 1$, so we'll need to require that $x^* \geq 1$, i.e. $0 \leq k \leq 1$.

- **infection:** $B > 0$

- **burnout:** $p = P_1 p$, $0 \leq P_1 \leq 1$
- **epidemic:** $0 \leq z \leq 1$ and $0 \leq D \leq 1$
- **movement:** $0 \leq c \leq 1$
- **birth:** $(1 + r)(1 - zD) \leq 1$

name	mode	parameter
incubation period	pneumonic	$1/\sigma_p$
infectious period	pneumonic	$1/\gamma_p$
rat-human ratio	rat-flea	N_r/N_h
rat infectious period	rat-flea	$1/\gamma_r$
rat recovery probability	rat-flea	g_r
incubation period	rat-flea	$1/\sigma_b$
infectious period	rat-flea	$1/\gamma_b$
recovery probability	rat-flea	g_b
prob of 2° pneumonic	rat-flea	p_s
rat-flea index	rat-flea	K_f
flea incubation period	rat-flea	$1/\sigma_f$
flea life away from host	rat-flea	$1/d_f$
louse growth rate	louse	r_l
louse index	louse	K_l
louse infectious period	louse	$1/\gamma_l$

Table 1: Model parameters and their notation.




Figure 2: parameters


2.1 in-patch dynamics (to get R_0)

$$\begin{aligned}
\frac{dS_r}{dt} &= -\beta_r S_r \frac{I_f}{N_r} (1 - e^{-a N_r}) \\
\frac{dI_r}{dt} &= \beta_r S_r \frac{I_f}{N_r} (1 - e^{-a N_r}) - \gamma_r I_r \\
\frac{dR_r}{dt} &= g_r \gamma_r I_r \\
\frac{dD_r}{dt} &= (1 - g_r) \gamma_r I_r \\
\frac{dN_f}{dt} &= r_f N_f \left(1 - \frac{N_f}{K_f N_r}\right) + d_f (1 - e^{-a N_r}) (E_f + I_f) - N_f \frac{(1 - g_r) \gamma_r I_r}{N_r} \\
\frac{dE_f}{dt} &= p_{f,\text{latent}} N_f \frac{(1 - g_r) \gamma_r I_r}{N_r} - (d_f + \sigma_f) E_f \\
\frac{dI_f}{dt} &= (1 - p_{f,\text{latent}}) N_f \frac{(1 - g_r) \gamma_r I_r}{N_r} + \sigma_f E_f - d_f I_f \\
\frac{dS_l}{dt} &= r_l S_l \left(1 - \frac{N_l}{K_l N_h}\right) - \beta_h (I_b + I_p) \frac{S_l}{N_l} \\
\frac{dI_l}{dt} &= \beta_h (I_b + I_p) \frac{S_l}{N_l} - \gamma_l I_l \\
\frac{dS_h}{dt} &= -\beta_b S_h \frac{I_f}{N_h} e^{-a N_r} - \beta_p S_h \frac{I_b}{N_h} - \beta_l S_h \frac{I_l}{N_h} \\
\frac{dE_b}{dt} &= \beta_b S_h \frac{I_f}{N_h} e^{-a N_r} + \beta_l S_h \frac{I_l}{N_h} - \sigma_b E_b \\
\frac{dI_b}{dt} &= \sigma_b E_b - \gamma_b I_b - \alpha I_b \\
\frac{dE_p}{dt} &= \beta_p S_h \frac{I_p}{N_h} - \sigma_p E_p \\
\frac{dI_p}{dt} &= \alpha I_b + \sigma_p E_p - \gamma_p I_p \\
\frac{dR_h}{dt} &= g_b \gamma_b I_b \\
\frac{dD_h}{dt} &= (1 - g_b) \gamma_b I_b + \gamma_p I_p
\end{aligned} \tag{1}$$

Simplified equations at DFE (Ignore searching efficiency, latent period and assume $\frac{N_f}{N_r} = K_f$ at DFE) :

$$\begin{aligned}
\frac{dI_r}{dt} &= \beta_r I_f - \gamma_r I_r \\
\frac{dI_f}{dt} &= \gamma_r K_f I_r - d_f I_f
\end{aligned} \tag{2}$$

This means an infected rat infects an average of K_f fleas and an infected flea infects an average of $\frac{\beta_r}{d_f}$ rats, so expected number of new infected rats generated by an infected rat is $R_{0\text{rat to rat}} = \frac{\beta_r K_f}{d_f}$. (By the way, R_0 in the burnout paper is $\frac{\mu}{\mu + \gamma}$ (μ is the natural death rate of rats, γ is the death rate caused by disease, $\mu \ll \gamma$), if we accept this R_0 , we might need to replace all γ with $\frac{d_f}{K_f}$, and this also makes a difference to ε)



We have priors on the three parameters:

$$\beta_r \sim \text{Gamma}(2, 0.25)$$

$$K_f \sim \text{Gamma}(5, 1.4)$$

$$\frac{1}{d_f} \sim \text{Gamma}(4, 1)$$

By sampling them we get distribution of R_0 and the peak is at $R_0 = 2.75$.

3 Disease Free Equilibrium

If we start at a point where $p > 0, N > 0$, all variables remain positive during the iterations as each step maps positive values to a positive values.

If we start at $p = 0, N > 0$ then population size ends up at

$$N_t = N_{t+1} = (1 + r)(1 - zD)(N_t + (p - 1)(S - N_t))$$

i.e.

$$N_{DFE} = \frac{c(1 + r)(1 - zD)S}{1 - (1 + r)(1 - zD)(1 - c)}$$

N_{DFE} increases with r, c, S and decreases with z, D .

4 invasion analysis

$$J_{f_s} = \begin{pmatrix} (1 + (1 - p)B)e^{-Bp} & 0 \\ -c(S - N) & 1 \end{pmatrix}$$

$$J_{f_e} = \begin{pmatrix} P_1 & p \frac{\partial P_1}{\partial N} \\ 0 & 1 - zD \end{pmatrix}$$

$$J_{f_g} = \begin{pmatrix} 1 & 0 \\ 0 & 1 + r \end{pmatrix}$$

And at DFE we have $p = 0, N = N_{DFE}$ so the matrices above are all lower triangular, so eigenvalues of $J_f = J_{f_g} J_{f_e} J_{f_s}$ are $\lambda_1 = (1 + B)P_1$ and $\lambda_2 = (1 - zD)(1 + r)$. We required previously that λ_2 is always smaller than 1. If $(1 + B)P_1|_{N=N_{DFE}+c(S-N_{DFE})} = (1 + B)P_1|_{N=\frac{cS}{1-(1+r)(1-zD)(1-c)}} < 1$, DFE is stable, so the disease cannot persist; if it is greater than 1, DFE is unstable. Simulation shows that the disease will persist and system ends up at endemic equilibrium.

5 Endemic equilibrium

In the hazard model version the endemic equilibrium is given by

$$\begin{aligned} P_1(p + (1 - e^{-Bp})(1 - p)) &= p \quad (p \neq 0) \\ (1 + r)(1 - zD)(N + c(1 - p)(S - N)) &= N \end{aligned}$$

where

$$P_1 = p_k(1 - q(x_{in})^{y_\star(N+c(1-p)(S-N))})$$

$p_k, q(x_{in}), y_\star$ are all defined in burnout paper. This is equivalent to

$$1 - \frac{N(t-1)}{c(S-N)} = p_k(1 - e^{y_\star t \ln q(x_{in})N})(1 - \frac{N(t-1)}{c(S-N)} e^{-B(1 - \frac{N(t-1)}{c(S-N)})})$$

We can't have an analytic solution even with W_r function (although we can always solve it numerically). (haven't proved EE is stable; we can always compute the eigenvalue numerically. I think there is a more elegant approach to prove it's stable but I just don't know ...)




Figure 3: unstable DFE

6 Order of events

We have $f = f_g \circ f_e \circ f_s$ and this leads to

$$N_{DFE} = \frac{c(1+r)(1-zD)S}{1 - (1+r)(1-zD)(1-c)}$$

and persistence condition

$$(1+B)P_1|_{N=\frac{cS}{1-(1+r)(1-zD)(1-c)}} > 1$$

We can do the same thing for $f_1 = f_s \circ f_g \circ f_e$ (cyclic permutation) and get

$$N_{DFE_1} = \frac{cS}{1 - (1+r)(1-zD)(1-c)}$$

and threshold for persistence is the same

$$(1+B)P_1|_{N=N_{DFE_1}} = (1+B)P_1|_{N=\frac{cS}{1-(1+r)(1-zD)(1-c)}} > 1$$

Equations for endemic equilibrium of f_1 is a little bit more complicated than f but we could also expect equivalent result since $\lim_{n \rightarrow \infty} f^n = \lim_{n \rightarrow \infty} f_1^n \circ f_s$

However if we consider non-cyclic permutation, for example, $f_2 = f_g \circ f_s \circ f_e$, then

$$N_{DFE_2} = \frac{cS(1+r)}{1 - (1+r)(1-zD)(1-c)}$$

And persistence condition is

$$(1 + B)P_1|_{N=N_{DFE_2}} > 1$$

$N_{DFE_2} > N_{DFE_1}$ and P_1 increases with N , so the disease becomes easier to persist. This makes sense biologically: with this order infected hosts breed before they die, and larger population size makes pathogens more likely to survive. (The previous order seems more reasonable than this one, as rats die very quickly after being infected.)

7 effect of R_0

According to the threshold, larger B and larger $P_1|_{N=\frac{cS}{1-(1+r)(1-zD)(1-c)}}$ is favored, and P_1 increases with N , i.e., increases with r, c, S and decreases with z, D . However, P_1 increases with R_0 so seems that smaller R_0 would not be favored.




Figure 4: P_1 increases with R_0

It might be the decrease of D that was favored in the evolution.




Figure 5: decrease in D makes disease persist