


Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.01, theta = 0; I0 = 1



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference



Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence


Exact Gillespie CTMC; rho = 0.01, theta = 0.5; I0 = 1



Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.01, theta = 1; I0 = 1



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.02$, $\theta = 0$; $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.02$, $\theta = 0.5$; $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence


Exact Gillespie CTMC; rho = 0.02, theta = 1; I0 = 1



Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.05$, $\theta = 0$; $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.05$, $\theta = 0.5$; $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- Standard: $\sqrt{y^*/K}$
- - - Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.05$, $\theta = 1$; $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.10$, $\theta = 0$; $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.10$, $\theta = 0.5$; $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.10$, $\theta = 1$; $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference

Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. Solid curves use the standard $\sqrt{y^*/K}$ height; dashed curves use the alternative y^* height. Each curve stops where its matching equation has no admissible root.