


Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.01, theta = 0; l0 = 1; matching height = 3/4 compromise



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.01, theta = 0.5; l0 = 1; matching height = 3/4 compromise



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.01, theta = 1; I0 = 1; matching height = 3/4 compromise



Analytical method


—	Exact Kendall + first-order entry	—	Exact Kendall + second-order entry
---	Laplace + first-order entry	---	Laplace + second-order entry

Matching height — $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.02, theta = 0; I0 = 1; matching height = 3/4 compromise



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.02, theta = 0.5; I0 = 1; matching height = 3/4 compromise



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height — $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.02, theta = 1; I0 = 1; matching height = 3/4 compromise



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle


Exact Gillespie CTMC; rho = 0.05, theta = 0; l0 = 1; matching height = 3/4 compromise



Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.05, theta = 0.5; I0 = 1; matching height = 3/4 compromise



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.05, theta = 1; I0 = 1; matching height = 3/4 compromise



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.10, theta = 0; l0 = 1; matching height = 3/4 compromise



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle


Exact Gillespie CTMC; rho = 0.10, theta = 0.5; l0 = 1; matching height = 3/4 compromise



Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.

Stochastic validation conditional on escaping early fizzle

Exact Gillespie CTMC; rho = 0.10, theta = 1; I0 = 1; matching height = 3/4 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = K^{-1/4} y_{star}^{3/4}$

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 3/4 compromise. Each curve stops where its matching equation has no admissible root.