


Unconditional persistence probability

$\rho = 0.01$, $\theta = 0$, $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Persistence probability conditional on escaping early fizzle

$\rho = 0.01$, $\theta = 0$, $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Unconditional persistence probability

$\rho = 0.01$, $\theta = 0.5$, $l_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Persistence probability conditional on escaping early fizzle

$\rho = 0.01$, $\theta = 0.5$, $l_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Unconditional persistence probability

$\rho = 0.01$, $\theta = 1$, $I_0 = 1$



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry


Matching height

- Standard: $\sqrt{y^*/K}$
- Alternative: y^*

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Persistence probability conditional on escaping early fizzle

$\rho = 0.01$, $\theta = 1$, $I_0 = 1$



Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.