

Epidemic Burnout under Susceptible Recruitment and Population-Size-Dependent Transmission

A Nonlinear Tail-Matching Theory from the First Epidemic Wave to Kendall Persistence Probability

Theory note

2026-09-08

Abstract

We develop a self-contained asymptotic and semi-analytical theory for pathogen burnout or persistence after the first epidemic wave in a susceptible–infectious–death model with susceptible recruitment $h_\theta(x) = (1 - x)x^\theta$ and population-size-dependent transmission. The normalized deterministic system is

$$\dot{x} = \rho h_\theta(x) - R_0 x y (x + y)^{-\psi}, \quad \dot{y} = \{R_0 x (x + y)^{-\psi} - 1\} y,$$

with $0 \leq \theta \leq 1$, $0 \leq \psi < 1$, $0 < \rho \ll 1$, and a large reference population K . The zero-recruitment first-wave geometry is explicit and supplies a regular outer approximation on any fixed interval away from the post-wave endpoint. The epidemic tail is then continued by the exact one-dimensional inverse-flow equation, written in logarithmic prevalence, without expanding the finite-prevalence transmission factor $(1 + y/x)^{-\psi}$. This nonlinear tail continuation remains well defined when y/x is not small and therefore does not require a Taylor expansion around a possibly very small final-size anchor. We derive explicit deterministic-to-stochastic overlap diagnostics, an action-based plateau quantity whose variation is rigorously controlled, the corresponding time-inhomogeneous Kendall branching formula, and its Laplace reduction. The construction recovers the usual logarithmic corner constants in the stronger local overlap, but those constants are asymptotic limits rather than required inputs. The special case $\psi = 0$ is regular and particularly simple because the per-infective transmission rate is independent of prevalence. The endpoint $\psi \rightarrow 1^-$ is regular at the level of the nonlinear deterministic tail but nonuniform for the full burnout theory because the low-prevalence threshold and the interior Laplace saddle collapse toward zero.

Contents

1	Executive formula sheet	5
1.1	Model and zero-recruitment geometry	5
1.2	Regular outer initialization	5
1.3	Nonlinear tail equation	6
1.4	Overlap diagnostics and the action plateau	6
1.5	Kendall prediction	7

1.6	Laplace and boundary-layer-independent form	7
1.7	Scope	8
2	Count-level model and normalization	8
3	Zero-recruitment first-wave geometry	9
3.1	Exact invariant and epidemic orbit	9
3.2	Final-size anchor	10
3.3	Low-prevalence threshold and positive equilibrium	11
4	Regular outer expansion on a fixed first-wave interval	13
5	Exact nonlinear continuation of the descending epidemic tail	14
5.1	Inverse-flow equation	14
5.2	Existence on the descending branch	15
5.3	Why the nonlinear continuation is uniform in finite prevalence	15
5.4	Alternative total-population representation	16
6	Deterministic-to-stochastic overlap	16
6.1	Finite-prevalence transmission error	16
6.2	Susceptible-feedback error	17
6.3	Population-size condition	17
7	Action identity and the nonlinear tail amplitude	17
7.1	Action primitive	17
7.2	Exact drift identity	18
7.3	Effective tail amplitude	18
7.4	Admissible handoff	19
8	Kendall branching layer	19
8.1	Reduced stochastic dynamics after handoff	19
8.2	Probabilistic status of the handoff	19
8.3	One-lineage extinction probability	20
8.4	Many infectives at handoff	20
9	Laplace reduction and handoff-independent leading form	21
9.1	Interior action saddle	21

9.2	Leading burnout exponent	22
9.3	Next saddle coefficient	22
10	Asymptotic reduction of the nonlinear tail	22
10.1	Local variables	23
10.2	Action reduction	23
10.3	Second-order interpretation	24
11	Special parameter limits	25
11.1	The case $\psi = 0$	25
11.2	Fixed $0 < \psi < 1$	25
11.3	The endpoint $\psi \rightarrow 1^-$	25
12	Validity conditions and identifiable failure modes	26
12.1	Regular outer validity	26
12.2	Descending inverse-flow validity	26
12.3	Deterministic-to-stochastic overlap	26
12.4	Near-threshold loss of saddle separation	27
12.5	Large R_0 and positive ψ	27
12.6	The singular endpoint $\psi = 1$	27
13	Computational workflow	27
13.1	Precomputation for fixed (R_0, θ, ψ)	27
13.2	Loop over ρ	28
13.3	Loop over K	28
14	Verification targets	29
14.1	Zero-recruitment checks	29
14.2	Outer-initialization independence	29
14.3	Exact deterministic regression	29
14.4	Action plateau	29
14.5	Handoff invariance	29
14.6	$\psi = 0$ regression	29
14.7	Positive- ψ stress test	30
15	Recruitment-law derivatives	30

16 Detailed derivation of the regular outer coefficient equations	30
16.1 Exact drift of the zero-recruitment invariant	30
16.2 Expansion of the invariant itself	31
16.3 First outer coefficient	31
16.4 Second outer coefficient	32
17 Detailed local inverse hierarchy and finite-part constants	32
17.1 Exact inverse-flow hierarchy	33
17.2 Analytic normal form at the anchor	33
17.3 First coefficient and the logarithmic amplitude	34
17.4 Second coefficient and the finite constant	35
17.5 Action cancellation at second order	36
18 Derivation of the Laplace correction	37
19 Summary of the theoretical chain	39

1 Executive formula sheet

1.1 Model and zero-recruitment geometry

The normalized deterministic equations are

$$\dot{x} = \rho h_\theta(x) - g(x, y)y, \quad \dot{y} = \{g(x, y) - 1\}y,$$

where

$$g(x, y) = R_0 x(x + y)^{-\psi}, \quad h_\theta(x) = (1 - x)x^\theta, \quad \eta = 1 - \psi > 0.$$

At $\rho = 0$, with $n = x + y$,

$$n_0(x) = \left(1 + \frac{\eta}{R_0} \log x\right)^{1/\eta}, \quad F_\psi(x) = n_0(x) - x.$$

The post-wave endpoint x_f is the nontrivial root

$$F_\psi(x_f) = 0,$$

and the low-prevalence growth threshold is

$$x_* = R_0^{-1/\eta}.$$

It is useful to define

$$z_f = x_f^\eta.$$

Then z_f is the unique nontrivial root in $(0, 1/R_0)$ of the ψ -independent equation

$$\boxed{z_f = 1 + \frac{1}{R_0} \log z_f}, \quad \boxed{x_f = z_f^{1/\eta}}.$$

Consequently

$$0 < x_f < x_* < 1, \quad q := R_0 x_f^\eta \in (0, 1), \quad \delta := 1 - q > 0.$$

Thus increasing ψ compresses x_f toward zero even when the transformed final size z_f is unchanged. The existence, uniqueness, and ordering are proved in Section 3.

1.2 Regular outer initialization

Choose a fixed outer point

$$\bar{x} \in (x_f, x_*)$$

away from both endpoints, and set

$$\bar{y} = F_\psi(\bar{x}).$$

On the regular outer interval, expand

$$y = F_\psi(x) + \rho Y_1(x) + \rho^2 Y_2(x) + O(\rho^3).$$

The coefficient equations are given explicitly in Section 4. At fixed $y = \bar{y}$, inversion gives

$$X(\bar{y}; \rho) = \bar{x} + \rho \bar{X}_1 + \rho^2 \bar{X}_2 + O(\rho^3),$$

where

$$\bar{X}_1 = -\frac{Y_1(\bar{x})}{F'_\psi(\bar{x})},$$

$$\bar{X}_2 = -\frac{\frac{1}{2}F''_\psi(\bar{x})\bar{X}_1^2 + Y'_1(\bar{x})\bar{X}_1 + Y_2(\bar{x})}{F'_\psi(\bar{x})}.$$

1.3 Nonlinear tail equation

On the descending branch $g < 1$, define the logarithmic prevalence coordinate

$$\ell = \log \frac{\bar{y}}{y}, \quad y = \bar{y}e^{-\ell}.$$

The exact phase-plane dynamics reduce to the scalar non-autonomous ODE

$$\boxed{\frac{dX}{d\ell} = \frac{\rho h_\theta(X) - g(X, y)y}{1 - g(X, y)}, \quad y = \bar{y}e^{-\ell},}$$

with initial condition

$$X(0) = \bar{x} + \rho \bar{X}_1 + \rho^2 \bar{X}_2.$$

No expansion of

$$(1 + y/X)^{-\psi}$$

is made in this tail equation.

1.4 Overlap diagnostics and the action plateau

Define the low-prevalence rate

$$g_0(x) = R_0 x^\eta, \quad g(x, y) = g_0(x) \left(1 + \frac{y}{x}\right)^{-\psi},$$

and the two dimensionless overlap ratios

$$s = \frac{y}{x}, \quad d = \frac{g(x, y)y}{\rho h_\theta(x)}.$$

The first measures finite-prevalence correction to the per-infective birth rate; the second measures infection depletion relative to susceptible recruitment.

Define the action through finite differences,

$$A_{\theta, \psi}(v) - A_{\theta, \psi}(u) = \int_u^v \frac{1 - R_0 z^\eta}{h_\theta(z)} dz.$$

Along the exact deterministic trajectory,

$$\boxed{\frac{d}{d\sigma} \left(\log y + \frac{A_{\theta, \psi}(x)}{\rho} \right) = g - g_0 - (1 - g_0)d.}$$

Since $\dot{\ell} = 1 - g$ on the descending branch,

$$\boxed{\frac{d}{d\ell} \log \mathcal{C}_\rho = \frac{g - g_0 - (1 - g_0)d}{1 - g},}$$

where

$$\boxed{\mathcal{C}_\rho(y) = y \exp \left\{ \frac{A_{\theta, \psi}(X(y)) - A_{\theta, \psi}(x_f)}{\rho} \right\} .}$$

For $0 \leq \psi \leq 1$,

$$0 \leq g_0 - g = g_0 \left[1 - (1 + s)^{-\psi} \right] \leq \psi g_0 s,$$

so

$$\left| \frac{d}{d\ell} \log \mathcal{C}_\rho \right| \leq \varepsilon_A := \frac{\psi g_0 s + |1 - g_0| d}{1 - g}.$$

An admissible deterministic-to-stochastic handoff point (x_h, y_h) satisfies

$$Ky_h \gg 1, \quad s_h \ll 1, \quad d_h \ll 1, \quad \varepsilon_{A,h} \ll 1, \quad g_h < 1.$$

A plateau of $\mathcal{C}_\rho(y)$ across several admissible handoff levels is the principal internal diagnostic.

1.5 Kendall prediction

At an admissible handoff, neglecting infectious feedback on x and finite- y/x correction to the per-infective birth rate gives

$$\dot{x} = \rho h_\theta(x), \quad b(\sigma) = R_0 x(\sigma)^\eta, \quad d(\sigma) = 1.$$

For one lineage starting at x_h define

$$\mathcal{I}(x_h) = \int_{x_h}^1 \frac{1}{\rho h_\theta(X)} \exp \left\{ \frac{A_{\theta,\psi}(X) - A_{\theta,\psi}(x_h)}{\rho} \right\} dX.$$

Then

$$q_1 = \frac{\mathcal{I}}{1 + \mathcal{I}}$$

is the asymptotic extinction probability of one lineage. With $m = Ky_h$ lineages,

$$P_{\text{persist}|\text{est}} \simeq 1 - \left(\frac{\mathcal{I}(x_h)}{1 + \mathcal{I}(x_h)} \right)^{Ky_h}.$$

For one initial infective before the first wave,

$$P_{\text{persist,uncond}} \simeq \left(1 - \frac{1}{R_0} \right) P_{\text{persist}|\text{est}}.$$

1.6 Laplace and boundary-layer-independent form

The action has its unique interior maximum at

$$x_* = R_0^{-1/\eta}, \quad A''_{\theta,\psi}(x_*) = -\frac{\eta}{x_* h_\theta(x_*)}.$$

Hence

$$\mathcal{I}(x_h) \sim \exp \left\{ \frac{A_{\theta,\psi}(x_*) - A_{\theta,\psi}(x_h)}{\rho} \right\} \sqrt{\frac{2\pi x_*}{\eta \rho h_*}}, \quad h_* = h_\theta(x_*).$$

When $\mathcal{I} \gg 1$,

$$B := -\log P_{\text{burn}|\text{est}} = Ky_h \log(1 + 1/\mathcal{I}) \sim \frac{Ky_h}{\mathcal{I}}.$$

Therefore

$$B_{\text{NL}} \sim K \mathcal{C}_{\rho,h} \sqrt{\frac{\eta \rho h_*}{2\pi x_*}} \exp \left\{ -\frac{\Delta A_f}{\rho} \right\},$$

where

$$\mathcal{C}_{\rho,h} = \mathcal{C}_{\rho}(y_h), \quad \Delta A_f = A_{\theta,\psi}(x_*) - A_{\theta,\psi}(x_f).$$

If $\mathcal{C}_{\rho}$ has reached a plateau, the leading Laplace prediction is asymptotically independent of the arbitrary handoff level.

1.7 Scope

The theory requires fixed

$$0 \leq \psi < 1, \quad R_0 > 1, \quad \rho \ll 1, \quad K \gg 1,$$

together with a nonempty deterministic-to-stochastic overlap. The nonlinear tail itself remains meaningful when $y/x = O(1)$; only the later branching handoff requires $y/x \ll 1$. At $\psi = 0$ the finite-prevalence transmission correction vanishes identically. The endpoint $\psi = 1$ is not covered by the full theory because $x_* \rightarrow 0$ as $\psi \rightarrow 1^-$, so the interior action saddle collapses to the boundary.

2 Count-level model and normalization

Let S and I be susceptible and infectious counts, $N = S + I$, K a reference population size, γ the infectious removal rate, rK the susceptible-recruitment scale, and β the transmission parameter. The infection event rate is

$$\lambda_{SI} = \frac{\beta}{K} SI \left(\frac{K}{N} \right)^{\psi}.$$

The deterministic count equations are

$$\frac{dS}{dt} = rK h_{\theta}(S/K) - \frac{\beta}{K} SI \left(\frac{K}{S+I} \right)^{\psi},$$

$$\frac{dI}{dt} = \frac{\beta}{K} SI \left(\frac{K}{S+I} \right)^{\psi} - \gamma I.$$

Set

$$x = \frac{S}{K}, \quad y = \frac{I}{K}, \quad \sigma = \gamma t, \quad \rho = \frac{r}{\gamma}, \quad R_0 = \frac{\beta}{\gamma}.$$

Since $N/K = x + y$,

$$\frac{\beta}{K} (Kx)(Ky)(x+y)^{-\psi} = \beta K xy (x+y)^{-\psi},$$

so division by $K\gamma$ gives

$$\dot{x} = \rho h_{\theta}(x) - R_0 xy (x+y)^{-\psi},$$

$$\dot{y} = \left\{ R_0 x (x+y)^{-\psi} - 1 \right\} y.$$

The deterministic fraction system is independent of K . Population size re-enters when a fraction y is converted to the infectious count Ky in the stochastic layer.

Define

$$g(x, y) = R_0 x(x + y)^{-\psi}.$$

Then

$$\dot{x} = \rho h_\theta(x) - g(x, y)y, \quad \dot{y} = \{g(x, y) - 1\}y.$$

For $x > 0$ write

$$s = \frac{y}{x}.$$

The per-infective birth factor has the exact decomposition

$$g(x, y) = R_0 x^{1-\psi} (1 + s)^{-\psi}.$$

With $\eta = 1 - \psi$ define

$$g_0(x) = R_0 x^\eta.$$

Then

$$g = g_0(1 + s)^{-\psi}.$$

This identity separates two effects: $g_0(x)$ is the low-prevalence growth factor, while $(1 + s)^{-\psi}$ is the finite-prevalence correction. When $\psi = 0$ the second factor is exactly one at every prevalence.

3 Zero-recruitment first-wave geometry

3.1 Exact invariant and epidemic orbit

Set $\rho = 0$ and $n = x + y$. Adding the equations gives

$$\dot{n} = -y.$$

Also

$$\dot{x} = -R_0 x y n^{-\psi}.$$

Therefore, wherever $y > 0$,

$$\frac{dn}{dx} = \frac{n^\psi}{R_0 x}.$$

Let

$$\eta = 1 - \psi > 0.$$

Then

$$n^{-\psi} dn = \frac{dx}{R_0 x},$$

so, using the epidemic initial state $(x, n) = (1, 1)$,

$$\frac{n^\eta - 1}{\eta} = \frac{1}{R_0} \log x.$$

Hence

$$n_0(x) = \left(1 + \frac{\eta}{R_0} \log x\right)^{1/\eta},$$

and the zero-recruitment epidemic branch is

$$F_\psi(x) = n_0(x) - x.$$

Equivalently,

$$H(x, y) = n^\eta - \frac{\eta}{R_0} \log x$$

is constant when $\rho = 0$.

3.2 Final-size anchor

At the end of the zero-recruitment epidemic, $y = 0$, so $n = x = x_f$. Therefore

$$x_f^\eta = 1 + \frac{\eta}{R_0} \log x_f.$$

Define

$$z_f = x_f^\eta.$$

Because $\log x_f = (1/\eta) \log z_f$,

$$z_f = 1 + \frac{1}{R_0} \log z_f.$$

The transformed final-size equation is independent of ψ , while

$$x_f = z_f^{1/\eta}.$$

The location and uniqueness of the nontrivial root require a short argument because the resulting ordering will be used throughout the theory. Define

$$\phi(z) = z - 1 - \frac{1}{R_0} \log z, \quad z > 0.$$

Then

$$\phi'(z) = 1 - \frac{1}{R_0 z}, \quad \phi''(z) = \frac{1}{R_0 z^2} > 0.$$

Hence ϕ is strictly convex and has its unique minimum at

$$z_* = \frac{1}{R_0}.$$

The point $z = 1$ is one root, since $\phi(1) = 0$. At the minimum,

$$\phi(z_*) = \frac{1}{R_0} - 1 + \frac{\log R_0}{R_0} = -\frac{R_0 - 1 - \log R_0}{R_0} < 0,$$

because $R_0 - 1 - \log R_0 > 0$ for $R_0 > 1$. Moreover,

$$\lim_{z \downarrow 0} \phi(z) = +\infty.$$

By continuity, ϕ therefore has a root in $(0, 1/R_0)$. Strict convexity implies that there can be at most two positive roots; since $z = 1$ is already the second one, the root in $(0, 1/R_0)$ is unique. Denote it by z_f . Thus

$$0 < z_f < \frac{1}{R_0} < 1.$$

Since $x_f = z_f^{1/\eta}$ and $x_* = R_0^{-1/\eta}$, monotonicity of $z \mapsto z^{1/\eta}$ gives

$$0 < x_f < x_* < 1.$$

Furthermore,

$$q = R_0 x_f^\eta = R_0 z_f,$$

so

$$0 < q < 1, \quad \delta = 1 - q > 0.$$

These inequalities establish, rather than assume, the ordering of the final-size anchor and the later low-prevalence threshold. They also ensure that all local coefficients containing δ^{-1} are well defined. Finally, because z_f depends only on R_0 , increasing ψ decreases η and compresses $x_f = z_f^{1/\eta}$ rapidly toward zero.

Differentiating n_0 gives

$$n'_0(x) = \frac{n_0(x)^\psi}{R_0 x}.$$

Thus

$$F'_\psi(x) = \frac{n_0(x)^\psi}{R_0 x} - 1.$$

At $x = x_f$, since $n_0(x_f) = x_f$,

$$F'_\psi(x_f) = \frac{1}{R_0 x_f^\eta} - 1.$$

Writing

$$q = R_0 x_f^\eta, \quad \delta = 1 - q,$$

we have

$$F'_\psi(x_f) = \frac{\delta}{q} > 0.$$

The positivity follows from the root-ordering argument above, which proved $0 < q < 1$ and $\delta > 0$.

3.3 Low-prevalence threshold and positive equilibrium

As $y/x \rightarrow 0$,

$$g(x, y) \rightarrow g_0(x) = R_0 x^\eta.$$

The low-prevalence growth threshold is therefore

$$x_* = R_0^{-1/\eta}.$$

At a positive deterministic equilibrium,

$$g(x_e, y_e) = 1, \quad \rho h_\theta(x_e) = y_e,$$

so

$$y_e = \rho h_\theta(x_e)$$

exactly and

$$R_0 x_e \{x_e + \rho h_\theta(x_e)\}^{-\psi} = 1.$$

Let

$$x_e = x_* + \rho \xi + O(\rho^2), \quad h_* = h_\theta(x_*).$$

Because $y_e = \rho h_\theta(x_e)$,

$$y_e = \rho h_* + O(\rho^2).$$

Take logarithms of the equilibrium condition:

$$0 = \log R_0 + \log x_e - \psi \log(x_e + y_e).$$

The two logarithms have the first-order expansions

$$\log x_e = \log x_* + \rho \frac{\xi}{x_*} + O(\rho^2),$$

$$\log(x_e + y_e) = \log x_* + \rho \frac{\xi + h_*}{x_*} + O(\rho^2).$$

Since $R_0 x_*^\eta = 1$,

$$\log R_0 + \eta \log x_* = 0.$$

Substitution therefore leaves, at order $O(\rho)$,

$$0 = \frac{\rho}{x_*} \{\xi - \psi(\xi + h_*)\} + O(\rho^2) = \frac{\rho}{x_*} (\eta \xi - \psi h_*) + O(\rho^2).$$

Hence

$$\boxed{\eta \xi - \psi h_* = 0},$$

and therefore

$$\boxed{x_e = x_* + \rho \frac{\psi h_*}{\eta} + O(\rho^2)},$$

$$\boxed{y_e = \rho h_* + O(\rho^2)}.$$

The demographic infectious scale is thus $O(\rho)$.

4 Regular outer expansion on a fixed first-wave interval

The nonlinear tail will be initialized from a point that remains a fixed positive distance from the zero-recruitment endpoint. On such an interval the perturbation expansion in ρ is regular and no corner expansion is needed.

Choose

$$\bar{x} \in (x_f, x_*)$$

with $\bar{x} - x_f = O(1)$ relative to the local perturbation scale, and define

$$\bar{y} = F_\psi(\bar{x}) > 0.$$

On $[\bar{x}, 1]$ write

$$\begin{aligned} y &= F_\psi(x) + \rho Y_1(x) + \rho^2 Y_2(x) + O(\rho^3), \\ n &= n_0(x) + \rho Y_1(x) + \rho^2 Y_2(x) + O(\rho^3). \end{aligned}$$

The exact phase-plane equation is

$$\frac{dy}{dx} = \frac{\{R_0 x(x+y)^{-\psi} - 1\}y}{\rho h_\theta(x) - R_0 x y(x+y)^{-\psi}}.$$

Introduce

$$W_1 = n_0^{-\psi} Y_1, \quad W_2 = n_0^{-\psi} Y_2 - \frac{\psi}{2} n_0^{-\psi-1} Y_1^2.$$

Expansion of $n^{-\psi}$ and collection of equal powers of ρ gives the sequential equations

$$\boxed{W_1' = \frac{h_\theta(x) F_\psi'(x)}{R_0 x F_\psi(x)}}.$$

At the next order,

$$\boxed{\begin{aligned} W_2' &= \frac{h_\theta(x)}{R_0^2 x^2 F_\psi(x)^2} \left[\psi n_0(x)^{\psi-1} F_\psi(x) - \{n_0(x)^\psi - R_0 x\} \right] Y_1(x) \\ &\quad + \frac{h_\theta(x)^2 n_0(x)^\psi \{n_0(x)^\psi - R_0 x\}}{R_0^3 x^3 F_\psi(x)^2}. \end{aligned}}$$

The original coefficients are recovered through

$$\boxed{Y_1 = n_0^\psi W_1},$$

$$\boxed{Y_2 = n_0^\psi W_2 + \frac{\psi}{2n_0} Y_1^2}.$$

Because $h_\theta(1) = 0$, the endpoint conditions

$$Y_1(1) = 0, \quad Y_2(1) = 0$$

are understood by continuous extension from the interior. The equations are integrated only to $\bar{x} > x_f$, where every coefficient is regular.

To initialize the inverse trajectory at the fixed prevalence $\bar{y} = F_\psi(\bar{x})$, write

$$X(\bar{y}; \rho) = \bar{x} + \rho \bar{X}_1 + \rho^2 \bar{X}_2 + O(\rho^3).$$

Substituting this into

$$\bar{y} = F_\psi(X) + \rho Y_1(X) + \rho^2 Y_2(X) + O(\rho^3)$$

and equating powers of ρ gives

$$0 = F'_\psi(\bar{x})\bar{X}_1 + Y_1(\bar{x}),$$

therefore

$$\boxed{\bar{X}_1 = -\frac{Y_1(\bar{x})}{F'_\psi(\bar{x})}.$$

At $O(\rho^2)$,

$$0 = F'_\psi(\bar{x})\bar{X}_2 + \frac{1}{2}F''_\psi(\bar{x})\bar{X}_1^2 + Y'_1(\bar{x})\bar{X}_1 + Y_2(\bar{x}),$$

so

$$\boxed{\bar{X}_2 = -\frac{\frac{1}{2}F''_\psi(\bar{x})\bar{X}_1^2 + Y'_1(\bar{x})\bar{X}_1 + Y_2(\bar{x})}{F'_\psi(\bar{x})}.$$

This produces an $O(\rho^3)$ outer initialization error at fixed $\bar{x}$ if the displayed regular expansion is carried through second order.

5 Exact nonlinear continuation of the descending epidemic tail

5.1 Inverse-flow equation

On any branch on which y is monotone and positive, divide the deterministic equations:

$$\frac{dx}{dy} = \frac{\rho h_\theta(x) - g(x, y)y}{\{g(x, y) - 1\}y}.$$

After the first epidemic peak and before the first trough,

$$g(x, y) < 1, \quad \dot{y} < 0.$$

Introduce

$$\ell = \log \frac{\bar{y}}{y}.$$

Because

$$\dot{\ell} = -\frac{\dot{y}}{y} = 1 - g(x, y),$$

we obtain directly

$$\frac{dx}{d\ell} = \frac{\dot{x}}{\dot{\ell}} = \frac{\rho h_\theta(x) - g(x, y)y}{1 - g(x, y)}.$$

Thus the descending tail is governed by the scalar equation

$$\boxed{\frac{dX}{d\ell} = \frac{\rho h_\theta(X) - R_0 X y (X + y)^{-\psi}}{1 - R_0 X (X + y)^{-\psi}}, \quad y = \bar{y} e^{-\ell}.$$

The initial condition is supplied by the regular outer solution:

$$X(0) = \bar{x} + \rho \bar{X}_1 + \rho^2 \bar{X}_2.$$

The scalar equation is exactly equivalent to the two-dimensional deterministic trajectory on the monotone descending branch; the only approximation in this stage is the outer initial value. If the initial value were exact, the inverse-flow continuation would reproduce the exact deterministic phase-plane curve up to the point where $g = 1$ and y ceases to be monotone.

5.2 Existence on the descending branch

Fix a compact region

$$\mathcal{D} = \{(x, y) : x \geq x_{\min} > 0, y \geq 0, g(x, y) \leq 1 - \gamma_0\}$$

with $\gamma_0 > 0$. On $\mathcal{D}$ the right-hand side of the scalar equation is smooth in x and continuous in ℓ , with denominator bounded away from zero. Standard ODE existence, uniqueness, and continuous-dependence results therefore apply. In particular, an $O(\rho^3)$ initialization error remains $O(\rho^3)$ on any fixed ℓ interval contained in such a region, with a constant determined by the corresponding Lipschitz bound.

The condition $g \leq 1 - \gamma_0$ is not required for numerical integration all the way to a practical handoff point; it is used here only to state a simple uniform continuous-dependence result. The inverse coordinate necessarily becomes singular at the exact trough $g = 1$, because y itself has a turning point there. The stochastic handoff is made before that turning point.

5.3 Why the nonlinear continuation is uniform in finite prevalence

Write

$$g(x, y) = g_0(x)(1 + s)^{-\psi}, \quad g_0(x) = R_0 x^\eta, \quad s = \frac{y}{x}.$$

The nonlinear tail equation keeps $(1 + s)^{-\psi}$ exactly. Consequently the construction does not require

$$s = \frac{y}{x} \ll 1$$

during the deterministic continuation from $\bar{y}$ down to the stochastic overlap.

This distinction matters when x_f is small. Since

$$x_f = z_f^{1/\eta},$$

a moderate reduction of $\eta = 1 - \psi$ can make x_f orders of magnitude smaller while leaving the transformed final size z_f unchanged. A prevalence that is absolutely small can then still satisfy $y/x_f = O(1)$ or larger. The nonlinear continuation is defined without expanding in that ratio.

At $\psi = 0$,

$$g(x, y) = R_0 x$$

exactly, so finite prevalence never modifies the per-infective transmission factor. For $\psi > 0$, the correction is monotone:

$$0 < (1 + s)^{-\psi} \leq 1.$$

5.4 Alternative total-population representation

For structural interpretation define

$$n = x + y, \quad p = \frac{x}{n}.$$

Then

$$x = pn, \quad y = (1 - p)n, \quad g = R_0 p n^\eta.$$

Adding the original equations gives

$$\dot{n} = \rho h_\theta(pn) - (1 - p)n.$$

Using $p = x/n$ gives

$$\dot{p} = (1 - p) \left[\frac{\rho h_\theta(pn)}{n} + p\{1 - R_0 n^\eta\} \right].$$

Thus the population-size-dependent transmission appears only through n^η . The pointwise deterministic vector field has a well-defined limit as $\psi \rightarrow 1^-$ ($\eta \rightarrow 0^+$), where $n^\eta \rightarrow 1$ for fixed $n > 0$. This observation concerns the deterministic tail only; the low-prevalence saddle structure of the burnout problem is not uniform at $\psi = 1$, as discussed later.

6 Deterministic-to-stochastic overlap

The nonlinear tail may be continued through a regime in which y/x is not small. The branching approximation is introduced only after the trajectory reaches a region in which two separate feedback mechanisms are small.

6.1 Finite-prevalence transmission error

The exact per-infective birth rate is

$$g = g_0(1 + s)^{-\psi}.$$

The relative finite-prevalence correction is

$$1 - \frac{g}{g_0} = 1 - (1 + s)^{-\psi}.$$

For $s \geq 0$ and $0 \leq \psi < 1$, the mean-value theorem gives

$$0 \leq 1 - (1 + s)^{-\psi} \leq \psi s.$$

Therefore

$$\frac{|g - g_0|}{g_0} \leq \psi \frac{y}{x}.$$

The first overlap condition is consequently

$$s_h = \frac{y_h}{x_h} \ll 1.$$

At $\psi = 0$ this error vanishes identically, regardless of s_h .

6.2 Susceptible-feedback error

The susceptible equation is

$$\dot{x} = \rho h_\theta(x) - g(x, y)y.$$

The approximation used in the branching layer is

$$\dot{x} \simeq \rho h_\theta(x).$$

Its natural relative error is

$$d(x, y) = \frac{g(x, y)y}{\rho h_\theta(x)}.$$

Thus the second overlap condition is

$$d_h \ll 1.$$

This condition is more precise than comparing y with ρ alone because it includes the local transmission factor and the local recruitment law.

6.3 Population-size condition

At handoff, the deterministic infectious fraction corresponds to

$$m = Ky_h$$

infectious individuals. To initialize the stochastic branching layer from a concentrated deterministic state one requires

$$Ky_h \gg 1.$$

The three primary overlap requirements are therefore

$$Ky_h \gg 1, \quad \frac{y_h}{x_h} \ll 1, \quad \frac{g_h y_h}{\rho h_\theta(x_h)} \ll 1.$$

The existence of a y_h satisfying all three conditions is a substantive assumption. If no such level exists, the deterministic and branching descriptions do not possess a clean common overlap at that finite K .

7 Action identity and the nonlinear tail amplitude

7.1 Action primitive

Define the action only through finite differences:

$$A_{\theta, \psi}(v) - A_{\theta, \psi}(u) = \int_u^v \frac{1 - R_0 z^\eta}{h_\theta(z)} dz.$$

Thus

$$A'_{\theta, \psi}(x) = \frac{1 - g_0(x)}{h_\theta(x)}.$$

The action is increasing for $x < x_*$, stationary at x_* , and decreasing for $x > x_*$.

7.2 Exact drift identity

Define

$$J(x, y) = \log y + \frac{A_{\theta, \psi}(x)}{\rho}.$$

Along the full deterministic model,

$$\frac{d}{d\sigma} \log y = g - 1.$$

Also

$$\frac{1}{\rho} \frac{d}{d\sigma} A_{\theta, \psi}(x) = \frac{1 - g_0}{\rho h_{\theta}(x)} \{\rho h_{\theta}(x) - gy\} = (1 - g_0) - (1 - g_0)d.$$

Adding the two identities yields

$$\dot{J} = g - g_0 - (1 - g_0)d.$$

This is exact. No low-prevalence expansion has been made.

The first term measures the finite-prevalence change in the infectious birth rate. The second term measures infectious feedback on susceptible recovery. In the common overlap both terms are small, and J becomes an approximate invariant.

7.3 Effective tail amplitude

Subtract the constant $A_{\theta, \psi}(x_f)/\rho$ and exponentiate. Define

$$\mathcal{C}_{\rho}(y) = y \exp \left\{ \frac{A_{\theta, \psi}(X(y)) - A_{\theta, \psi}(x_f)}{\rho} \right\}.$$

Then

$$\log \mathcal{C}_{\rho} = J - \frac{A_{\theta, \psi}(x_f)}{\rho}.$$

Since $\dot{\ell} = 1 - g$ on the descending branch,

$$\frac{d}{d\ell} \log \mathcal{C}_{\rho} = \frac{g - g_0 - (1 - g_0)d}{1 - g}.$$

This identity quantifies handoff dependence directly.

For $s \geq 0$,

$$g_0 - g = g_0 \{1 - (1 + s)^{-\psi}\} \leq \psi g_0 s.$$

Therefore

$$\left| \frac{d}{d\ell} \log \mathcal{C}_{\rho} \right| \leq \frac{\psi g_0 s + |1 - g_0|d}{1 - g} =: \varepsilon_A.$$

If $y_1 > y_2$ are two handoff levels on the descending branch, corresponding to $\ell_1 < \ell_2$, integration gives the rigorous bound

$$\left| \log \frac{\mathcal{C}_{\rho}(y_2)}{\mathcal{C}_{\rho}(y_1)} \right| \leq \int_{\ell_1}^{\ell_2} \varepsilon_A(\ell) d\ell.$$

Thus a plateau in $\mathcal{C}_{\rho}$ is not merely a numerical convenience; it is the direct signature that the action invariant has entered its overlap regime.

7.4 Admissible handoff

A practical handoff point (x_h, y_h) on the nonlinear tail should satisfy

$$\boxed{Ky_h \gg 1, \quad s_h \ll 1, \quad d_h \ll 1, \quad \varepsilon_{A,h} \ll 1, \quad g_h < 1.}$$

The first three conditions validate the deterministic-to-branching approximation. The ε_A condition additionally ensures that the action amplitude is locally insensitive to the arbitrary handoff level. In computation, it is preferable to identify an interval of admissible levels and verify a plateau rather than to rely on one prescribed boundary.

A traditional geometric-mean candidate,

$$y_{\text{geo}} = \sqrt{\frac{\rho h_*}{K}},$$

may be used as an initial guess, but it is accepted only if the displayed overlap diagnostics are satisfied there.

8 Kendall branching layer

8.1 Reduced stochastic dynamics after handoff

Inside the overlap, infectious feedback on x is negligible to leading order, so the susceptible background follows

$$\boxed{\dot{x} = \rho h_\theta(x).}$$

The finite-prevalence correction to infectious transmission is also negligible, so each infectious individual evolves as a time-inhomogeneous linear birth–death lineage with

$$\boxed{b(\sigma) = R_0 x(\sigma)^\eta, \quad d(\sigma) = 1.}$$

This is a branching approximation because lineages no longer interact through the susceptible equation at leading order.

8.2 Probabilistic status of the handoff

The preceding reduction is an asymptotic branching approximation. A stochastic weak-convergence theorem is not proved in this note. The conditions

$$Ky_h \gg 1, \quad s_h \ll 1, \quad d_h \ll 1$$

have distinct roles: $Ky_h \gg 1$ places the deterministic handoff above the few-particle scale, $s_h \ll 1$ makes the finite-prevalence correction to the per-capita birth rate small, and $d_h \ll 1$ makes infectious depletion negligible in the susceptible equation. Together they identify a deterministic-to-branching overlap in which the reduced birth–death description is asymptotically consistent with the deterministic drift. They do not, by themselves, constitute a pathwise coupling, a functional central-limit theorem, or a Kurtz-type weak-convergence result for the full density-dependent jump process. Establishing such a theorem would require specifying the underlying finite- K Markov chain and a joint limit for K , ρ , and the handoff scale. The persistence formulas below should therefore be read as semi-analytical branching approximations conditional on this overlap.

8.3 One-lineage extinction probability

For a time-inhomogeneous birth–death process with one initial individual, Kendall’s formula may be written

$$q_1 = \frac{\mathcal{I}}{1 + \mathcal{I}},$$

where

$$\mathcal{I} = \int_0^\infty \exp \left\{ - \int_0^t [b(u) - d(u)] du \right\} d(t) dt.$$

Here $d(t) = 1$. Along the susceptible recovery path,

$$d\sigma = \frac{dX}{\rho h_\theta(X)}.$$

Moreover,

$$\int_0^{\sigma(X)} [b(u) - d(u)] du = \int_{x_h}^X \frac{R_0 z^\eta - 1}{\rho h_\theta(z)} dz = - \frac{A_{\theta,\psi}(X) - A_{\theta,\psi}(x_h)}{\rho}.$$

Therefore

$$\mathcal{I}(x_h) = \int_{x_h}^1 \frac{1}{\rho h_\theta(X)} \exp \left\{ \frac{A_{\theta,\psi}(X) - A_{\theta,\psi}(x_h)}{\rho} \right\} dX.$$

The upper endpoint is integrable. As $X \uparrow 1$, $h_\theta(X) \sim 1 - X$ and

$$A_{\theta,\psi}(X) = A_1 + (R_0 - 1) \log(1 - X) + o(1),$$

more precisely the exponential factor contributes $(1 - X)^{(R_0 - 1)/\rho}$, so the full integrand behaves as

$$\rho^{-1} (1 - X)^{(R_0 - 1)/\rho - 1},$$

which is integrable for $R_0 > 1$.

8.4 Many infectives at handoff

With

$$m = Ky_h$$

approximately independent lineages, the burnout probability is

$$P_{\text{burn|est}} \simeq \left(\frac{\mathcal{I}(x_h)}{1 + \mathcal{I}(x_h)} \right)^{Ky_h}.$$

Hence

$$P_{\text{persist|est}} \simeq 1 - \left(\frac{\mathcal{I}(x_h)}{1 + \mathcal{I}(x_h)} \right)^{Ky_h}.$$

For numerical stability define

$$B = -\log P_{\text{burn|est}} = Ky_h \log \left(1 + \frac{1}{\mathcal{I}(x_h)} \right),$$

so

$$P_{\text{persist}|\text{est}} = 1 - e^{-B}.$$

If the epidemic starts from one infective in an otherwise susceptible population, the classical early establishment probability is approximately

$$1 - \frac{1}{R_0}.$$

The unconditional persistence probability through the first trough is therefore

$$P_{\text{persist,uncond}} \simeq \left(1 - \frac{1}{R_0}\right) P_{\text{persist}|\text{est}}.$$

9 Laplace reduction and handoff-independent leading form

9.1 Interior action saddle

The action derivative is

$$A'_{\theta,\psi}(x) = \frac{1 - R_0 x^\eta}{h_\theta(x)}.$$

Hence the unique interior stationary point is

$$x_* = R_0^{-1/\eta}.$$

At $x = x_*$ the numerator vanishes, and differentiation gives

$$A''_{\theta,\psi}(x_*) = -\frac{\eta}{x_* h_*}, \quad h_* = h_\theta(x_*).$$

Thus x_* is a nondegenerate maximum for fixed $\eta > 0$.

Assume the lower endpoint is separated from the saddle by more than the Gaussian width,

$$x_* - x_h \gg \sqrt{\rho},$$

and define

$$\Lambda_h = \frac{A_{\theta,\psi}(x_*) - A_{\theta,\psi}(x_h)}{\rho}.$$

For $\Lambda_h \gg 1$, Laplace's method gives

$$\mathcal{I}(x_h) \sim \exp(\Lambda_h) \sqrt{\frac{2\pi x_*}{\eta \rho h_*}}.$$

9.2 Leading burnout exponent

When $\mathcal{I} \gg 1$,

$$\log(1 + 1/\mathcal{I}) = \frac{1}{\mathcal{I}} + O(\mathcal{I}^{-2}).$$

Therefore

$$B \sim K y_h \exp \left\{ -\frac{A_{\theta,\psi}(x_*) - A_{\theta,\psi}(x_h)}{\rho} \right\} \sqrt{\frac{\eta \rho h_*}{2\pi x_*}}.$$

Insert

$$y_h \exp \left\{ \frac{A_{\theta,\psi}(x_h) - A_{\theta,\psi}(x_f)}{\rho} \right\} = \mathcal{C}_{\rho,h}.$$

With

$$\Delta A_f = A_{\theta,\psi}(x_*) - A_{\theta,\psi}(x_f),$$

we obtain

$$B_{\text{NL}} \sim K \mathcal{C}_{\rho,h} \sqrt{\frac{\eta \rho h_*}{2\pi x_*}} \exp \left(-\frac{\Delta A_f}{\rho} \right).$$

If $\mathcal{C}_{\rho,h}$ lies on an action plateau, the leading expression is independent of the arbitrary handoff level up to the integral plateau error derived in Section 7.

9.3 Next saddle coefficient

Write the Laplace expansion as

$$\mathcal{I}(x_h) \sim \exp(\Lambda_h) \sqrt{\frac{2\pi x_*}{\eta \rho h_*}} (1 + \rho c_L + O(\rho^2)).$$

For $h = h_\theta$, the coefficient is

$$c_L = \frac{(2\eta^2 - \eta - 1)h_*^2 + (\eta - 1)x_*h_*h'_* + 2x_*^2(h'_*)^2 - 3x_*^2h_*h''_*}{24\eta h_*x_*}.$$

The derivatives and term-by-term algebra reducing the general Laplace coefficient to this closed form are displayed in Section 18. Consequently,

$$B_{\text{NL,next}} \sim B_{\text{NL}} (1 - \rho c_L + O(\rho^2) + O(\mathcal{I}^{-1})).$$

An exponentialized form,

$$B_{\text{NL,next}} \approx B_{\text{NL}} e^{-\rho c_L},$$

is equivalent through relative order $O(\rho)$.

10 Asymptotic reduction of the nonlinear tail

The nonlinear tail equation is the primary deterministic bridge. In the stronger regime where the trajectory also becomes local relative to x_f , it reduces to the familiar logarithmic corner structure. This section derives that reduction and identifies the asymptotic meaning of the action amplitude.

10.1 Local variables

Let

$$x = x_f + \xi, \quad y \rightarrow 0, \quad \frac{|\xi| + y}{x_f} \ll 1.$$

At the anchor,

$$q = R_0 x_f^\eta, \quad \delta = 1 - q > 0.$$

Also define

$$h_f = h_\theta(x_f), \quad \boxed{a = \frac{h_f}{\delta}}, \quad \boxed{b = \frac{q}{\delta}}.$$

The zero-recruitment inverse branch satisfies

$$X_0(y) = x_f + by + O(y^2),$$

because

$$X'_0(0) = \frac{1}{F'_\psi(x_f)} = \frac{q}{\delta} = b.$$

For sufficiently small relative displacement,

$$g(x, y) = q + O\left(\frac{|\xi| + y}{x_f}\right).$$

The logarithmic-tail equation therefore has leading balance

$$\frac{dX}{d\ell} = \frac{\rho h_f - qy}{\delta} + \text{higher-order terms}.$$

The qy contribution reproduces the zero-recruitment displacement by , while the recruitment term integrates to a logarithm. Hence

$$\boxed{X(y; \rho) = x_f + by + \rho a \log \frac{C}{y} + \text{higher-order terms},}$$

where $C > 0$ is fixed by matching to the regular outer trajectory.

10.2 Action reduction

At the anchor,

$$A'_{\theta, \psi}(x_f) = \frac{1 - q}{h_f} = \frac{1}{a}.$$

Therefore, to first order,

$$A_{\theta, \psi}(X) - A_{\theta, \psi}(x_f) = \frac{X - x_f}{a} + \dots = \rho \log \frac{C}{y} + \frac{b}{a} y + \dots.$$

Thus

$$\mathcal{C}_\rho(y) = y \exp \left\{ \frac{A_{\theta, \psi}(X) - A_{\theta, \psi}(x_f)}{\rho} \right\} = C \exp \left\{ \frac{b}{a} \frac{y}{\rho} + \dots \right\}.$$

In the strict overlap

$$y \ll \rho,$$

we obtain

$$\boxed{\mathcal{C}_\rho(y) \rightarrow C.}$$

Thus the logarithmic matching amplitude C is the leading small- ρ limit of the nonlinear action plateau.

10.3 Second-order interpretation

The local inverse expansion may be written

$$X(y; \rho) = X_0(y) + \rho X_1(y) + \rho^2 X_2(y) + \cdots,$$

with

$$X_1(y) = a \log \frac{C}{y} + O(y \log y).$$

Define

$$r = \frac{\eta q}{x_f}, \quad c = \delta h'_f + r h_f, \quad \kappa = \frac{h_f c}{2\delta^3}.$$

The second inverse coefficient has the local form

$$X_2(y) = \kappa \log^2 \frac{C}{y} + D + O(y \log^2 y),$$

where D is a global finite part fixed by the regular trajectory.

The action derivatives at x_f are

$$A'_{\theta,\psi}(x_f) = \frac{1}{a}, \quad A''_{\theta,\psi}(x_f) = -\frac{c}{h_f^2}.$$

The quadratic logarithms cancel because

$$\frac{\kappa}{a} = \frac{c}{2\delta^2}, \quad \frac{1}{2} A''_{\theta,\psi}(x_f) a^2 = -\frac{c}{2\delta^2}.$$

Consequently, in the stricter overlap in which explicit y -dependent quadratic terms are negligible,

$$\boxed{A_{\theta,\psi}(X) - A_{\theta,\psi}(x_f) = \rho \log \frac{C}{y} + \rho^2 \frac{D}{a} + o(\rho^2).}$$

It follows that

$$\boxed{\mathcal{C}_\rho = C \exp \left(\rho \frac{D}{a} + o(\rho) \right).}$$

Thus the nonlinear plateau contains the first and second logarithmic matching constants as the first two terms in its small- ρ expansion, while retaining the finite-prevalence tail nonlinearly when that local expansion is not well ordered.

11 Special parameter limits

11.1 The case $\psi = 0$

When $\psi = 0$, $\eta = 1$ and

$$g(x, y) = R_0 x$$

is independent of y . The zero-recruitment geometry reduces to

$$n_0(x) = 1 + \frac{1}{R_0} \log x, \quad x_* = \frac{1}{R_0}.$$

The nonlinear tail equation becomes

$$\boxed{\frac{dX}{d\ell} = \frac{\rho h_\theta(X) - R_0 X y}{1 - R_0 X}.$$

The action-drift identity simplifies to

$$\boxed{\dot{J} = -(1 - R_0 x)d.}$$

The entire finite-prevalence transmission term $g - g_0$ vanishes exactly. Correspondingly,

$$\varepsilon_A = \frac{|1 - R_0 x|d}{1 - R_0 x} = d$$

on the descending branch. This is the structural reason that the $\psi = 0$ case possesses a particularly broad action overlap: only susceptible feedback must become small.

In the local limit, the nonlinear tail reduces to the standard $\psi = 0$ logarithmic corner and the plateau amplitude reduces to the corresponding matching constant C_θ .

11.2 Fixed $0 < \psi < 1$

For positive ψ the finite-prevalence correction is present:

$$g = g_0(1 + s)^{-\psi}.$$

No assumption on s is required during the nonlinear deterministic tail. The small- s assumption enters only at the stochastic handoff. Consequently a trajectory may pass through a substantial interval with $y/x = O(1)$ and still be propagated deterministically without a transmission Taylor expansion.

11.3 The endpoint $\psi \rightarrow 1^-$

For fixed positive (x, y) , the exact deterministic transmission factor has the finite limit

$$g(x, y) = R_0 x(x + y)^{-\psi} \longrightarrow R_0 \frac{x}{x + y}.$$

Likewise, in (n, p) coordinates,

$$g = R_0 p n^\eta \rightarrow R_0 p.$$

Thus the nonlinear deterministic tail equation itself has a regular pointwise limit.

The full burnout asymptotics are nevertheless nonuniform because

$$x_* = R_0^{-1/\eta} \rightarrow 0 \quad (\eta \rightarrow 0^+),$$

and

$$x_f = z_f^{1/\eta} \rightarrow 0.$$

At exactly $\psi = 1$, the low-prevalence rate for fixed $x > 0$ is

$$g_0(x) = R_0,$$

so there is no positive interior solution of $g_0(x) = 1$ when $R_0 > 1$. The interior Laplace saddle therefore disappears into the boundary. A theory uniform all the way to $\psi = 1$ would require a separate joint scaling in η and the small host fraction. The present theory is consequently stated for fixed $0 \leq \psi < 1$.

12 Validity conditions and identifiable failure modes

12.1 Regular outer validity

The outer expansion is used only at a fixed point $\bar{x}$ away from the zero-recruitment endpoint and the low-prevalence saddle. It requires

$$\rho \ll 1$$

and bounded outer coefficient corrections, for example

$$|\rho Y_1(\bar{x})| \ll \bar{y}, \quad |\rho^2 Y_2(\bar{x})| \ll |\rho Y_1(\bar{x})|$$

when a strict second-order hierarchy is desired.

12.2 Descending inverse-flow validity

The scalar nonlinear continuation requires

$$y > 0, \quad g < 1,$$

so that y is monotone decreasing. It is stopped before the first trough $g = 1$. No condition $y/x \ll 1$ is imposed during this stage.

12.3 Deterministic-to-stochastic overlap

A clean handoff requires a nonempty interval on which

$$Ky \gg 1, \quad s = \frac{y}{x} \ll 1, \quad d = \frac{gy}{\rho h_\theta(x)} \ll 1.$$

For action-plateau independence one also requires the accumulated drift to be small:

$$\int \varepsilon_A d\ell \ll 1$$

over the range of candidate handoff levels.

Failure to find such an interval is not a failure of the nonlinear deterministic tail. It indicates that, at the specified finite K , stochasticity becomes important before the deterministic trajectory enters a clean low-prevalence, recruitment-dominated overlap.

12.4 Near-threshold loss of saddle separation

Laplace's method requires

$$\Lambda_h = \frac{A_{\theta,\psi}(x_*) - A_{\theta,\psi}(x_h)}{\rho} \gg 1,$$

and, equivalently in the fixed-parameter regime,

$$x_* - x_h \gg \sqrt{\rho}.$$

If the handoff lies too close to the saddle, the exact Kendall integral should be retained.

12.5 Large R_0 and positive ψ

At fixed R_0 , z_f is independent of ψ , while

$$x_f = z_f^{1/(1-\psi)}.$$

Thus positive ψ can make the post-wave susceptible fraction extremely small. The nonlinear tail remains applicable because it retains $(1 + y/x)^{-\psi}$ exactly. However, the branching handoff still requires eventual entry into $y/x \ll 1$. At finite K this may force the handoff to very low prevalence, potentially conflicting with $Ky \gg 1$. The relevant diagnostic is therefore the existence of the overlap itself, not the smallness of x_f alone.

12.6 The singular endpoint $\psi = 1$

The deterministic inverse-flow representation has a limit at $\psi = 1$, but the interior threshold x_* is lost. The Kendall/Laplace theory derived here therefore does not extend uniformly to $\psi = 1$.

13 Computational workflow

13.1 Precomputation for fixed (R_0, θ, ψ)

1. Set

$$\eta = 1 - \psi, \quad h_\theta(x) = (1 - x)x^\theta.$$

2. Solve

$$z_f = 1 + \frac{1}{R_0} \log z_f$$

for the nontrivial root $z_f \in (0, 1)$ and set

$$x_f = z_f^{1/\eta}.$$

3. Compute

$$x_* = R_0^{-1/\eta}, \quad h_* = h_\theta(x_*), \quad \Delta A_f = \int_{x_f}^{x_*} \frac{1 - R_0 z^\eta}{h_\theta(z)} dz.$$

4. Choose several test values of $\bar{x} \in (x_f, x_*)$ away from the endpoints and compute

$$\bar{y} = F_\psi(\bar{x}).$$

5. Integrate the regular outer equations for Y_1, Y_2 from $x = 1$ to $\bar{x}$ and convert to $\bar{X}_1, \bar{X}_2$.

13.2 Loop over ρ

For each ρ :

1. Initialize

$$X(0) = \bar{x} + \rho \bar{X}_1 + \rho^2 \bar{X}_2.$$

2. Integrate

$$\frac{dX}{d\ell} = \frac{\rho h_\theta(X) - g(X, y)y}{1 - g(X, y)}, \quad y = \bar{y}e^{-\ell},$$

down the descending tail.

3. Along the trajectory record

$$s = \frac{y}{X}, \quad d = \frac{gy}{\rho h_\theta(X)}, \quad \varepsilon_A = \frac{\psi g_0 s + |1 - g_0|d}{1 - g},$$

$$\mathcal{C}_\rho(y) = y \exp \left\{ \frac{A_{\theta, \psi}(X) - A_{\theta, \psi}(x_f)}{\rho} \right\}.$$

4. Verify that results are insensitive to the chosen outer initialization point $\bar{x}$.

13.3 Loop over K

For each population size K :

1. Find candidate handoff points satisfying

$$Ky \gg 1, \quad s \ll 1, \quad d \ll 1, \quad \varepsilon_A \ll 1.$$

2. Require $\mathcal{C}_\rho(y)$ to be approximately flat across a range of such points. Use the plateau range to quantify deterministic handoff sensitivity.

3. For a representative handoff (x_h, y_h) compute the exact Kendall integral

$$\mathcal{I}(x_h) = \int_{x_h}^1 \frac{1}{\rho h_\theta(X)} \exp \left\{ \frac{A_{\theta, \psi}(X) - A_{\theta, \psi}(x_h)}{\rho} \right\} dX.$$

4. Compute

$$B = Ky_h \log(1 + 1/\mathcal{I}), \quad P_{\text{persist}|\text{est}} = 1 - e^{-B}.$$

5. If $\Lambda_h \gg 1$, also compute

$$B_{\text{NL}} = KC_{\rho, h} \sqrt{\frac{\eta \rho h_*}{2\pi x_*}} \exp \left(-\frac{\Delta A_f}{\rho} \right)$$

as the fully asymptotic predictor.

14 Verification targets

A production implementation should retain the following checks.

14.1 Zero-recruitment checks

Verify numerically that

$$F_\psi(x_f) = 0, \quad x_f^\eta = z_f, \quad z_f = 1 + \frac{1}{R_0} \log z_f, \quad x_* = R_0^{-1/\eta}.$$

14.2 Outer-initialization independence

Repeat the calculation for several $\bar{x}$ values. The nonlinear tail and final persistence prediction should agree to the order of the regular outer truncation.

14.3 Exact deterministic regression

Because the nonlinear inverse flow is mathematically equivalent to the two-dimensional deterministic ODE on the descending branch, compare $X(y)$ against a direct deterministic trajectory. Any discrepancy larger than the expected outer initialization error indicates an implementation error.

14.4 Action plateau

Plot

$$\log \mathcal{C}_\rho(y) = \log y + \frac{A_{\theta,\psi}(X(y)) - A_{\theta,\psi}(x_f)}{\rho}$$

against $\ell = \log(\bar{y}/y)$ together with s , d , and ε_A . The plateau should emerge where the overlap diagnostics become small.

14.5 Handoff invariance

Evaluate the exact Kendall prediction at several handoff points inside the admissible plateau. The variation is a direct finite-parameter measure of handoff error.

14.6 $\psi = 0$ regression

At $\psi = 0$, verify

$$g = R_0 x$$

for all y and confirm that the finite-prevalence term in the action drift is identically zero. The nonlinear theory should reduce to the corresponding $\psi = 0$ prediction as $\rho \rightarrow 0$.

14.7 Positive- ψ stress test

For large R_0 and positive ψ , record

$$\frac{y}{x}$$

along the tail. The nonlinear continuation should remain stable through regions where this ratio is $O(1)$, while the stochastic handoff should be delayed until a genuine low-prevalence overlap is reached.

15 Recruitment-law derivatives

For

$$h_\theta(x) = (1 - x)x^\theta,$$

we have

$$h'_\theta(x) = x^{\theta-1}\{\theta - (\theta + 1)x\}.$$

For $x > 0$,

$$h''_\theta(x) = \theta x^{\theta-2}\{(\theta - 1) - (\theta + 1)x\}.$$

These formulas enter the next Laplace coefficient and local asymptotic reductions.

16 Detailed derivation of the regular outer coefficient equations

The compact formulas in Section 4 follow most transparently from the perturbed zero-recruitment invariant. This derivation is included so that the regular initialization is completely reproducible.

16.1 Exact drift of the zero-recruitment invariant

Recall

$$H(x, y) = n^\eta - \frac{\eta}{R_0} \log x, \quad n = x + y.$$

For the full system,

$$\dot{n} = \rho h_\theta(x) - y.$$

Hence

$$\dot{H} = \eta n^{-\psi}(\rho h_\theta - y) - \frac{\eta}{R_0 x}(\rho h_\theta - gy).$$

Using

$$\frac{g}{R_0 x} = n^{-\psi},$$

the two terms proportional to y cancel exactly, leaving

$$\dot{H} = \eta \rho h_\theta(x) \left[n^{-\psi} - \frac{1}{R_0 x} \right].$$

Dividing by

$$\dot{x} = \rho h_\theta(x) - R_0 x y n^{-\psi}$$

gives the exact phase derivative

$$\boxed{\frac{dH}{dx} = \eta \rho h_\theta(x) \frac{n^{-\psi} - 1/(R_0 x)}{\rho h_\theta(x) - R_0 x y n^{-\psi}}.}$$

16.2 Expansion of the invariant itself

On the regular outer interval set

$$y = F_\psi + \rho Y_1 + \rho^2 Y_2 + O(\rho^3), \quad n = n_0 + \rho Y_1 + \rho^2 Y_2 + O(\rho^3).$$

Taylor expansion of n^η gives

$$\begin{aligned} n^\eta &= n_0^\eta + \eta n_0^{-\psi} (\rho Y_1 + \rho^2 Y_2) + \frac{\eta(\eta-1)}{2} n_0^{\eta-2} \rho^2 Y_1^2 + O(\rho^3) \\ &= n_0^\eta + \eta \rho n_0^{-\psi} Y_1 + \eta \rho^2 \left[n_0^{-\psi} Y_2 - \frac{\psi}{2} n_0^{-\psi-1} Y_1^2 \right] + O(\rho^3), \end{aligned}$$

because $\eta - 1 = -\psi$. Since

$$n_0^\eta - \frac{\eta}{R_0} \log x = 1$$

on the zero-recruitment orbit, define

$$W_1 = n_0^{-\psi} Y_1, \quad W_2 = n_0^{-\psi} Y_2 - \frac{\psi}{2} n_0^{-\psi-1} Y_1^2.$$

Then

$$\boxed{H = 1 + \eta \rho W_1 + \eta \rho^2 W_2 + O(\rho^3),}$$

and therefore

$$\frac{dH}{dx} = \eta \rho W_1' + \eta \rho^2 W_2' + O(\rho^3).$$

16.3 First outer coefficient

Write

$$N_0 = n_0^{-\psi}, \quad B_0 = N_0 - \frac{1}{R_0 x}, \quad A_0 = R_0 x F_\psi N_0.$$

At leading order in the exact formula for H_x ,

$$\frac{1}{\eta \rho} \frac{dH}{dx} = h_\theta(x) \frac{B_0}{-A_0} + O(\rho).$$

Now

$$B_0 = \frac{R_0 x - n_0^\psi}{R_0 x n_0^\psi}, \quad A_0 = \frac{R_0 x F_\psi}{n_0^\psi}.$$

Thus

$$h_\theta \frac{B_0}{-A_0} = \frac{h_\theta(n_0^\psi - R_0 x)}{R_0^2 x^2 F_\psi}.$$

Using

$$F'_\psi(x) = \frac{n_0^\psi}{R_0 x} - 1 = \frac{n_0^\psi - R_0 x}{R_0 x},$$

we obtain

$$W'_1 = \frac{h_\theta(x) F'_\psi(x)}{R_0 x F_\psi(x)}.$$

16.4 Second outer coefficient

To one further order,

$$n^{-\psi} = N_0 + \rho N_1 + O(\rho^2), \quad N_1 = -\psi n_0^{-\psi-1} Y_1.$$

Also

$$R_0 x y n^{-\psi} = A_0 + \rho A_1 + O(\rho^2),$$

with

$$A_1 = R_0 x \{Y_1 N_0 + F_\psi N_1\} = R_0 x N_0 Y_1 \left(1 - \psi \frac{F_\psi}{n_0}\right).$$

The numerator factor is

$$n^{-\psi} - \frac{1}{R_0 x} = B_0 + \rho N_1 + O(\rho^2),$$

while the denominator is

$$\rho h_\theta - R_0 x y n^{-\psi} = -A_0 + \rho(h_\theta - A_1) + O(\rho^2).$$

Therefore

$$\frac{1}{-A_0 + \rho(h_\theta - A_1)} = -\frac{1}{A_0} \left[1 + \rho \frac{h_\theta - A_1}{A_0}\right] + O(\rho^2).$$

Multiplication gives

$$W'_2 = -\frac{h_\theta}{A_0} \left[N_1 + B_0 \frac{h_\theta - A_1}{A_0}\right].$$

Substituting A_0, A_1, B_0, N_1 and simplifying yields

$$W'_2 = \frac{h_\theta(x)}{R_0^2 x^2 F_\psi(x)^2} \left[\psi n_0(x)^{\psi-1} F_\psi(x) - \{n_0(x)^\psi - R_0 x\} \right] Y_1(x) + \frac{h_\theta(x)^2 n_0(x)^\psi \{n_0(x)^\psi - R_0 x\}}{R_0^3 x^3 F_\psi(x)^2}.$$

Finally,

$$Y_1 = n_0^\psi W_1, \quad Y_2 = n_0^\psi W_2 + \frac{\psi}{2n_0} Y_1^2,$$

which completes the regular outer derivation.

17 Detailed local inverse hierarchy and finite-part constants

The nonlinear tail does not require a local expansion about x_f . Nevertheless, the local hierarchy is useful because it proves the reduction stated in Section 10 and identifies the asymptotic coefficients contained in the plateau amplitude.

17.1 Exact inverse-flow hierarchy

Write

$$\frac{dx}{dy} = \Phi(x, y) + \rho\Theta(x, y),$$

where

$$\Phi(x, y) = -\frac{g(x, y)}{g(x, y) - 1}, \quad \Theta(x, y) = \frac{h_\theta(x)}{\{g(x, y) - 1\}y}.$$

At fixed y , expand

$$X(y; \rho) = X_0(y) + \rho X_1(y) + \rho^2 X_2(y) + O(\rho^3).$$

Taylor expansion of the right-hand side about X_0 gives

$$\begin{aligned} \Phi(X, y) + \rho\Theta(X, y) &= \Phi(X_0, y) \\ &\quad + \rho\{\Phi_x(X_0, y)X_1 + \Theta(X_0, y)\} \\ &\quad + \rho^2\left\{\Phi_x(X_0, y)X_2 + \frac{1}{2}\Phi_{xx}(X_0, y)X_1^2 + \Theta_x(X_0, y)X_1\right\} + O(\rho^3). \end{aligned}$$

Equating coefficients gives

$$\boxed{X'_0 = \Phi(X_0, y),}$$

$$\boxed{X'_1 = \Phi_x(X_0, y)X_1 + \Theta(X_0, y),}$$

$$\boxed{X'_2 = \Phi_x(X_0, y)X_2 + \frac{1}{2}\Phi_{xx}(X_0, y)X_1^2 + \Theta_x(X_0, y)X_1.}$$

The leading solution is the inverse zero-recruitment orbit,

$$F_\psi(X_0(y)) = y.$$

17.2 Analytic normal form at the anchor

Define

$$\Psi(x, y) = \frac{h_\theta(x)}{g(x, y) - 1}.$$

Then

$$\Theta = \frac{\Psi}{y}, \quad \Theta_x = \frac{\Psi_x}{y}.$$

At $(x_f, 0)$,

$$g(x_f, 0) = q < 1,$$

so $g - 1$ is nonzero and Φ, Ψ are analytic in a neighborhood of the anchor. Because

$$F'_\psi(x_f) = \frac{\delta}{q} > 0,$$

the inverse-function theorem gives an analytic $X_0(y)$ near $y = 0$.

Define

$$\varphi(y) = \Phi_x(X_0(y), y), \quad \chi(y) = \frac{1}{2}\Phi_{xx}(X_0(y), y),$$

$$\tilde{\Psi}(y) = \Psi(X_0(y), y), \quad \omega(y) = \Psi_x(X_0(y), y).$$

Then

$$X_1' = \varphi X_1 + \frac{\tilde{\Psi}}{y},$$

$$X_2' = \varphi X_2 + \chi X_1^2 + \frac{\omega}{y} X_1.$$

All four coefficient functions are analytic at $y = 0$.

The required local values follow directly from

$$g_x = g \left(\frac{1}{x} - \frac{\psi}{x+y} \right).$$

At the anchor,

$$g_x(x_f, 0) = \frac{\eta q}{x_f} =: r.$$

Since

$$\Phi_x = \frac{g_x}{(g-1)^2},$$

we have

$$\varphi(0) = \frac{r}{\delta^2}.$$

Also

$$\tilde{\Psi}(0) = -\frac{h_f}{\delta} = -a.$$

Differentiating Ψ gives

$$\Psi_x = \frac{h'(g-1) - h g_x}{(g-1)^2}.$$

Therefore

$$\omega(0) = -\frac{\delta h'_f + r h_f}{\delta^2} = -\frac{c}{\delta^2}.$$

17.3 First coefficient and the logarithmic amplitude

Let

$$E(y) = \exp \left\{ \int_0^y \varphi(t) dt \right\}.$$

Then $E(0) = 1$ and

$$\left(\frac{X_1}{E} \right)' = \frac{\tilde{\Psi}(y)}{E(y)y}.$$

Set

$$\mu(y) = \frac{\tilde{\Psi}(y)}{E(y)}.$$

Because μ is analytic and $\mu(0) = -a$,

$$\frac{\mu(y)}{y} = -\frac{a}{y} + \frac{\mu(y) + a}{y},$$

and the second term is regular at zero. Integration therefore gives

$$\frac{X_1(y)}{E(y)} = -a \log y + C_1 + O(y).$$

Since $E(y) = 1 + O(y)$,

$$X_1(y) = -a \log y + C_1 + O(y \log y).$$

Writing

$$C_1 = a \log C$$

defines the positive global matching amplitude C and gives

$$X_1(y) = a \log \frac{C}{y} + O(y \log y).$$

Equivalently,

$$a \log C = \lim_{y \downarrow 0} \{X_1(y) + a \log y\}.$$

17.4 Second coefficient and the finite constant

The second equation is a linear ODE in X_2 :

$$X_2' - \varphi X_2 = \chi X_1^2 + \frac{\omega}{y} X_1.$$

Using the same integrating factor,

$$\left(\frac{X_2}{E} \right)' = \frac{\chi X_1^2}{E} + \frac{\omega X_1}{Ey}.$$

Since

$$X_1 = aL + O(yL), \quad L = \log \frac{C}{y},$$

the first forcing term is $O(L^2)$ and is integrable at zero. The only nonintegrable contribution comes from the second term:

$$\frac{\omega X_1}{Ey} = -\frac{ac}{\delta^2} \frac{L}{y} + O(L).$$

Define

$$\kappa = \frac{ac}{2\delta^2} = \frac{h_f c}{2\delta^3}.$$

Because

$$\frac{d}{dy}(\kappa L^2) = -\frac{2\kappa L}{y} = -\frac{ac}{\delta^2} \frac{L}{y},$$

the complete nonintegrable singularity is the derivative of κL^2 . Hence

$$X_2(y) = \kappa L^2 + D + O(yL^2),$$

where

$$D = \lim_{y \downarrow 0} \left\{ X_2(y) - \kappa \log^2 \frac{C}{y} \right\}.$$

No independent single-log constant occurs at this order: the coefficient of the possible single logarithm is tied to the first-order finite part through the same singular forcing.

For numerical evaluation define

$$Z_1 = X_1 + a \log y, \quad Z_2 = X_2 - \kappa \log^2 \frac{C}{y}.$$

Their derivatives are integrable at zero, so for any regular $\bar{y} > 0$,

$$Z_j(0^+) = Z_j(\bar{y}) - \int_0^{\bar{y}} Z'_j(y) dy, \quad j = 1, 2.$$

Thus C and D are global finite parts determined by convergent one-dimensional quadratures.

17.5 Action cancellation at second order

At the anchor,

$$A'_{\theta,\psi}(x_f) = \frac{1}{a}.$$

Differentiating once more and using $g'_0(x_f) = r$ gives

$$A''_{\theta,\psi}(x_f) = -\frac{c}{h_f^2}.$$

In the strict local overlap, the logarithmic part of the displacement is

$$X - x_f = \rho a L + \rho^2(\kappa L^2 + D) + \dots.$$

Taylor expansion of the action gives

$$\begin{aligned} A_{\theta,\psi}(X) - A_{\theta,\psi}(x_f) &= \frac{1}{a} \{ \rho a L + \rho^2(\kappa L^2 + D) \} + \frac{1}{2} A''_{\theta,\psi}(x_f) \rho^2 a^2 L^2 + \dots \\ &= \rho L + \rho^2 \frac{D}{a} + \rho^2 L^2 \left\{ \frac{\kappa}{a} + \frac{1}{2} A''_{\theta,\psi}(x_f) a^2 \right\} + \dots. \end{aligned}$$

Now

$$\frac{\kappa}{a} = \frac{c}{2\delta^2}, \quad \frac{1}{2} A''_{\theta,\psi}(x_f) a^2 = -\frac{c}{2\delta^2},$$

so the squared logarithm cancels exactly. Therefore

$$A_{\theta,\psi}(X) - A_{\theta,\psi}(x_f) = \rho \log \frac{C}{y} + \rho^2 \frac{D}{a} + \dots,$$

which proves the second-order plateau expansion used in Section 10.

18 Derivation of the Laplace correction

Write the Kendall integral as

$$\mathcal{I}(x_h) = \frac{1}{\rho} e^{-A_{\theta,\psi}(x_h)/\rho} \int_{x_h}^1 f(X) e^{A_{\theta,\psi}(X)/\rho} dX, \quad f(X) = \frac{1}{h_{\theta}(X)}.$$

Let

$$A_0 = A_{\theta,\psi}(x_*), \quad \lambda = -A''_{\theta,\psi}(x_*) > 0, \quad A_3 = A'''_{\theta,\psi}(x_*), \quad A_4 = A''''_{\theta,\psi}(x_*).$$

Set $z = X - x_*$. Near the saddle,

$$A_{\theta,\psi}(X) = A_0 - \frac{\lambda}{2} z^2 + \frac{A_3}{6} z^3 + \frac{A_4}{24} z^4 + O(z^5),$$

$$f(X) = f_0 + f_1 z + \frac{f_2}{2} z^2 + O(z^3).$$

Scale

$$z = \sqrt{\frac{\rho}{\lambda}} t.$$

Expanding the exponential and retaining terms whose Gaussian averages are $O(\rho)$ gives

$$\begin{aligned} \int f(X) e^{A_{\theta,\psi}(X)/\rho} dX &\sim e^{A_0/\rho} \sqrt{\frac{2\pi\rho}{\lambda}} f_0 \\ &\times \left[1 + \rho \left(\frac{f_2}{2f_0\lambda} + \frac{f_1 A_3}{2f_0\lambda^2} + \frac{A_4}{8\lambda^2} + \frac{5A_3^2}{24\lambda^3} \right) + O(\rho^2) \right]. \end{aligned}$$

Here the numerical coefficients follow from the Gaussian moments

$$\langle t^2 \rangle = 1, \quad \langle t^4 \rangle = 3, \quad \langle t^6 \rangle = 15.$$

Therefore

$$c_L = \frac{f_2}{2f_0\lambda} + \frac{f_1 A_3}{2f_0\lambda^2} + \frac{A_4}{8\lambda^2} + \frac{5A_3^2}{24\lambda^3}.$$

For $f = 1/h_{\theta}$, write $h = h_{\theta}$ temporarily and define

$$r(x) = 1 - R_0 x^{\eta}, \quad A'_{\theta,\psi}(x) = \frac{r(x)}{h(x)}.$$

At $x = x_*$ we have $r(x_*) = 0$ and $R_0 x_*^{\eta} = 1$. The derivatives of r at the saddle are

$$r'_* = -\frac{\eta}{x_*}, \quad r''_* = -\frac{\eta(\eta-1)}{x_*^2}, \quad r'''_* = -\frac{\eta(\eta-1)(\eta-2)}{x_*^3}.$$

Consequently

$$\begin{aligned} \lambda &= -A''_{\theta,\psi}(x_*) = \frac{\eta}{x_* h_*}, \\ A_3 &= -\frac{\eta(\eta-1)}{x_*^2 h_*} + \frac{2\eta h'_*}{x_* h_*^2}, \end{aligned}$$

and, using

$$\left(\frac{1}{h}\right)' = -\frac{h'}{h^2}, \quad \left(\frac{1}{h}\right)'' = \frac{2(h')^2}{h^3} - \frac{h''}{h^2},$$

$$A_4 = -\frac{\eta(\eta-1)(\eta-2)}{x_*^3 h_*} + \frac{3\eta(\eta-1)h'_*}{x_*^2 h_*^2} - \frac{6\eta(h'_*)^2}{x_* h_*^3} + \frac{3\eta h_*''}{x_* h_*^2}.$$

For the prefactor,

$$f_0 = \frac{1}{h_*}, \quad f_1 = -\frac{h'_*}{h_*^2}, \quad f_2 = \frac{2(h'_*)^2}{h_*^3} - \frac{h_*''}{h_*^2}.$$

Substituting these expressions into

$$c_L = \frac{f_2}{2f_0\lambda} + \frac{f_1 A_3}{2f_0\lambda^2} + \frac{A_4}{8\lambda^2} + \frac{5A_3^2}{24\lambda^3}$$

gives the four contributions

$$\frac{f_2}{2f_0\lambda} = \frac{x_*}{\eta} \left\{ \frac{(h'_*)^2}{h_*} - \frac{1}{2}h_*'' \right\},$$

$$\frac{f_1 A_3}{2f_0\lambda^2} = \frac{\eta-1}{2\eta} h'_* - \frac{x_*(h'_*)^2}{\eta h_*},$$

$$\frac{A_4}{8\lambda^2} = -\frac{(\eta-1)(\eta-2)h_*}{8\eta x_*} + \frac{3(\eta-1)h'_*}{8\eta} - \frac{3x_*(h'_*)^2}{4\eta h_*} + \frac{3x_* h_*''}{8\eta},$$

$$\frac{5A_3^2}{24\lambda^3} = \frac{5(\eta-1)^2 h_*}{24\eta x_*} - \frac{5(\eta-1)h'_*}{6\eta} + \frac{5x_*(h'_*)^2}{6\eta h_*}.$$

Collecting the coefficients of h_*/x_* , h'_* , $x_*(h'_*)^2/h_*$, and $x_* h_*''$ yields

$$c_L = \frac{(2\eta^2 - \eta - 1)h_*}{24\eta x_*} + \frac{(\eta-1)h'_*}{24\eta} + \frac{x_*(h'_*)^2}{12\eta h_*} - \frac{x_* h_*''}{8\eta}.$$


Putting these terms over the common denominator $24\eta h_* x_*$ gives

$$c_L = \frac{(2\eta^2 - \eta - 1)h_*^2 + (\eta-1)x_* h_* h'_* + 2x_*^2 (h'_*)^2 - 3x_*^2 h_* h_*''}{24\eta h_* x_*}.$$

Thus every algebraic step from the general Gaussian correction to the closed coefficient used in Section 9 is explicit.

19 Summary of the theoretical chain

The theory can be summarized as



The key structural separation is that the deterministic tail is allowed to pass through finite relative prevalence $y/x = O(1)$ without expanding the transmission factor. The condition $y/x \ll 1$ is imposed only where it is actually needed: at the deterministic-to-stochastic handoff. The action identity then supplies both a quantitative overlap diagnostic and the handoff-independent amplitude that enters the leading Laplace burnout exponent.