


Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.01$, $\theta = 0$; $l_0 = 1$; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.01, theta = 0.5; I0 = 1; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.01, theta = 1; l0 = 1; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.02, theta = 0; l0 = 1; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.02, theta = 0.5; I0 = 1; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.02$, $\theta = 1$; $l_0 = 1$; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.05, theta = 0; l0 = 1; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.05, theta = 0.5; I0 = 1; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.05, theta = 1; l0 = 1; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; $\rho = 0.10$, $\theta = 0$; $l_0 = 1$; matching height = $2/3$ compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height $2/3$ compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.10, theta = 0.5; I0 = 1; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference


Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.

Full stochastic validation of unconditional persistence

Exact Gillespie CTMC; rho = 0.10, theta = 1; l0 = 1; matching height = 2/3 compromise



Analytical method

- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry
- Early-establishment reference

Matching height

- $y_{BL} = K^{-1/3} y_{star}^{2/3}$
- Early-establishment reference

Points and 95% Wilson intervals are stochastic estimates. All analytical curves use matching height 2/3 compromise. Each curve stops where its matching equation has no admissible root.