


Unconditional persistence probability

$\rho = 0.01$, $\theta = 0$, $l_0 = 1$; matching height = $\sqrt{y^*/K}$



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = \sqrt{y_{star}/K}$

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. All analytical curves use matching height $\sqrt{y^*/K}$. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Persistence probability conditional on escaping early fizzle

$\rho = 0.01$, $\theta = 0$, $l_0 = 1$; matching height = $\sqrt{y^*/K}$



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = \sqrt{y_{star}/K}$

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. All analytical curves use matching height $\sqrt{y^*/K}$. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Unconditional persistence probability

$\rho = 0.01$, $\theta = 0.5$, $l_0 = 1$; matching height = $\sqrt{y^*/K}$



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = \sqrt{y_{star}/K}$

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. All analytical curves use matching height $\sqrt{y^*/K}$. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Persistence probability conditional on escaping early fizzle

$\rho = 0.01$, $\theta = 0.5$, $l_0 = 1$; matching height = $\sqrt{y^*/K}$



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = \sqrt{y_{star}/K}$

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. All analytical curves use matching height $\sqrt{y^*/K}$. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Unconditional persistence probability

$\rho = 0.01$, $\theta = 1$, $l_0 = 1$; matching height = $\sqrt{y^*/K}$



Analytical method


- Exact Kendall + first-order entry
- Exact Kendall + second-order entry
- Laplace + first-order entry
- Laplace + second-order entry

Matching height $y_{BL} = \sqrt{y_{star}/K}$

Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. All analytical curves use matching height $\sqrt{y^*/K}$. Curves use the analytical D_θ expression. Missing curve segments indicate no admissible matching root.

Persistence probability conditional on escaping early fizzle

$\rho = 0.01$, $\theta = 1$, $l_0 = 1$; matching height = $\sqrt{y^*/K}$



Points and 95% Wilson intervals: adaptive tau-leaping (epsilon = 0.01); 3,000 attempts per point, expanded to 10,000 when needed. All analytical curves use matching height $\sqrt{y^*/K}$. Curves use the analytical D_{θ} expression. Missing curve segments indicate no admissible matching root.