

Bridging the gap between theory and data: the Red Queen

Hypothesis for sex

Abstract

Sexual reproduction persists in nature despite its large cost. The Red Queen Hypothesis postulates that parasite pressure maintains sexual reproduction in the host population by selecting for the ability to produce rare genotypes that are resistant to infection. Mathematical models have been used to lay theoretical foundations for the hypothesis; empirical studies have confirmed these predictions. For example, Lively used a simple host-parasite model to predict that the frequency of sexual hosts should be positively correlated with the prevalence of infection. Lively *et al.* later confirmed the prediction through numerous field studies of snail-trematode systems in New Zealand. In this study, we fit a simple metapopulation host-parasite coevolution model to three data sets, each representing a different snail-trematode system, by matching the observed prevalence of sexual reproduction and trematode infection among hosts. Using the estimated parameters, we perform a power analysis to test the feasibility of observing the positive correlation predicted by Lively. We discuss anomalies in the data that are poorly explained by the model and provide practical guidance to both modelers and empiricists. Overall, our study suggests that a simple Red Queen model can only partially explain the observed relationships between parasite infection and the maintenance of sexual reproduction.

1 Introduction

Despite being the dominant mode of reproduction in multicellular organisms (Vrijenhoek 1998, Whitton et al. 2008, Otto 2009), sexual reproduction entails numerous costs (Lehtonen et al. 2012). The most commonly mentioned is the cost of producing males (Smith 1978): as males cannot produce offspring, asexual lineages are expected to outgrow their sexual counterparts. This *two-fold cost of sex* (Smith 1978) relies on the assumption that everything else is equal. Then, what drives the observed prevalence of sexual reproduction?

One explanation for the persistence of sexual reproduction is the Red Queen Hypothesis (RQH) (Bell 1982). The RQH predicts that sexually reproducing hosts overcome the cost of sex under strong parasitic pressure by producing genetically diverse offspring that can escape infection (Haldane 1949, Jaenike 1978, Hamilton 1980, Hamilton et al. 1990). Limited genetic diversity of asexually reproducing hosts allows for sexual reproduction to be maintained in the host population (Ashby and King 2015).

Much of the theoretical literature has focused on determining qualitative conditions under which parasite selection can maintain sexual reproduction in the host population. Here, we describe a few important criteria. First, hosts and parasites must coevolve (Bell 1982). Host-parasite coevolution creates a time-lagged selective advantage for rare host genotypes, creating an oscillation in genotypic frequencies (Jaenike 1978, Hamilton 1980, Agrawal and Lively 2001); this process promotes host genetic diversity and not necessarily sex (King et al. 2009, Dagan et al. 2013a, Ashby and King 2015). Second, parasites must be highly virulent (May and Anderson 1983). Although sexual and asexual hosts can coexist at intermediate virulence, sexual reproduction is unable to provide enough advantage to overcome the cost of sex against avirulent parasites (Howard et al. 1994). Finally, sexual hosts must be genetically more diverse than asexual hosts, as high clonal diversity may diminish the advantage of sexual reproduction (Lively and Howard 1994, Lively 2010c, Ashby and King 2015).

Some theoretical studies have departed from the classical population genetics framework to study the effects of ecological and epidemiological context on the Red Queen dynamics. A few studies have suggested that incorporating more realistic ecological and epidemiological details may assist in supporting sexual

reproduction in the host population (Lively 2009, 2010b) and maintaining coevolutionary cycles (Ashby and Gupta 2014). In contrast, MacPherson and Otto (2018) showed that Red Queen dynamics (i.e., cycles in allele frequencies) fail to persist when an explicit epidemiological structure is taken into account with coevolutionary dynamics. Given the wide range of assumptions underlying these eco-evolutionary models, it may be difficult to understand how different ecological details of these models affect the overall maintenance of sexual reproduction; however, it is clear is that these details play important roles in shaping coevolutionary dynamics (Song et al. 2015, Haafke et al. 2016, Ashby et al. 2019).

Many empirical studies have focused on confirming predictions that stem from the RQH. Typical among them are rapid parasite evolution (Rauch et al. 2006), local adaptation (Lively 1989, Morran et al. 2014, King et al. 2011, Gibson et al. 2016), time-lagged selection (Buckling and Rainey 2002, Decaestecker et al. 2007, Koskella and Lively 2009, Thrall et al. 2012, Koskella 2013, 2014), and association between parasite prevalence and host reproductive mode (Lively 1992, Vergara et al. 2013, Verhoeven and Biere 2013). A key example is the snail population in New Zealand that serves as intermediate hosts for trematodes (Winterbourn 1974, McArthur and Featherston 1976). Through decades of work, Lively *et al.* have demonstrated that the population satisfies necessary conditions for the host-parasite coevolutionary dynamics and provides support for the hypothesis (e.g., Lively (1987, 1989), Dybdahl and Lively (1995b, 1998), Jokela et al. (2009), Vergara et al. (2014b), Gibson et al. (2016)). While field studies often provide only indirect evidence for the hypothesis, experimental systems can be used to directly test the hypothesis (Auld et al. 2016, Slowinski et al. 2016, Lynch et al. 2018, Zilio et al. 2018).

Even though the RQH has gained both theoretical and empirical support, there still remains a gap between theory and data. Many theoretical models rely on simplifying assumptions that are not applicable to natural populations and make indirect connections to empirical studies. For example, none of the Red Queen models reviewed by Ashby and King (2015) make statistical connections to empirical data. It is unclear how well these models capture coevolutionary dynamics in natural systems.

Theoretical models can still be used to make *qualitative* predictions about nature. Lively (1992) initially postulated that infection prevalence should be positively correlated with the frequency of sexual hosts and

later formalized the idea using a mathematical model (Lively 2001). The prediction has since been confirmed in several empirical studies, most of which are based on the snail-trematode system (Lively and Jokela 2002, Kumpulainen et al. 2004, King et al. 2011, Vergara et al. 2013, McKone et al. 2016, Gibson et al. 2016). Surprisingly, the predicted correlation was not observed in a different snail-trematode system (Heller and Farstey 1990, Ben-Ami and Heller 2005, 2007, 2008, Dagan et al. 2013a). Here, we try to help bridge the gap between theory and data in understanding the role of parasites in maintaining sexual reproduction in host populations and make quantitative inference about empirical systems. We extend the model used by Lively (2010b) to account for demographic stochasticity and include a simple metapopulation structure. Then, we fit the model to data sets from three field studies (Dagan et al. 2013a, McKone et al. 2016, Vergara et al. 2014b) using Approximate Bayesian Computation (ABC) to estimate biologically relevant parameters. We assess model fits and discuss discrepancies between a theoretical model and the observed data. Using biologically realistic parameters, we test for power (probability of observing a significant effect) to detect a positive correlation between frequency of sexual reproduction and prevalence of infection Lively (2001). We provide guidance for studying the RQH and discuss underlying factors that drive the correlation.

2 Material and Methods

2.1 Data

We analyze three observational data sets (Vergara et al. 2014b, McKone et al. 2016, Dagan et al. 2013a) that represent interactions between freshwater snails and sterilizing trematodes. The New Zealand snail populations (Vergara et al. 2014b, McKone et al. 2016) consist of *Potamopyrgus antipodarum*, which serves as an intermediate host for trematodes, among which *Microphallus* sp. is most commonly found (Winterbourn 1974, Lively 1987). The Israeli snail populations (Dagan et al. 2013a) consist of *Melanooides tuberculata*, which also serves as an intermediate host for trematodes, including *Centrocestus* sp. (Ben-Ami and Heller 2005), *Philophthalmus* sp. (Ben-Ami 2006), and *Transversotrema patialense* (Ben-Ami et al.

2005). Diploid sexual females and polyploid asexual females coexist in both the New Zealand (Phillips and Lambert 1989, Wallace 1992, Dybdahl and Lively 1995a) and the Israeli (Samadi et al. 1999) snail populations.

We choose to analyze snail-trematode systems because they have been extensively studied under the context of the RQH. The snail-trematode systems in New Zealand exhibit positive correlation between prevalence of infection and frequency of sexual hosts (Lively and Jokela 2002, King et al. 2011, Vergara et al. 2013, McKone et al. 2016, Gibson et al. 2016); local adaptation (Lively 1989, Lively et al. 2004, King et al. 2011); negative frequency-dependent selection (Dybdahl and Lively 1995b, 1998, Jokela et al. 2009, Koskella and Lively 2009); and periodic selection for sexual reproduction (Vergara et al. 2014b, Gibson et al. 2018). These results are consistent with the host-parasite coevolutionary theory behind the RQH. Therefore, we expect a basic Red Queen model to be able to mimic the observed dynamics reasonably well. We included the Israeli population because there has been a consistent lack of positive correlation between the prevalence of infection and frequency of sexual hosts (Heller and Farstey 1990, Ben-Ami and Heller 2005, 2007, 2008, Dagan et al. 2013a), which sharply contrasts the observations from the New Zealand population. Comparison of these two populations with a mathematical model may allow us to better understand the underlying cause of the difference.

The data sets include proportions of infected snails, proportions of sexual/asexual snails, and locations of sampling sites; the data set from Vergara et al. (2014b) also includes sampling year. Whereas Vergara et al. (2014b) report infection status of snails specific to *Microphallus* sp., McKone et al. (2016) and Dagan et al. (2013a) do not distinguish among different species. The data sets collected by Dagan et al. (2013a) and Vergara et al. (2014b) are obtained from their Dryad repositories (Dagan et al. 2013b, Vergara et al. 2014a). The data set collected by McKone et al. (2016) is extracted from their figure.

2.2 Model

We model obligately sexual hosts competing with obligately asexual hosts in a metapopulation by extending the model introduced by Lively (2010b). Our model is a discrete time susceptible-infected (SI)

model with natural mortality and virulence (defined as a reduction in offspring production among infected hosts). Metapopulation structure is included to model unobserved dynamics among different habitats; equivalently, each population can be considered as a sampling site. A similar metapopulation model was developed by Lively (2018) in order to study the local adaptation of parasites.

We do not model the life history of snail-trematode interactions explicitly. Modeling interactions of multiple trematodes species, each of which goes through a different life cycle, with a single snail species is extremely complicated. Instead, testing a basic model against data will allow us to identify the difference between theory and data more clearly.

All hosts are diploids with two biallelic loci controlling resistance; parasites are assumed to be haploids.

Let $S_{ij}^k(t)$ and $A_{ij}^k(t)$ be the number of sexual and asexual hosts with genotype ij from subpopulation k at generation t , where the subscripts, i and j , represents host haplotypes: $i, j \in \{AB, Ab, aB, ab\}$. For simplicity, we drop the superscript representing the subpopulation and write $S_{ij}(t)$ and $A_{ij}(t)$; every population is governed by the same set of equations unless noted otherwise (e.g., when we account for the interaction between populations). Following Lively (2010b), the expected genotypic contribution (before recombination or outcrossing) by sexual hosts is given by

$$S'_{ij} = c_b(1 - s) \left(W_U S_{ij,U}(t) + W_I S_{ij,I}(t) \right), \quad (1)$$

where s is the proportion of males produced by sexual hosts, and $S_{ij,U}$ and $S_{ij,I}$ are the number of uninfected and infected sexual hosts in a population. W_U and W_I represent their corresponding fitnesses where virulence is defined as $V = 1 - W_I/W_U$. We allow for the cost of sex to vary by multiplying the growth rate by a scale parameter, c_b , where $2/c_b$ corresponds to a two-fold cost of sex when $c_b = 1$ (Ashby and King 2015). Recombination and outcrossing are modeled after incorporating genotypic contributions from other populations.

We define

$$W_U = \frac{b_U}{1 + a_U N(t)}, \quad W_I = \frac{b_I}{1 + a_I N(t)}$$

where b_U and b_I are the maximum number of offspring produced by uninfected and infected hosts, respectively, and a_U and a_I determine their corresponding strengths of density dependence (Smith and

143 Slatkin 1973, Lively 2010b). For simplicity, we assume that $a_U = a_I$ so that virulence can be defined
 144 strictly in terms of decrease in offspring production: $V = 1 - b_I/b_U$.
 145 Asexual hosts are strictly clonal. The expected genotypic contribution by asexual hosts is given by

$$A'_{ij} = W_U A_{ij,U}(t) + W_I A_{ij,I}(t), \quad (2)$$

146 where $A_{ij,U}$ and $A_{ij,I}$ are the number of uninfected and infected asexual hosts in a population.
 147 A proportion ϵ_{mix} of a population mixes with other populations. The expected number of offspring in the
 148 next generation (accounting for contributions from all populations) is given by

$$\begin{aligned} \mathbb{E}(S_{ij}^k(t+1)) &= f_{\text{sex}} \left((1 - \epsilon_{\text{mix}}) (S_{ij}^k)' + \frac{\epsilon_{\text{mix}}}{n_{\text{pop}} - 1} \sum_{h \neq k} (S_{ij}^h)' \right), \\ \mathbb{E}(A_{ij}^k(t+1)) &= (1 - \epsilon_{\text{mix}}) (A_{ij}^k)' + \frac{\epsilon_{\text{mix}}}{n_{\text{pop}} - 1} \sum_{h \neq k} (A_{ij}^h)', \end{aligned} \quad (3)$$

149 where $f_{\text{sex}}(x)$ is the function that models sexual reproduction, including recombination probability r_{host} and
 150 outcrossing, and n_{pop} is the number of populations modeled. The total number of sexual and asexual hosts
 151 in the next generation is given by a Poisson random variables with means specified previously. We also
 152 allow for stochastic migration in order to avoid fixation:

$$\begin{aligned} S_{ij}^k(t+1) &\sim \text{Poisson}(\lambda = \mathbb{E}(S_{ij}^k(t+1))) + \text{Bernoulli}(p = p_{ij,\text{sex}}), \\ A_{ij}^k(t+1) &\sim \text{Poisson}(\lambda = \mathbb{E}(A_{ij}^k(t+1))) + \text{Bernoulli}(p = p_{ij,\text{asex}}), \end{aligned} \quad (4)$$

153 where $p_{ij,\text{sex}}$ and $p_{ij,\text{asex}}$ are the probabilities of a single sexual or asexual host with genotype ij entering a
 154 subpopulation.

155 Infection is modeled using the matching alleles model (Otto and Michalakis 1998). We assume that snails
 156 are equally susceptible to parasites that match either haplotype. However, parasites must match the host
 157 haplotype at both loci in order to successfully infect a host. The expected number of infected hosts that
 158 are infected with parasite with genotype i at generation t is given by:

$$I_i(t) = f_{\text{mutation}} \left(\sum_j \left(S_{ij,i,I}(t) + A_{ij,i,I}(t) \right) \right). \quad (5)$$

159 $S_{ij,i,I}(t)$ and $A_{ij,i,I}(t)$ represent the expected numbers of sexual and asexual hosts that have genotype ij
 160 and are infected with parasites that have genotype i . The function f_{mutation} models mutation at a single

locus with probability r_{parasite} (Ashby and King 2015). We further account for stochastic migration with probability $p_{i,\text{parasite}}$ to avoid fixation:

$$I'_i(t) \sim I_i(t) + \text{Bernoulli}(p = p_{i,\text{parasite}}). \quad (6)$$

The total expected number of infectious contacts made by infected hosts within a population is given by $\lambda_i^k = \beta^k I_i^k(t)$, where β^k is the transmission rate of each population. We model mixing between subpopulations by allowing infected hosts to make contact with susceptible hosts in other populations; we assume that a proportion ϵ_{mix} of the infectious contacts are equally distributed among other subpopulations. The total amount of infectious contact, coming from hosts that carry genotype i parasite, that is received by susceptible hosts in population k is given by

$$\lambda_{i,\text{total}}^k = (1 - \epsilon_{\text{mix}})\lambda_i^k + \frac{\epsilon_{\text{mix}}}{n_{\text{pop}} - 1} \sum_{l \neq k} \lambda_i^l \quad (7)$$

Then, the force of infection that a susceptible host with genotype ij experiences in generation $t + 1$ is given by

$$\text{FOI}_{ij}^k = \frac{\lambda_{i,\text{total}}^k + \lambda_{j,\text{total}}^k}{2N^k(t + 1)}, \quad (8)$$

where $N^k(t + 1) = \sum_{i,j} S_{ij}^k(t + 1) + A_{ij}^k(t + 1)$ is the total number of hosts in generation $t + 1$. The probability that a susceptible host with genotype ij in population k becomes infected in the next generation is given by

$$P_{ij}^k(t + 1) = 1 - \exp(-\text{FOI}_{ij}^k). \quad (9)$$

Finally, the number of infected hosts in the next generation is determined by a binomial process:

$$\begin{aligned} S_{ij,I}^k(t + 1) &\sim \text{Binom}(S_{ij}^k(t + 1), P_{ij}^k(t + 1)), \\ A_{ij,I}^k(t + 1) &\sim \text{Binom}(A_{ij}^k(t + 1), P_{ij}^k(t + 1)). \end{aligned} \quad (10)$$

The expected number of infected hosts that have genotype ij and are infected by parasites with genotype i in the next generation is proportional to the amount of infectious contact that was made in the current generation:

$$\begin{aligned} S_{ij,i,I}^k(t + 1) &= \frac{2^{\delta_{ij}} \lambda_{i,\text{total}}^k}{\lambda_{i,\text{total}}^k + \lambda_{j,\text{total}}^k} S_{ij,I}^k(t + 1) \\ A_{ij,i,I}^k(t + 1) &= \frac{2^{\delta_{ij}} \lambda_{i,\text{total}}^k}{\lambda_{i,\text{total}}^k + \lambda_{j,\text{total}}^k} A_{ij,I}^k(t + 1) \end{aligned} \quad (11)$$

where a δ_{ij} is a Kronecker delta (δ_{ij} equals 1 when $i = j$ and 0 otherwise).

2.3 Simulation design and parameterization

Many Red Queen models have focused on competition between a single asexual genotype and multiple sexual genotypes or have assumed equal genetic diversity between asexual and sexual hosts (see Ashby and King (2015) for a review of previous Red Queen models) but neither of these assumptions is realistic. In contrast, Ashby and King (2015) adopted a more realistic approach by incorporating stochastic external migration of an asexual genotype to a population and allowing for asexual genetic diversity to vary over time. Here, we combine these methods. We allow for stochastic external migration of asexual hosts with different genotypes into the system but fix the number of asexual genotypes (denoted by G_{asex}) that can be present in the system. Since sexual hosts have limited genetic diversity in our model (diploid hosts with two biallelic loci yields a total of 10 genotypes), allowing for unlimited migration of asexual hosts will cause the sexual population to be easily outcompeted by the asexual population. Limiting asexual genetic diversity allows us to account for the intrinsic difference in sexual and asexual diversity and make a compromise between simple and realistic models. At the beginning of each simulation, a pool of G_{asex} asexual genotypes are sampled at random from the entire genotypic space; these are the genotypes that are available for external immigration into subpopulations. Then, we use our model-based comparison with data to estimate G_{asex} to test whether the data are consistent with the dynamics of a model that allows for more than a single asexual clone.

To account for differing number of sexual and asexual genotypes, we let

$$p_{ij,\text{sex}} = 1 - (1 - p_{\text{host}})^{1/G_{\text{sex}}},$$

$$p_{ij,\text{asex}} = \begin{cases} 1 - (1 - p_{\text{host}})^{1/G_{\text{asex}}} & \text{if } ij \in \{\text{asexual genotypes}\} \\ 0 & \text{otherwise} \end{cases}, \quad (12)$$

where p_{host} is the probability that at least one sexual or asexual host enters the population in a generation.

We scale the probability of an infected host carrying parasite genotype i in a similar way for

interpretability:

$$p_{i,\text{parasite}} = 1 - (1 - p_{\text{infected}})^{1/G_{\text{parasite}}}, \quad (13)$$

where p_{infected} is the probability that at least one infected host enters the population in a generation. The number of parasite genotypes G_{parasite} is equal to 4 (because parasites are assumed to be haploids with two loci).

Each simulation consists of 40 subpopulations. Every subpopulation is initialized with 2000 sexual hosts, of which 80 are infected. They are assumed to be in Hardy-Weinberg equilibrium where the allele frequency in each locus is equal to 0.5. In order to account for variation in infection rates among subpopulations, the transmission rate, β^k , is randomly drawn for each subpopulation from a Gamma distribution with mean β_{mean} and coefficient of variation β_{CV} . The simulation runs for 500 generations without the introduction of asexuals. At generation 501, 10 asexual hosts of a single genotype are introduced to each population (the asexual genotype introduced can vary across subpopulations) and simulation runs for a further 600 generations while allowing for stochastic migration of asexuals.

2.4 Approximate Bayesian Computation

We use Approximate Bayesian Computation (ABC) to estimate parameters from data (Toni et al. 2009). ABC relies on comparing summary statistics of observed data with those of simulated data; it is particularly useful when the exact likelihood function is not available. Each summary statistic (also referred to as a *probe*) represents an aspect of a system Kendall et al. (1999); probe matching can be more powerful than classical likelihood-based approaches because it allows to evaluate model fits based on biologically meaningful aspects of a system (Wood 2010). We consider the mean proportion of infected and sexually reproducing snails in the system and variation in these proportions — measured by a coefficient of variation (CV) — across space (population) and time as our probes. As Dagan et al. (2013a) and McKone et al. (2016) only reported the proportion of males, the proportion of sexual hosts is assumed to be twice the proportion of males.

CV across space is calculated by first calculating mean proportions by averaging across time (generation)

223 for each site (subpopulation) and then taking the the CV of these mean proportions. CV across time
 224 (generation) is calculated by first averaging proportions across space (subpopulation) at each generation
 225 and then taking the CV. For purely spatial data (Dagan et al. (2013a) and McKone et al. (2016)), CV
 226 across space is calculated without averaging across time.

227 Because ABC is a Bayesian method, we must specify prior distributions for all parameters. We use weakly
 228 informative priors for all estimated parameters except c_b , a scale parameter for the cost of sex (see Table 1
 229 for prior distributions used and parameters assumed). The prior distribution for the scale parameter is
 230 chosen so that 95% prior quantile range of cost of sex ($2/c_b$) is approximately equal to the 95% confidence
 231 interval reported by Gibson et al. (2017). For brevity, all other parameters are assumed to be fixed. While
 232 it is a common practice to fix several parameters of a Red Queen model (e.g., Lively (2010b), Ashby and
 233 King (2015), Haafke et al. (2016), Ashby et al. (2019)), failing to explicitly account for uncertainty in all
 234 parameters of a model can lead to overly confident conclusions (Elderd et al. 2006).

235 We use the Population Monte Carlo approach (Turner and Van Zandt 2012), which allows for efficient
 236 sampling while ensuring that final result still converges to a correct (approximate) Bayesian posterior. The
 237 Population Monte Carlo approach begins with a basic ABC. For each random parameter sample drawn
 238 from the prior distribution, the model is simulated and a sample of a subpopulation is drawn from the
 239 simulated system, equal to the number of sites collected in a study. Summary statistics are calculated
 240 based on the last 100 generations out of 1100 generations; the current parameter set is accepted if the
 241 distance between simulated and observed data is less than a specified tolerance value. Distance is measured
 242 by the sum of absolute differences in summary statistics between simulated and observed data. This
 243 process is repeated until 100 parameter sets are accepted.

244 After the first run ($t = 1$), equal weights ($w_{i,1} = 1/100$) are assigned to each accepted parameter set $\theta_{i,1}$
 245 ($i = 1, 2, \dots, 100$). A weighted random sample (θ^*) is drawn from the accepted parameters of the previous
 246 run ($t - 1$) with weights $w_{i,t-1}$. For any run $t > 1$, a parameter sample ($\theta_{i,t}$) is proposed from a
 247 multivariate normal distribution with a mean θ^* and a variance covariance matrix that is equal to
 248 $\sigma_{t-1}^2 = 2\text{Var}(\theta_{1:N,t-1})$, where $\text{Var}(\theta_{1:N,t-1})$ is the weighted variance covariance matrix of the accepted

Notation	Description	Prior distribution/parameter values	Source
β_{mean}	Mean transmission rate	Gamma($k = 2, \theta = 10$)	Assumption
β_{CV}	CV transmission rate	Gamma($k = 2, \theta = 0.5$)	Assumption
V	Virulence	Beta($\alpha = 6, \beta = 2$)	Assumption
ϵ_{mix}	Mixing proportion	Beta($\alpha = 1, \beta = 9$)	Assumption
G_{asex}	Number of asexual genotypes	$1 + \text{BetaBinomial}(N = 9, p = 3/9, \theta = 5)$	Assumption
c_b	Cost of sex scale	LogNormal($\mu = -0.07, \sigma = 0.09$)	Gibson et al. (2017)
s	Proportion of male offspring produced	0.5	Assumption
b_U	Number of offspring produced by an uninfected host	20	Lively (2010b)
b_I	Number of offspring produced by an infected host	$(1 - V)b_U$	Lively (2010b)
a_U	Density dependent effect coefficient of uninfected hosts	0.001	Lively (2010b)
a_I	Density dependent effect coefficient of infected hosts	0.001	Lively (2010b)
r_{host}	Host recombination probability	0.2	Lively (2010b)
r_{parasite}	Parasite mutation probability	0.05	Assumption
p_{host}	Probability that at least one sexual or asexual host enters the population	0.1	Assumption
p_{infected}	Probability that at least one infected host enters the population	0.02	Assumption

Table 1: Parameter descriptions and values. Parameters with prior distributions are estimated via Approximate Bayesian Computation (ABC). Parameters k and θ in Gamma distribution represent shape and scale parameters where mean and squared CV are given by $k\theta$ and $1/k$, respectively. α and β in Beta distribution represent shape parameters where mean and squared CV are given by $\alpha/(\alpha + \beta)$ and $\beta/(\alpha^2 + \alpha\beta + \alpha)$. N , p and θ in Beta binomial distributions represent number of trials, probability of success, and overdispersion parameters (Morris et al. 1983). Parameters μ and σ in a log-normal distribution represent mean and standard deviation on a log scale. All other parameters are fixed throughout simulations.

parameters from the previous run and N is the total number of accepted parameters from the previous run.

The parameter G_{asex} is rounded to the nearest integer and the model is simulated. If a proposed parameter is accepted, the following weight is assigned:

$$w_{i,t} = \frac{\pi(\boldsymbol{\theta}_{i,t})}{\sum_{i=1}^{100} w_{j,t-1} q(\boldsymbol{\theta}_{j,t-1} | \boldsymbol{\theta}_{i,t}, \sigma_{t-1}^2)}$$

where $\pi(\cdot)$ is a prior density and $q(\cdot | \boldsymbol{\theta}_{i,t}, \sigma_{t-1}^2)$ is a multivariate normal density with mean $\boldsymbol{\theta}_{i,t}$ and variance covariance matrix σ_{t-1}^2 . For each run, 100 parameters are accepted and weights are normalized at the end to sum to 1.

For each observed data set, we perform 4 runs with decreasing tolerance. First three tolerance values are chosen in a decreasing sequence to reach the final step quicker. The tolerance value of the final run is chosen so that a parameter set will be accepted if its each simulated summary statistic deviates from the corresponding observed summary statistic by 0.1 units on average. For spatial data (Dagan et al. 2013a, McKone et al. 2016), four summary statistics are compared: the mean proportion of infected and sexually reproducing hosts and CV in these proportions across populations. Tolerance values of 1.6, 0.8, 0.6 and 0.4 are used for each run. For spatiotemporal data (Vergara et al. 2014b), we additionally compare the CV of proportions of infected and sexually reproducing hosts across generations. In this case, larger tolerance values (2.4, 1.2, 0.9 and 0.6) are used for each run to account for a higher number of summary statistics being compared.

All statistical results are weighted by parameter weights of the final run; confidence intervals are obtained by taking weighted quantiles. However, it is always not possible to obtain the exact 95% confidence interval when the smallest (or the largest) value has a weight greater than 0.025. In these cases, we take the lowest (or largest) possible quantile that is closest to the 2.5% (or the 97.5%) quantile. While this procedure results in a slightly narrower confidence interval, the rounding errors in quantiles are negligible. This rule leads to one instance where we use the 2.82% quantile instead of the 2.5% quantile as the lower end of a confidence interval.

2.5 Power analysis

Using estimated parameters for each data set, we calculate the power to detect a significant positive correlation between infection prevalence and frequency of sexual hosts. For each parameter sample from the final run of the ABC, 10 simulations are run. For each simulation, we take the last two generations — assuming that a year contains two snail generations (estimated generation time is 4-9 months (Neiman et al. 2005)) — from the simulation and choose n populations at random from 40 simulated populations. For each selected population, hosts are divided into four categories based on their infection status (infected/uninfected) and reproductive mode (asexual/sexual), and the mean proportion of hosts in each category is calculated by averaging over two generations. Independent multinomial samples of size m are drawn from each selected population based on the proportions in every four categories. Correlation between the proportion of infected hosts and the proportion of sexual hosts is tested using the Spearman's rank correlation at a 5% significance level.

While we do not explicitly try to match observed correlations, this procedure is essentially a retrospective power analysis. The main purpose of a power analysis is to design an experiment and determine an appropriate sample size (Cohen 1992). If instead, power is calculated based on an observed effect size after an experiment, the estimated power is likely to be correlated with the observed significance of the result; therefore, interpreting significance (or a lack of significance) of a statistical result based on a retrospective power analysis is inappropriate (Goodman and Berlin 1994, Senn 2002).

Here, we do not try to justify the significant correlation observed by (McKone et al. 2016) or the nonsignificant correlation observed by (Dagan et al. 2013a) based on this power analysis. Instead, we use a power analysis to better understand how a study design or the dynamics of the model might affect the correlation between the prevalence of infection and sexual reproduction while using biologically realistic parameters.

3 Results

First, we compare observed summary statistics with either fitted or predicted summary statistics (Fig. 1).

Fitted (filtered) summary statistics (Fig. 1) are values that have been accepted by the ABC procedure as being sufficiently close to observed summary statistics; these can be interpreted as model-based estimates of the summary statistics (and associated confidence intervals) of the study sites. *Predicted* (unfiltered) summary statistics, in contrast (Fig. 2), use parameters from the posterior distribution sampled by ABC, but generate new simulations of the dynamics and calculate summary statistics from randomly selected subpopulations. These values represent summary statistics sampled from the full distribution of coevolutionary dynamics that are consistent with the observed dynamics; they represent the expected range of dynamics from randomly sampled host-parasite metapopulations with a given distribution of parameters. Because they include an additional level of dynamical variability, the predicted summary statistics range much more widely than the fitted summary statistics.

Our simple meta-population Red Queen model can capture observed variation in infection prevalence and frequency of sexual hosts reasonably well; both temporal and spatial variation (measured by CV across mean proportions) are well-matched by the model. However, as model fitting is performed by minimizing the sum of absolute distance between observed and simulated summary statistics, our method does not guarantee that all summary statistics are equally well-fitted. The model tends to overestimate the mean proportion of infected hosts. The observed mean proportions of infected hosts are 0.175 (Dagan et al. 2013a), 0.051 (McKone et al. 2016), and 0.440 (Vergara et al. 2014b), whereas their ABC-based estimates are 0.240 (95% CI: 0.174 - 0.286), 0.313 (95% CI: 0.233 - 0.404), and 0.542 (95% CI: 0.360 - 0.730), respectively. The model underestimates the mean proportion of sexual hosts for Dagan et al. (2013a) and Vergara et al. (2014b). Observed mean proportions of sexual hosts are 0.045 and 0.704, respectively, whereas their ABC-based estimates are 0.026 (95% CI: 0.007 - 0.048) and 0.596 (95% CI: 0.441 - 0.679). To further diagnose the fit, we compare the predicted relationships between the mean proportion of infected hosts and the mean proportion of sexual hosts across subpopulations with the observed data (Fig. 2). Despite its accuracy in reproducing summary statistics reported by Dagan et al. (2013a), our




Figure 1: Summary statistics of the observed data vs. distribution of summary statistics of the simulated data from the posterior samples. Dotted horizontal lines represent observed summary statistics. Violin plots show the weighted distribution of fitted summary statistics (i.e., summary statistics that were accepted during ABC). Error bars show 95% weighted quantiles of predicted summary statistics. The weights correspond to the posterior distribution weights from ABC. For each posterior sample, 10 simulations are run and each simulated system is sampled at random 100 times so that each sample consists of the same number of populations as in the fitted data. All univariate summary statistics are matched reasonably well, except for the mean proportion of infected hosts in McKone *et al.* (2016) (indicated by an asterisk).

model poorly captures the relationship between the mean proportion of infected hosts and the mean
 proportion of sexual hosts (Fig. 2; Dagan et al. (2013a)). The model predicts sexual reproduction to be
 well maintained when infection prevalence is high ($> 40\%$) whereas the observed data (Dagan et al. 2013a)
 suggests that sexual reproduction can only be supported when infection prevalence is low ($< 20\%$).
 McKone et al. (2016) also found sexually reproducing snails in sites with low infection prevalence ($< 20\%$).
 On the other hand, Vergara et al.'s (2014b) data set suggests that sexually reproducing populations are
 likely to have a relatively high prevalence of infection (approximately $20\% - 80\%$). This observation is
 broadly consistent with our model prediction. Most of the data points fall within the range of model
 predictions (Fig. 2; Vergara et al. (2014b)). However, there is one site in which more than 90% of the snails
 were found to be sexual throughout the study period of 5 years (Vergara et al. 2014b); our model fails to
 maintain such high levels of sexual reproduction.

There is a high posterior density region in which the proportion of infected hosts remains almost constant
 (around 0.5) among subpopulations while the proportion of sexual hosts can range from 0 to 0.3 (visible in
 the fits to McKone et al. (2016)). As transmission rate (β) increases, selection for sexual hosts increases
 but the increasing number of resistant offspring prevents further infection from occurring and can decrease
 overall infection prevalence. This trend is consistent with previous results of Lively (2001) who noted that
 there is a region in which either sexual and asexual reproduction can be selected exclusively under the
 same infection prevalence.

The proportion of sexual hosts decreases when infection prevalence is extremely high (visible in the fits to
 Vergara et al. (2014b)). This pattern can be explained by the decrease in fitness of sexual hosts with the
 increase in infection prevalence, predicted by Ashby and King (2015). The same pattern can be seen in
 earlier work by Lively (2010b) although it was not discussed.

Fig. 3 presents parameter estimates. Keeping in mind that we do not obtain good fits to data from Dagan
 et al. (2013a) and McKone et al. (2016), we still find that high virulence and a low ratio of asexual to
 sexual genetic diversity are necessary to explain the observed dynamics. Moreover, we are able to capture
 observed differences in mean and variation in infection prevalence among studies in our estimates of




Figure 2: Predicted relationship between mean infection prevalence and mean proportion of sexual hosts in each population. For each posterior sample, 10 simulations are run. For each population within a simulation, mean infection prevalence and mean proportion of sexual hosts is calculated by averaging across last two generations. Each point, consisting of mean infection prevalence and meal proportion of sexual hosts, is assigned the weight of the parameter used to simulated the population. The density at each cell is calculated as the sum of weights of the points that lie within it. Cell densities are normalized by dividing by the maximum grid densities for each fit. Dashed contours are calculated at $\log(\text{density}) = -4$. Open triangles represent observed data; proportion of sexual hosts is computed by doubling proportion of male hosts.

transmission rate parameters (β_{mean} and β_{CV}).

Our fits to McKone et al. (2016) suggest that the scale parameter for the cost of sex, c_b , should be higher than our prior assumption based on Gibson et al. (2017) that estimated the cost of sex to be 2.14 (95% CI: 1.81 - 2.55). Ashby and King (2015) defined c_b as additional costs and benefits of sex, where $c_b = 1$ corresponds the two fold cost. Under their interpretation, our estimate of c_b corresponds to a slightly lower estimate of the cost of sex: 1.95 (95% CI: 1.68 - 2.4). (We propose an alternate interpretation to this parameter estimate below).

Finally, our power analyses suggest that there is high power to detect a positive correlation between infection prevalence and frequency of sexual hosts in the systems studied by Dagan et al. (2013a) and




Figure 3: Parameter estimates from Sequential Monte Carlo Approximate Bayesian

Computation. Violin plots represent weighted distribution of 100 posterior samples obtained from ABC. Violin plots for prior distribution is obtained by drawing 10000 random parameter samples from the prior distribution. G_{asex} is a discrete variable but is drawn on a continuous scale for convenience. See Table 1 for full parameter descriptions and their prior distributions.

McKone et al. (2016) (Fig. 4). Such high power predicted for Dagan et al. (2013a) is particularly surprising given that they were not able to observe the expected correlation. This discrepancy implies that the snail populations studied by Dagan et al. (2013a) show sufficient variation in infection prevalence in order for the correlation to be observed under pure Red queen selection, but other underlying factors that are neglected by our model may have caused the populations to deviate from their expected behaviors. On the other hand, our model predicts low power for detecting the positive correlation for the system studied by Vergara et al. (2014b) (Fig. 4).

Overall, our analysis suggests that increasing the number of study sites is a more effective way to increase power than increasing the number of samples per site (Fig. 4). The correlation between the proportion of infected and sexual hosts depends on the spatial distribution of parasite infection; increasing the number of

sites can better capture this distribution and increase the power to observe the correlation. On the other hand, collecting more samples from a site only gives us a more accurate estimate of the two proportions for that specific site; it does not tell us how these proportions are distributed across space. As a result, the power quickly saturates once we have a sufficient number of samples (> 100) to estimate the two proportions reliably at each site.

While we originally planned to perform power analysis using Spearman's rank correlation, we repeated the analysis using Pearson's correlation after applying arcsine square root transformation (Lively 1992) to see whether this procedure improves power. Using Pearson correlation gives slightly higher power to detect the positive correlation between frequency of sexual hosts and infection prevalence (see Appendix).




Figure 4: Power to detect a statistically significant positive correlation between infection prevalence and frequency of sexual hosts. Spearman's rank correlation was used to test for correlation between infection prevalence and frequency of sexual hosts in simulated data from the posterior distributions.

4 Discussion

A simple metapopulation model for host-parasite coevolution is able to match the observed prevalence of sexual reproduction and trematode infection in snail populations (Fig. 1). A direct comparison between a model and a data set allows us to infer biologically meaningful parameters of a model (Fig. 3) and estimate the power to observe a significant, positive correlation between the proportion of infected hosts and the proportion of sexual hosts (Fig. 4), as predicted by Lively (1992). However, discrepancies between the model predictions and observed data suggest that a simple host-parasite coevolution model may not be able to sufficiently explain the maintenance of sexual reproduction observed in snail populations (Fig. 2). A model that fits poorly can sometimes tell us more about a biological system than a model that fits well. Our model clearly failed to match the data presented by Dagan et al. (2013a) (Fig. 2). The snail populations studied by Dagan et al. (2013a) live in qualitatively different environments from the two other snail populations that we considered (Vergara et al. 2014b, McKone et al. 2016). For example, their habitats are subject to seasonal flash floods (Ben-Ami and Heller 2007), which can affect reproductive strategies of snails and interfere with the host-parasite coevolution (Lytle 2000, Ben-Ami and Heller 2007). As a result, a positive correlation between infection prevalence and frequency of sexual reproduction could not be detected from the system even though our model (which does not incorporate disturbance) predicts high power. These results suggest that a significant correlation (or the lack of it) between infection prevalence and the frequency of sexual reproduction may not provide sufficient evidence for (or against) the role of parasites in maintaining sexual reproduction in the host population. The model fits to Dagan et al. (2013a) and McKone et al. (2016) suggest that the cost of sex can be overcome and sexual reproduction can be maintained only if infection prevalence is much higher than the observed prevalence (Fig. 2). In other words, the benefit of producing offspring with novel genotype is relatively small when infection prevalence is low. In order to support sexual reproduction at lower infection prevalence, the benefit of sex must be greater or other mechanisms must compensate for the difference. As our model relies on a simple structure and strong parametric assumptions, an additional benefit of sex can only be provided by lowering the cost of sex (i.e., increasing the scale parameter, c_b).

The simple structure of the model and limited genetic diversity can explain the discrepancy between model prediction and the observed data by McKone et al. (2016). Here, we assumed that host resistance to infection is determined entirely by two biallelic loci, resulting in 10 possible genotypes. However, it is unlikely that such a simple model can capture the genetic interaction between hosts and parasites observed in nature. Although exact genetic architecture that determines trematode infection in snails (e.g., the number and inheritance of loci involved in parasite resistance) is not known, the documented genetic diversity of snails is far greater than what our model assumes (Fox et al. 1996, King et al. 2011, Dagan et al. 2013a). Increasing the maximum possible genetic diversity of the model would have allowed sexual hosts to escape infection more easily and maintained sexual reproduction at a lower prevalence of infection (Lively 2010a, King and Lively 2012, Ashby and King 2015).

While the positive correlation between frequency of sexual hosts and prevalence of infection can provide evidence for the effect of the parasite on the maintenance of sexual reproduction in the host population, it does not fully capture the coevolutionary process. In particular, the positive correlation represents a contrast between populations that undergo Red Queen dynamics and those that do not (rather than a continuous relationship between the frequency of sexual hosts and prevalence of infection): low-risk populations will be dominated by asexual individuals whereas high-risk populations will consist of both sexual and asexual individuals. Lively (2001) gave a similar explanation and predicted that large variation in the risk of infection is required to observe the predicted correlation.

The similarity between the correlations predicted by the model and the correlations observed in nature may provide us more confidence that the observed populations provide evidence for the RQH (Fig. 5). However, the strength of the correlation is not a direct measure of the strength of selection imposed on the host population. Moreover, when populations are actively coevolving, wide range of correlations can be detected (Fig. 5: Vergara et al. (2014b)). While simple statistical summaries, such as a correlation coefficient, are easily accessible, more sophisticated and mechanistic statistical models that yield biologically interpretable parameters may be better suited for understanding the effect of parasites on the maintenance of sexual reproduction in the host population.




Figure 5: Predicted vs. observed correlation between infection prevalence and frequency of sexual hosts. Violin plots show weighted distribution of predicted strength of correlation based on ABC fits. Predicted correlation was measured for each simulation from the posterior by taking into account the last two generations from each simulated population. Triangles and circles represent observed correlation sizes from previous studies. Dagan *et al.* (2013a) and Vergara *et al.* (2014b) do not report the size of the correlation; correlations are re-calculated from their data sets using both Pearson (after applying arcsine square root transformation) and Spearman correlations.

Mathematical models have used extensively to build theoretical foundations for the evolution of sex but only a few models have been confronted with data. The idea of the two-fold cost of sex was only recently tested directly by fitting a theoretical model to observed data (Gibson *et al.* 2017). Making statistical inference on the observed systems and testing theoretical models against data may provide deeper insight into underpinnings of maintenance of sexual reproduction.

References

- Agrawal, A. F. and C. M. Lively (2001). Parasites and the evolution of self-fertilization. *Evolution* 55(5), 869–879.
- Ashby, B. and S. Gupta (2014). Parasitic castration promotes coevolutionary cycling but also imposes a cost on sex. *Evolution* 68(8), 2234–2244.
- Ashby, B., R. Iritani, A. Best, A. White, and M. Boots (2019). Understanding the role of eco-evolutionary feedbacks in host-parasite coevolution. *J. Theor. Biol.* 464, 115–125.
- Ashby, B. and K. C. King (2015). Diversity and the maintenance of sex by parasites. *J. Evol. Biol.* 28(3), 511–520.
- Auld, S. K., S. K. Tinkler, and M. C. Tinsley (2016). Sex as a strategy against rapidly evolving parasites. *Proc. R. Soc. Lond., B, Biol. Sci.* 283(1845).
- Bell, G. (1982). *The Masterpiece of Nature: The Evolution and Genetics of Sexuality*. University of California Press.
- Ben-Ami, F. (2006). First report of the invasive freshwater snail *Tarebia granifera* (Lamarck, 1816) (Gastropoda: Thiariidae) from Israel. *Nautilus* 120(4), 156–161.
- Ben-Ami, F., D. Gold, and B. Fried (2005). Differential infectivity of *Transversotrema patialense* for naive fish. *J. Parasitol.* 91(4), 949–951.
- Ben-Ami, F. and J. Heller (2005). Spatial and temporal patterns of parthenogenesis and parasitism in the freshwater snail *Melanoides tuberculata*. *J. Evol. Biol.* 18(1), 138–146.
- Ben-Ami, F. and J. Heller (2007). Temporal patterns of geographic parthenogenesis in a freshwater snail. *Biol. J. Linn. Soc.* 91(4), 711–718.
- Ben-Ami, F. and J. Heller (2008). Sex versus parasitism versus density. *Biol. J. Linn. Soc.* 93(3), 537–544.

- Buckling, A. and P. B. Rainey (2002). Antagonistic coevolution between a bacterium and a bacteriophage. *Proc. R. Soc. Lond., B, Biol. Sci.* 269(1494), 931–936.
- Cohen, J. (1992). Statistical power analysis. *Curr. Dir. Psychol. Sci.* 1(3), 98–101.
- Dagan, Y., K. Liljeroos, J. Jokela, and F. Ben-Ami (2013a). Clonal diversity driven by parasitism in a freshwater snail. *J. Evol. Biol.* 26(11), 2509–2519.
- Dagan, Y., K. Liljeroos, J. Jokela, and F. Ben-Ami (2013b). Data from: Clonal diversity driven by parasitism in a freshwater snail. <https://doi.org/10.5061/dryad.f5t56>.
- Decaestecker, E., S. Gaba, J. A. Raeymaekers, R. Stoks, L. Van Kerckhoven, D. Ebert, and L. De Meester (2007). Host–parasite ‘Red Queen’ dynamics archived in pond sediment. *Nature* 450(7171), 870.
- Dybdahl, M. F. and C. M. Lively (1995a). Diverse, endemic and polyphyletic clones in mixed populations of a freshwater snail (*Potamopyrgus antipodarum*). *J. Evol. Biol.* 8(3), 385–398.
- Dybdahl, M. F. and C. M. Lively (1995b). Host-parasite interactions: infection of common clones in natural populations of a freshwater snail (*Potamopyrgus antipodarum*). *Proc. R. Soc. Lond., B, Biol. Sci.* 260(1357), 99–103.
- Dybdahl, M. F. and C. M. Lively (1998). Host-parasite coevolution: evidence for rare advantage and time-lagged selection in a natural population. *Evolution*, 1057–1066.
- Elder, B. D., V. M. Dukic, and G. Dwyer (2006). Uncertainty in predictions of disease spread and public health responses to bioterrorism and emerging diseases. *Proc. Natl. Acad. Sci. U.S.A.* 103(42), 15693–15697.
- Fox, J. A., M. F. Dybdahl, J. Jokela, and C. M. Lively (1996). Genetic structure of coexisting sexual and clonal subpopulations in a freshwater snail (*Potamopyrgus antipodarum*). *Evolution* 50(4), 1541–1548.
- Gibson, A. K., L. F. Delph, and C. M. Lively (2017). The two-fold cost of sex: Experimental evidence from a natural system. *Evol. Lett.* 1(1), 6–15.

475 Gibson, A. K., L. F. Delph, D. Vergara, and C. M. Lively (2018). Periodic, parasite-mediated selection for
476 and against sex. *Am. Nat.* 192(5), 537–551.

477 Gibson, A. K., J. Y. Xu, and C. M. Lively (2016). Within-population covariation between sexual
478 reproduction and susceptibility to local parasites. *Evolution* 70(9), 2049–2060.

479 Goodman, S. N. and J. A. Berlin (1994). The use of predicted confidence intervals when planning
480 experiments and the misuse of power when interpreting results. *Ann. Intern. Med.* 121(3), 200–206.

481 Haafke, J., M. Abou Chakra, and L. Becks (2016). Eco-evolutionary feedback promotes Red Queen
482 dynamics and selects for sex in predator populations. *Evolution* 70(3), 641–652.

483 Haldane, J. B. S. (1949). Disease and evolution. *La Ricerca Scientifica Supplement* 19, 68–76. Reproduced
484 in *Malaria: Genetic and Evolutionary Aspects* (2016), eds. Dronamraju, K. R., and P. Arese. Springer.

485 Hamilton, W. D. (1980). Sex versus non-sex versus parasite. *Oikos*, 282–290.

486 Hamilton, W. D., R. Axelrod, and R. Tanese (1990). Sexual reproduction as an adaptation to resist
487 parasites (a review). *Proc. Natl. Acad. Sci. U.S.A.* 87(9), 3566–3573.

488 Heller, J. and V. Farstey (1990). Sexual and parthenogenetic populations of the freshwater snail
489 *Melanoides tuberculata* in Israel. *Isr. J. Ecol. Evol.* 37(2), 75–87.

490 Howard, R. S., C. M. Lively, et al. (1994). Parasitism, mutation accumulation and the maintenance of sex.
491 *Nature* 367(6463), 554–557.

492 Jaenike, J. (1978). An hypothesis to account for the maintenance of sex within populations. *Evol.*
493 *Theory*. 3, 191–194.

494 Jokela, J., M. F. Dybdahl, and C. M. Lively (2009). The maintenance of sex, clonal dynamics, and
495 host-parasite coevolution in a mixed population of sexual and asexual snails. *Am. Nat.* 174(S1), S43–S53.

496 Kendall, B. E., C. J. Briggs, W. W. Murdoch, P. Turchin, S. P. Ellner, E. McCauley, R. M. Nisbet, and

- S. N. Wood (1999). Why do populations cycle? A synthesis of statistical and mechanistic modeling approaches. *Ecology* 80(6), 1789–1805.
- King, K. and C. M. Lively (2012). Does genetic diversity limit disease spread in natural host populations? *Heredity* 109(4), 199–203.
- King, K. C., L. F. Delph, J. Jokela, and C. M. Lively (2009). The geographic mosaic of sex and the red queen. *Curr. Biol.* 19(17), 1438–1441.
- King, K. C., L. F. Delph, J. Jokela, and C. M. Lively (2011). Coevolutionary hotspots and coldspots for host sex and parasite local adaptation in a snail-trematode interaction. *Oikos* 120(9), 1335–1340.
- King, K. C., J. Jokela, and C. M. Lively (2011). Parasites, sex, and clonal diversity in natural snail populations. *Evolution* 65(5), 1474–1481.
- Koskella, B. (2013). Phage-mediated selection on microbiota of a long-lived host. *Curr. Biol.* 23(13), 1256–1260.
- Koskella, B. (2014). Bacteria-phage interactions across time and space: merging local adaptation and time-shift experiments to understand phage evolution. *Am. Nat* 184(S1), S9–S21.
- Koskella, B. and C. M. Lively (2009). Evidence for negative frequency-dependent selection during experimental coevolution of a freshwater snail and a sterilizing trematode. *Evolution* 63(9), 2213–2221.
- Kumpulainen, T., A. Grapputo, J. Mappes, and M. Björklund (2004). Parasites and sexual reproduction in psychid moths. *Evolution* 58(7), 1511–1520.
- Lehtonen, J., M. D. Jennions, and H. Kokko (2012). The many costs of sex. *Trends Ecol. Evol.* 27(3), 172–178.
- Lively, C. (2009). The maintenance of sex: host–parasite coevolution with density-dependent virulence. *J. Evol. Biol.* 22(10), 2086–2093.

519 Lively, C. M. (1987). Evidence from a New Zealand snail for the maintenance of sex by parasitism.
520 *Nature* 328(6130), 519–521.

521 Lively, C. M. (1989). Adaptation by a parasitic trematode to local populations of its snail host.
522 *Evolution* 43(8), 1663–1671.

523 Lively, C. M. (1992). Parthenogenesis in a freshwater snail: reproductive assurance versus parasitic release.
524 *Evolution* 46(4), 907–913.

525 Lively, C. M. (2001). Trematode infection and the distribution and dynamics of parthenogenetic snail
526 populations. *Parasitology* 123(07), 19–26.

527 Lively, C. M. (2010a). The effect of host genetic diversity on disease spread. *Am. Nat* 175(6), E149–E152.

528 Lively, C. M. (2010b). An epidemiological model of host–parasite coevolution and sex. *J. Evol. Biol.* 23(7),
529 1490–1497.

530 Lively, C. M. (2010c). A review of Red Queen models for the persistence of obligate sexual reproduction.
531 *J. Hered.* 101(suppl.1), S13–S20.

532 Lively, C. M. (2018). Habitat heterogeneity, host population structure, and parasite local adaptation. *J.*
533 *Hered.* 109(1), 29–37.

534 Lively, C. M., M. F. Dybdahl, J. Jokela, E. E. Osnas, and L. F. Delph (2004). Host sex and local
535 adaptation by parasites in a snail-trematode interaction. *Am. Nat* 164(S5), S6–S18.

536 Lively, C. M. and R. S. Howard (1994). Selection by parasites for clonal diversity and mixed mating. In
537 *Infection, Polymorphism and Evolution*, pp. 1–11. Springer.

538 Lively, C. M. and J. Jokela (2002). Temporal and spatial distributions of parasites and sex in a freshwater
539 snail. *Evol. Ecol. Res.* 4(2), 219–226.

540 Lynch, Z. R., M. J. Penley, and L. T. Morran (2018). Turnover in local parasite populations temporarily
541 favors host outcrossing over self-fertilization during experimental evolution. *Ecol. Evol.* 8(13), 6652–6662.

542 Lytle, D. A. (2000). Biotic and abiotic effects of flash flooding in a montane desert stream. *Arch.*
 543 *Hydrobiol.*, 85–100.

544 MacPherson, A. and S. P. Otto (2018). Joint coevolutionary–epidemiological models dampen Red Queen
 545 cycles and alter conditions for epidemics. *Theor. Popul. Biol.* 122, 137–148.

546 May, R. M. and R. M. Anderson (1983). Epidemiology and genetics in the coevolution of parasites and
 547 hosts. *Proc. R. Soc. Lond., B, Biol. Sci.* 219(1216), 281–313.

548 McArthur, C. P. and D. Featherston (1976). Suppression of egg production in *Potamopyrgus antipodarum*
 549 (Gastropoda: Hydrobiidae) by larval trematodes. *N. Z. J. Zool.* 3(1), 35–38.

550 McKone, M. J., A. K. Gibson, D. Cook, L. A. Freymiller, D. Mishkind, A. Quinlan, J. M. York, C. M.
 551 Lively, and M. Neiman (2016). Fine-scale association between parasites and sex in *Potamopyrgus*
 552 *antipodarum* within a New Zealand lake. *N. Z. J. Ecol.* 40(3), 1.

553 Morran, L. T., R. C. Parrish, I. A. Gelarden, M. B. Allen, and C. M. Lively (2014). Experimental
 554 coevolution: rapid local adaptation by parasites depends on host mating system. *Am. Nat* 184(S1),
 555 S91–S100.

556 Morris, C. N. et al. (1983). Natural exponential families with quadratic variance functions: statistical
 557 theory. *Ann. Stat.* 11(2), 515–529.

558 Neiman, M., J. Jokela, and C. Lively (2005). Variation in asexual lineage age in *Potamopyrgus*
 559 *antipodarum*, a New Zealand snail. *Evolution* 59(9), 1945–1952.

560 Otto, S. P. (2009). The evolutionary enigma of sex. *Am. Nat* 174(S1), S1–S14.

561 Otto, S. P. and Y. Michalakis (1998). The evolution of recombination in changing environments. *Trends*
 562 *Ecol. Evol.* 13(4), 145–151.

563 Phillips, N. R. and D. M. Lambert (1989). Genetics of *Potamopyrgus antipodarum* (Gastropoda:
 564 Prosobranchia): evidence for reproductive modes. *N. Z. J. Zool.* 16(3), 435–445.

565 Rauch, G., M. Kalbe, and T. B. Reusch (2006). One day is enough: rapid and specific host–parasite
 566 interactions in a stickleback-trematode system. *Biol. Lett.* 2(3), 382–384.

567 Samadi, S., J. Mavárez, J.-P. Pointier, B. Delay, and P. Jarne (1999). Microsatellite and morphological
 568 analysis of population structure in the parthenogenetic freshwater snail *Melanooides tuberculata*: insights
 569 into the creation of clonal variability. *Mol. Ecol.* 8(7), 1141–1153.

570 Senn, S. J. (2002). Power is indeed irrelevant in interpreting completed studies. *BMJ* 325(7375), 1304.

571 Slowinski, S. P., L. T. Morran, R. C. Parrish, E. R. Cui, A. Bhattacharya, C. M. Lively, and P. C. Phillips
 572 (2016). Coevolutionary interactions with parasites constrain the spread of self-fertilization into
 573 outcrossing host populations. *Evolution* 70(11), 2632–2639.

574 Smith, J. M. (1978). *The Evolution of Sex*, Volume 54. Cambridge Univ. Press.

575 Smith, J. M. and M. Slatkin (1973). The stability of predator-prey systems. *Ecology* 54(2), 384–391.

576 Song, Y., C. S. Gokhale, A. Papkou, H. Schulenburg, and A. Traulsen (2015). Host-parasite coevolution in
 577 populations of constant and variable size. *BMC Evol. Biol.* 15(1), 212.

578 Thrall, P. H., A.-L. Laine, M. Ravensdale, A. Nemri, P. N. Dodds, L. G. Barrett, and J. J. Burdon (2012).
 579 Rapid genetic change underpins antagonistic coevolution in a natural host-pathogen metapopulation.
 580 *Ecol. Lett.* 15(5), 425–435.

581 Toni, T., D. Welch, N. Strelkowa, A. Ipsen, and M. P. Stumpf (2009). Approximate Bayesian computation
 582 scheme for parameter inference and model selection in dynamical systems. *J. R. Soc. Interface.* 6(31),
 583 187–202.

584 Turner, B. M. and T. Van Zandt (2012). A tutorial on approximate Bayesian computation. *J. Math.*
 585 *Psychol.* 56(2), 69–85.

586 Vergara, D., J. Jokela, and C. Lively (2014a). Data from: Infection dynamics in coexisting sexual and
 587 asexual host populations: support for the Red Queen hypothesis.
 588 <https://doi.org/10.5061/dryad.29nk3.2>.

589 Vergara, D., J. Jokela, and C. M. Lively (2014b). Infection dynamics in coexisting sexual and asexual host
590 populations: support for the Red Queen hypothesis. *Am. Nat* 184(S1), S22–S30.

591 Vergara, D., C. M. Lively, K. C. King, and J. Jokela (2013). The geographic mosaic of sex and infection in
592 lake populations of a New Zealand snail at multiple spatial scales. *Am. Nat* 182(4), 484–493.

593 Verhoeven, K. J. and A. Biere (2013). Geographic parthenogenesis and plant-enemy interactions in the
594 common dandelion. *BMC Evol. Biol.* 13(1), 23.

595 Vrijenhoek, R. C. (1998). Animal clones and diversity. *Bioscience* 48(8), 617–628.

596 Wallace, C. (1992). Parthenogenesis, sex and chromosomes in *Potamopyrgus*. *J. Molluscan Stud.* 58(2),
597 93–107.

598 Whitton, J., C. J. Sears, E. J. Baack, and S. P. Otto (2008). The dynamic nature of apomixis in the
599 angiosperms. *Int. J. Plant Sci.* 169(1), 169–182.

600 Winterbourn, M. (1974). Larval Trematoda parasitizing the New Zealand species of *Potamopyrgus*
601 (Gastropoda: Hydrobiidae). *Mauri Ora* 2, 17–30.

602 Wood, S. N. (2010). Statistical inference for noisy nonlinear ecological dynamic systems.
603 *Nature* 466(7310), 1102.

604 Zilio, G., L. Moesch, N. Bovet, A. Sarr, and J. C. Koella (2018). The effect of parasite infection on the
605 recombination rate of the mosquito *Aedes aegypti*. *PloS ONE* 13(10), e0203481.

Appendix




Figure A1: Power to detect a statistically significant positive correlation between infection prevalence and frequency of sexual hosts. Pearson correlation was used to test for correlation between square root arcsine transformed infection prevalence and frequency of sexual hosts in simulated data from the posterior distributions.