

Splines and basis expansion (week 3?)

6 Feb 2023

Table of contents

linear basis expansion	1
polynomial basis	2
piecewise polynomial bases	2
splines	3
spline terminology	3
truncated power basis	3
B-spline basis	3
natural cubic splines	3
smoothing splines	5
degrees of freedom and smoother matrix	5
multidimensional splines	6
tensor produce	6
thin-plate splines	6
null space	6
reduced-rank	6

linear basis expansion


- transformations of various kinds
- quadratic expansion
- nonlinear transformations
- indicator variables

Select or regularize from the expanded set.

polynomial basis

- polynomial basis: $y_i = \sum_{j=0}^n \beta_j x_i^j$

```
## replicate figure 3.2 from Wood
library(ggplot2)
x <- seq(0, 1, length = 101)
n <- 4
y <- sapply(0:n, \(j) x^j)
beta <- c(4.31, -10.72, 16.8, 2.22, -10.88)
y <- cbind(y, fx = y %*% beta)
dimnames(y) <- list(x = x, j = c(0:n, "f(x)"))
yy <- as.data.frame(as.table(y))
yy$x <- as.numeric(as.character(yy$x))
ggplot(yy, aes(x, Freq)) + geom_line() + facet_wrap(~j, scale = "free")
```



piecewise polynomial bases

- constant, linear, continuous
- basis functions
- translate from x_i to columns of \mathbf{X}

splines

- **piecewise** polynomials with continuity/smoothness constraints
- very useful for function approximation
- convert a single numeric predictor into a flexible basis
- efficient
- with multiple predictors, consider **additive models**
- handle interactions (multidim smooth surfaces) *if reasonably low-dimensional*: tensor products etc.

spline terminology

- **knots**: breakpoints (boundary, interior)
- order- M (ESL): continuous derivatives up to order $M - 2$ (cubic, $M = 4$)
- typically $M = 1, 2, 4$
- number of knots = df (degrees of freedom) -1 -intercept

truncated power basis

- $X^0 \dots X^n$
- remaining columns are $(x - \xi_\ell)^{M-1}$ where ℓ are the *interior knots*

B-spline basis

- splines of a given order with *minimal support* (i.e., local)
- basis functions defined by recursion (not pretty)
- convenient for regression splines (see below)

natural cubic splines

- linear constraints beyond boundary knots (so 2d and 3d derivatives are 0 at the boundaries)

```
library(splines)
bb <- bs(1:20, df = 5)
attributes(bb)[c("degree", "knots", "Boundary.knots")]
```

```
$degree
[1] 3
```

```
$knots
33.33333% 66.66667%
 7.333333 13.666667
```

```
$Boundary.knots
[1] 1 20
```


```
nn <- ns(1:20, df = 7)
attributes(nn)[c("degree", "knots", "Boundary.knots")]
```

```
$degree
[1] 3
```

```
$knots
14.28571% 28.57143% 42.85714% 57.14286% 71.42857% 85.71429%
 3.714286  6.428571  9.142857 11.857143 14.571429 17.285714
```

```
$Boundary.knots
[1] 1 20
```

```
par(mfrow = c(1,2), las = 1, bty = "l")
matplot(ns(1:20, df = 5), type = "l", main = "natural spline")
matplot(bs(1:20, df = 5), type = "l", main = "B-spline")
```



smoothing splines

- as many knots as data points
- plus squared-second-derivative (“wiggleness”) penalty

$$\text{RSS} + \lambda \int (f''(t))^2 dt$$

* defined on an infinite-dimensional space * minimizer is a natural cubic spline with knots at x_i

$$(\mathbf{y} - \mathbf{N}\theta)^\top (\mathbf{y} - \mathbf{N}\theta) + \lambda \theta^\top \Omega_N \theta$$

with $\{\Omega_N\}_{jk} = \int N_j''(t) N_k''(t) dt$ **generalized** ridge regression: penalize by $\lambda \Omega_N$ rather than λI * same data augmentation methods as before except that now we use $\sqrt{\lambda} C$ where C is a matrix square root of Ω_N

See Wood (2017), Perperoglou et al. (2019)

degrees of freedom and smoother matrix

- The equivalent of the hat matrix is

Wood, Simon N. 2017. *Generalized Additive Models: An Introduction with R*. CRC Texts in Statistical Science. Chapman & Hall. https://www.amazon.com/Generalized-Additive-Models-Introduction-Statistical-ebook/dp/B071Z9L5D5/ref=sr_1_1?ie=UTF8&qid=1511887995&sr=8-1&keywords=wood+additive+models.

Perperoglou, Aris, Willi Sauerbrei, Michal Abrahamowicz, and Matthias Schmid. 2019. “A Review of Spline Function Procedures in R.” *BMC Medical Research Methodology* 19 (1): 46. <https://doi.org/10.1186/s12874-019-0666-3>.

multidimensional splines


tensor produce

thin-plate splines


null space

reduced-rank

```
library(Matrix)
dd <- data.frame(x=1:200)
Xb <- model.matrix(~splines::bs(x, df = 10), data = dd)
Xn <- model.matrix(~splines::ns(x, df = 10), data = dd)
gridExtra::grid.arrange(
  ncol = 2,
  image(Matrix(Xb), aspect = "fill"),
  image(Matrix(Xn), aspect = "fill")
)
```



Dimensions: 200 x 11



Dimensions: 200 x 11