Eco-evolutionary dynamics of pathogen virulence

Ben Bolker, McMaster University

Departments of Mathematics & Statistics and Biology

University of Puerto Rico - Rio Piedras

26 April 2016

Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- Transient virulence of HIV
 - abc
- 4 Conclusions

Acknowledgements

People Arjun Nanda and Dharmini Shah; Christophe Fraser;
Marm Kilpatrick; Anson Wong

Support NSF IRCEB grant 9977063; QSE³ IGERT; NSERC Discovery grant

Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging disease
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions

Why is it interesting?

- Intellectual merit
 - Coevolutionary loops
 - Cryptic effects
 - Eco-evolutionary dynamics (Luo and Koelle, 2013)
 - Cool stories
 - Lots of data (sometimes)
- Broader applications
 - Medical
 - Conservation and management
 - Outreach

Why is it interesting?

- Intellectual merit
 - Coevolutionary loops
 - Cryptic effects
 - Eco-evolutionary dynamics (Luo and Koelle, 2013)
 - Cool stories
 - Lots of data (sometimes)
- Broader applications
 - Medical
 - Conservation and management
 - Outreach

- General public: badness
- Plant biologists: infectivity
- Evolutionists: loss of host fitness
- Theoreticians: rate of host mortality (mortality rate vs. case mortality vs. clearance)

Classical dogma monotonic trend toward avirulence

Ewald era virulence as an evolved (adaptive) trait. Tradeoff theory, modes of transmission.

post-Ewald more formal tradeoff models, mostly based on \mathcal{R}_0 optimization. Adaptive dynamics

Nov

- tradeoff backlash
- within-host dynamics/multi-level models
- eco-evolutionary dynamics
- host effects: resistance vs tolerance vs virulence

Classical dogma monotonic trend toward avirulence

Ewald era virulence as an evolved (adaptive) trait. Tradeoff theory, modes of transmission.

post-Ewald more formal tradeoff models, mostly based on \mathcal{R}_0 optimization. Adaptive dynamics

Nov

- tradeoff backlash
- within-host dynamics/multi-level models
- eco-evolutionary dynamics
- host effects: resistance vs tolerance vs virulence

Classical dogma monotonic trend toward avirulence

Ewald era virulence as an evolved (adaptive) trait. Tradeoff theory, modes of transmission.

post-Ewald more formal tradeoff models, mostly based on \mathcal{R}_0 optimization. Adaptive dynamics

Nov

- tradeoff backlash
- within-host dynamics/multi-level models
- eco-evolutionary dynamics
- host effects: resistance vs tolerance vs virulence

Classical dogma monotonic trend toward avirulence

Ewald era virulence as an evolved (adaptive) trait. Tradeoff theory, modes of transmission.

post-Ewald more formal tradeoff models, mostly based on \mathcal{R}_0 optimization. Adaptive dynamics

Now

- tradeoff backlash
- within-host dynamics/multi-level models
- eco-evolutionary dynamics
- host effects: resistance vs tolerance vs virulence


Overview


Basic tradeoff theory: assumptions

- Homogeneous, non-evolving hosts
- No superinfection/coinfection
- Horizontal, direct transmission
- Tradeoff between rate of transmission and length of infectious period


4D > 4A > 4E > 4E > 900

Tradeoffs, \mathcal{R}_0 , and r





Tradeoffs, \mathcal{R}_0 , and r




Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions

Conclusions

Epidemiological model


- SIR model



Overview

000000

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, N = 1(time units of host lifespan




Overview

000000

SIR model


- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, N = 1 (time units of host lifespan



Conclusions

Epidemiological model

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, N = 1 (time units of host lifespan)



Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value $(\bar{\alpha})$ and ecological context (proportion susceptible)
- ullet Constant additive genetic variance V_g
- Trait evolves toward increased fitness: rate proportional to Δfitness/Δtrait

Alternatives

Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value $(\bar{\alpha})$ and ecological context (proportion susceptible)
- ullet Constant additive genetic variance V_g
- Trait evolves toward increased fitness: rate proportional to Δfitness/Δtrait

Alternatives

Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value $(\bar{\alpha})$ and ecological context (proportion susceptible)
- Constant additive genetic variance V_g
- Trait evolves toward increased fitness: rate proportional to Δfitness/Δtrait

Alternatives

The model (2): evolutionary dynamics

Incorporate trait dynamics

Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value $(\bar{\alpha})$ and ecological context (proportion susceptible)
- ullet Constant additive genetic variance V_g
- Trait evolves toward increased fitness: rate proportional to Δfitness/Δtrait


Alternatives

Standard quantitative genetics model (Abrams, 2001):


- Fitness depends on mean trait value $(\bar{\alpha})$ and ecological context (proportion susceptible)
- ullet Constant additive genetic variance V_g
- Trait evolves toward increased fitness: rate proportional to Δfitness/Δtrait

Alternatives:


Evolutionary dynamics, cont.



Evolutionary dynamics, cont.



Power-law tradeoff curves



Outline

Overview

- 1 Overview
 - The evolution of host-pathogen theor
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

Sampling bias

Overview

- Lower host resistance/tolerance
- High transmission \rightarrow frequent coinfection \rightarrow selection for virulence
- ullet Disease-induced decrease in host density o selection for lowered virulence (Lenski and May, 1994)

Conclusions

(Why) are emerging pathogens more virulent?

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

- Sampling bias
- Lower host resistance/tolerance
- High transmission \rightarrow frequent coinfection \rightarrow selection for virulence
- ullet Disease-induced decrease in host density o selection for lowered virulence (Lenski and May, 1994)

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

Sampling bias

Overview

- Lower host resistance/tolerance
- High transmission \rightarrow frequent coinfection \rightarrow selection for virulence

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

Sampling bias

Overview

- Lower host resistance/tolerance
- ullet High transmission o frequent coinfection o selection for virulence
- ullet Disease-induced decrease in host density o selection for lowered virulence (Lenski and May, 1994)

Overview

Selection differs between the **epidemic** and **endemic** phases of an outbreak (Frank, 1996; Day and Proulx, 2004)

endemic phase selection for per-generation offspring production: maximize \mathcal{R}_0 , $\beta N/(\alpha + \mu)$

epidemic phase selection for per-unit-time offspring production: maximize r, $\beta N - (\alpha + \mu)$

Transient virulence

Selection differs between the **epidemic** and **endemic** phases of an outbreak (Frank, 1996; Day and Proulx, 2004)

endemic phase selection for per-generation offspring production: maximize \mathcal{R}_0 , $\beta N/(\alpha + \mu)$

epidemic phase selection for per-unit-time offspring production: maximize r, $\beta N - (\alpha + \mu)$

Transient virulence

Selection differs between the **epidemic** and **endemic** phases of an outbreak (Frank, 1996; Day and Proulx, 2004)

endemic phase selection for per-generation offspring production: maximize \mathcal{R}_0 , $\beta N/(\alpha + \mu)$

epidemic phase selection for per-unit-time offspring production: maximize r, $\beta N - (\alpha + \mu)$

Transient emerging virulence

When a parasite previously in eco-evolutionary equilibrium emerges in a new host population (at low density) it will show a transient peak in virulence as it spreads

How big is the peak? Does it matter?

Outline

- 1 Overview
 - The evolution of host-pathogen theor
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions


Model parameters

Overview


00000000

Parameter		Alternative	
С	Transmission scale	\mathcal{R}_0^*	Equilibrium \mathcal{R}_0
γ	Transmission shape	α^*	Equilibrium virulence
<i>I</i> (0)	Initial epidemic size	$1/N_0$	Inverse population size
Va	Genetic variance		

Example




Response variables



Time

Peak height



Estimates for emerging pathogens


Overview

Order of magnitude estimates for some emerging high-virulence pathogens:

Pathogen	\mathcal{R}_0^*	α^*	Reference
SARS	3	640	Anderson et al. (2004)
West Nile	1.61 - 3.24	639	Wonham et al. (2004)
HIV	1.43	6.36	Velasco-Hernandez et al. (2002)
myxomatosis	3	5	Dwyer et al. (1990)

Emerging pathogens: where are we?


$$CV_g = 0.5$$
, $I(0) = 10^{-3}$ (middle panel):



Outline

- 1 Overview
 - The evolution of host-pathogen theor
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions

- Mosquito-borne viral disease of rabbits
- Benign in South American rabbits, quickly fatal in European rabbits
- Well characterized (Fenner et al., 1956; Dwyer et al., 1990)



- Key parameter: genetic variance in virulence (evolvability)

- Key parameter: genetic variance in virulence (evolvability)
- Despite case studies of rapid pathogen evolution:
 - myxomatosis (Dwyer et al., 1990)
 - syphilis (Knell, 2004)
 - serial passage experiments (Ebert, 1998)
 - Plasmodium chabaudi (Mackinnon and Read, 1999)

we rarely have enough information to estimate V_{φ}


Estimating evolvability (V_g)

- Key parameter: genetic variance in virulence (evolvability)
- Despite case studies of rapid pathogen evolution:
 - myxomatosis (Dwyer et al., 1990)
 - syphilis (Knell, 2004)
 - serial passage experiments (Ebert, 1998)
 - Plasmodium chabaudi (Mackinnon and Read, 1999)


we rarely have enough information to estimate V_g

 Only (?) for myxomatosis do we know the variation in virulence among circulating strains


Myxomatosis grades vs. time




Myxomatosis variance vs. time




Myxomatosis virulence dynamics: power-law tradeoff




Myxomatosis virulence dynamics: realistic tradeoff



Myxo virulence: equilibrium start, power-law tradeoff




Myxo virulence: equilibrium start, realistic tradeoff



- eco-evo virulence dynamics are at least in the ballpark of what we would expect
- detailed shape of virulence curve very important ...
- ... as is genetic variance
- more realistic models (esp. of variance) needed?


Phage dynamics (Berngruber et al., 2013)



Outline


- 1 Overview
 - The evolution of host-pathogen theor
 - Toy models
- 2 Transient virulence and emerging disease
 - Overview
 - Toy model
 - Myxomatosis data
- Transient virulence of HIVabc
- 4 Conclusions

HIV tradeoff curve (Fraser et al., 2007)



- virulence/transmission from set-point viral load (SPVL)
- Fraser et al. (2007): plausible tradeoff observed from Rakai data
 - SPVL heritable
 - correlated with rate of progression to AIDS
 - correlated with probability (rate) of within-couple transmission

Model results



References

Shirreff et al. (2011): within-host assumptions

- generally based on Rakai estimates
- Normally distributed mutation of SPVL between hosts
- Normally distributed phenotypic variation
- different infectivity for early, mid, late stages of HIV
- tradeoff applies in mid stage: Hill functions $(x^d/(a+bx^d))$
- Weibull-distributed time in mid stage

Shirreff et al. (2011) epidemiological assumptions

implicit partnership-formation model


$$foi = \frac{\beta c}{\beta + c + 1/d}$$

• derived from \mathcal{R}_0 for an instantaneous-partnership-formation model, e.g. Hollingsworth et al. (2008); Diekmann and Heesterbeek (2000)

Epidemiological models for HIV Champredon et al. (2013)

Susceptible

Infectious



- need to track $\{S, SS, I(\alpha), SI(\alpha), II(\alpha, \alpha')\}$
- simplifications:
 - single-stage, exponential infectious period $(\beta = \text{time-weighted average of } \beta_1, \beta_2(\alpha), \beta_3)$
 - no phenotypic variation

Model variations

- 6 models:
 - extra-pair contact (yes/no) ×
 - instantaneous partnership formation (yes/no)
- implicit (Shirreff/Hollingsworth) model
- random-mixing model

Latin hypercube sampling


 sample evenly, randomly across (potentially many) uncertain parameters


 following Champredon et al. (2013), earlier Blower et al. (1991) x X X X X X X

Details

- r varies widely across parameters/model sets
- ullet scale parameters to a common r
- scale both uncoupled and extra-coupled rates relative to within-coupled rate
- response variables:
 - equilibrium virulence
 - peak virulence time (years)
 - peak virulence
 - relative peak virulence (peak/eq)

Virulence trajectories





- Eco-evolutionary dynamics is still a reasonable framework (Alizon et al., 2009; Luo and Koelle, 2013)
- need to know: genetic variance, shapes of tradeoffs
- theory meets molecular biology: mutations of large effect vs. quantitative variability
- next steps in connecting with data? general theory

- Eco-evolutionary dynamics is still a reasonable framework (Alizon et al., 2009; Luo and Koelle, 2013)
- need to know: genetic variance, shapes of tradeoffs
- theory meets molecular biology: mutations of large effect vs. quantitative variability
- next steps in connecting with data? general theory?

Crome (1997) on theory

When we regard theories as tight, real entities and devote ourselves to their analysis, we can limit our horizons and, worse, attempt to make the world fit them. A lot of ecological discussion is not about nature, but about theories, generalizations, or models supposed to represent nature . . .

References

- Abrams, P.A., 2001. Ecol Lett, 4:166-175.
- Alizon, S., Hurford, A., et al., 2009. J. Evol. Biol., 22:245–259. doi:10.1111/j.1420-9101.2008.01658.x.
- Anderson, R.M., Fraser, C., et al., 2004. Phil Trans R Soc London B, 359(1447):1091-1105.
- Berngruber, T.W., Froissart, R., et al., 2013. *PLoS Pathog*, 9(3):e1003209. doi:10.1371/journal.ppat.1003209.
- Blower, S.M., Hartel, D., et al., 1991. Philosophical Transactions of the Royal Society of London B: Biological Sciences, 331(1260):171-187. ISSN 0962-8436, 1471-2970. doi:10.1098/rstb.1991.0006.
- Champredon, D., Bellan, S., and Dushoff, J., 2013. PLoS ONE, 8(12):e82906. doi:10.1371%2Fjournal.pone.0082906.
- Crome, F.H.J., 1997. In W.F. Laurance and J. Richard O. Bierregard, editors, Tropical Forest Remnants: Ecology, Management and Conservation of Fragmented Communities, chapter 31, pages 485-501. University of Chicago Press, Chicago.
- Day, T. and Proulx, S.R., 2004. Amer Nat, 163(4):E40-E63.
- Diekmann, O. and Heesterbeek, J.A.P., 2000. Mathematical Epidemiology of Infectious Diseases: Model Building, Analysis and Interpretation. Wiley. ISBN 978-0-471-49241-2.
- Dwyer, G., Levin, S., and Buttel, L., 1990. *Ecol Monog.* 60:423–447.

- Ebert, D., 1998. Science, 282(5393):1432-1435.
- Fenner, F., Day, M.F., and Woodroofe, G.M., 1956. J Hyg (London), 54(2):284-302.
- Frank, S.A., 1996. Q Rev Biol, 71(1):37-78.
- Fraser, C., Hollingsworth, T.D., et al., 2007. PNAS, 104:17441–17446.
- Herbeck, J.T., Mittler, J.E., et al., 2014. *PLoS Computational Biology*, 10(6):e1003673. doi:10.1371/journal.pcbi.1003673.
- Hollingsworth, T., Anderson, R., and Fraser, C., 2008.
 Journal of Infectious Diseases, 198(5):687–693.
 ISSN 0022-1899, 1537-6613. doi:10.1086/590501.
- Knell, R.J., 2004. Proc R Soc London B, 271:S174-S176.
- Lenski, R.E. and May, R.M., 1994. J Theor Biol, 169:253-265.
- Luo, S. and Koelle, K., 2013. The American Naturalist, 181(S1):S58–S75. ISSN 0003-0147. doi:10.1086/669952.
- Mackinnon, M.J. and Read, A.F., 1999. *Evolution*, 53(3):689-703.
- Shirreff, G., Pellis, L., et al., 2011. PLoS Computational Biology, 7(10). ISSN 1553-734X. doi:10.1371/journal.pcbi.1002185. WOS:000297262700019.
- Velasco-Hernandez, J.X., Gershgorn, H.B., and Blower, S.M., 2002. Lancet, 2:487–493.
- Wonham, M.J., de Camino-Beck, T., and Lewis, M.A., 2004. Proc R Soc London B. 271:501-507




HIV eco-evo dynamics

Conclusions 00

References


extras

Baseline results



900

Pairs plots



Sensitivity

