

Eco-evolutionary dynamics of pathogen virulence

Ben Bolker, McMaster University

Departments of Mathematics & Statistics and Biology

University of Puerto Rico - Rio Piedras

26 April 2016

Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions

Acknowledgements

People Daniel Park; Arjun Nanda and Dharmini Shah;
Christophe Fraser; Marm Kilpatrick; Anson Wong

Support NSF IRCEB grant 9977063; QSE³ IGERT; NSERC
Discovery grant

Outline

1 Overview

- The evolution of host-pathogen theory
- Toy models

2 Transient virulence and emerging diseases

- Overview
- Toy model
- Myxomatosis data

3 Transient virulence of HIV

- abc

4 Conclusions

Host-pathogen evolutionary biology

Why is it interesting?

- Intellectual merit
 - Coevolutionary loops
 - Cryptic effects
 - Eco-evolutionary dynamics (Luo and Koelle, 2013)
 - Cool stories
 - Lots of data (sometimes)
- Broader applications
 - Medical
 - Conservation and management
 - Outreach

Host-pathogen evolutionary biology

Why is it interesting?

- Intellectual merit
 - Coevolutionary loops
 - Cryptic effects
 - Eco-evolutionary dynamics (Luo and Koelle, 2013)
 - Cool stories
 - Lots of data (sometimes)
- Broader applications
 - Medical
 - Conservation and management
 - Outreach

Virulence: definitions

- General public: badness
- Plant biologists: infectivity
- Evolutionists: loss of host fitness
- Theoreticians: *rate* of host mortality
(mortality rate vs. case mortality vs. clearance)

Evolution of virulence evolution theory

Classical dogma monotonic trend toward avirulence

Ewald era virulence as an evolved (adaptive) trait. Tradeoff theory, modes of transmission.

post-Ewald more formal tradeoff models, mostly based on \mathcal{R}_0 optimization. Adaptive dynamics

- Now**
- tradeoff backlash
 - within-host dynamics/multi-level models
 - eco-evolutionary dynamics
 - host effects: resistance vs tolerance vs virulence

Evolution of virulence evolution theory

Classical dogma monotonic trend toward avirulence

Ewald era virulence as an evolved (adaptive) trait. Tradeoff theory, modes of transmission.

post-Ewald more formal tradeoff models, mostly based on \mathcal{R}_0 optimization. Adaptive dynamics

- Now
- tradeoff backlash
 - within-host dynamics/multi-level models
 - eco-evolutionary dynamics
 - host effects: resistance vs tolerance vs virulence

Evolution of virulence evolution theory

Classical dogma monotonic trend toward avirulence

Ewald era virulence as an evolved (adaptive) trait. Tradeoff theory, modes of transmission.

post-Ewald more formal tradeoff models, mostly based on \mathcal{R}_0 optimization. Adaptive dynamics

- Now
- tradeoff backlash
 - within-host dynamics/multi-level models
 - eco-evolutionary dynamics
 - host effects: resistance vs tolerance vs virulence

Evolution of virulence evolution theory

Classical dogma monotonic trend toward avirulence

Ewald era virulence as an evolved (adaptive) trait. Tradeoff theory, modes of transmission.


post-Ewald more formal tradeoff models, mostly based on \mathcal{R}_0 optimization. Adaptive dynamics

- Now**
- tradeoff backlash
 - within-host dynamics/multi-level models
 - eco-evolutionary dynamics
 - host effects: resistance vs tolerance vs virulence


Basic tradeoff theory: assumptions

- Homogeneous, non-evolving hosts
- No superinfection/coinfection
- Horizontal, direct transmission
- Tradeoff between *rate* of transmission and length of infectious period
- Infectious period $\propto 1/\text{clearance}$
(= recovery+*disease-induced mortality*+*natural mortality*)


Tradeoffs, \mathcal{R}_0 , and r



Tradeoffs, \mathcal{R}_0 , and r



Tradeoffs, \mathcal{R}_0 , and r




Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions


Epidemiological model

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, $N = 1$ (time units of host lifespan)




Epidemiological model

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, $N = 1$ (time units of host lifespan)




Epidemiological model

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, $N = 1$
(time units of host lifespan)



Epidemiological model

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, $N = 1$ (time units of host lifespan)



The model (2): evolutionary dynamics

Incorporate *trait dynamics*

Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value ($\bar{\alpha}$)
and ecological context (proportion susceptible)
- Constant additive genetic variance V_g
- Trait evolves toward increased fitness:
rate proportional to $\Delta\text{fitness}/\Delta\text{trait}$

Alternatives:

multi-strain, adaptive dynamics, PDEs, agent-based models ...

The model (2): evolutionary dynamics

Incorporate *trait dynamics*

Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value ($\bar{\alpha}$)
and ecological context (proportion susceptible)
- Constant additive genetic variance V_g
- Trait evolves toward increased fitness:
rate proportional to $\Delta\text{fitness}/\Delta\text{trait}$

Alternatives:

multi-strain, adaptive dynamics, PDEs, agent-based models ...

The model (2): evolutionary dynamics

Incorporate *trait dynamics*

Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value ($\bar{\alpha}$) and ecological context (proportion susceptible)
- Constant additive genetic variance V_g
- Trait evolves toward increased fitness: rate proportional to $\Delta\text{fitness}/\Delta\text{trait}$

Alternatives:

multi-strain, adaptive dynamics, PDEs, agent-based models ...

The model (2): evolutionary dynamics

Incorporate *trait dynamics*

Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value ($\bar{\alpha}$) and ecological context (proportion susceptible)
- Constant additive genetic variance V_g
- Trait evolves toward increased fitness: rate proportional to $\Delta\text{fitness}/\Delta\text{trait}$

Alternatives:

multi-strain, adaptive dynamics, PDEs, agent-based models ...

The model (2): evolutionary dynamics

Incorporate *trait dynamics*


Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value ($\bar{\alpha}$) and ecological context (proportion susceptible)
- Constant additive genetic variance V_g
- Trait evolves toward increased fitness: rate proportional to $\Delta\text{fitness}/\Delta\text{trait}$


Alternatives:

multi-strain, adaptive dynamics, PDEs, agent-based models ...

Evolutionary dynamics, cont.



Evolutionary dynamics, cont.



A graph showing the relationship between Transmission (Y-axis) and Virulence (X-axis). The graph displays three curves representing different parameter values for c and γ :

- Blue curve: $c=2, \gamma=2$
- Black curve: $c=1, \gamma=2$
- Red curve: $c=1, \gamma=3$

The equation $\beta(\alpha) = c\alpha^{1/\gamma}$ is shown at the top of the graph.

Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 **Transient virulence and emerging diseases**
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions

(Why) are emerging pathogens more virulent?

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

- Sampling bias
- Lower host resistance/tolerance
- High transmission → frequent coinfection → selection for virulence
- Disease-induced decrease in host density → selection for lowered virulence (Lenski and May, 1994)

(Why) are emerging pathogens more virulent?

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

- Sampling bias
- Lower host resistance/tolerance
- High transmission → frequent coinfection → selection for virulence
- Disease-induced decrease in host density → selection for lowered virulence (Lenski and May, 1994)

(Why) are emerging pathogens more virulent?

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

- Sampling bias
- Lower host resistance/tolerance
- High transmission → frequent coinfection → selection for virulence
- Disease-induced decrease in host density → selection for lowered virulence (Lenski and May, 1994)

(Why) are emerging pathogens more virulent?

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

- Sampling bias
- Lower host resistance/tolerance
- High transmission → frequent coinfection → selection for virulence
- Disease-induced decrease in host density → selection for lowered virulence (Lenski and May, 1994)

Transient virulence

Selection differs between the **epidemic** and **endemic** phases of an outbreak (Frank, 1996; Day and Proulx, 2004)

endemic phase selection for per-generation offspring production:
maximize $\mathcal{R}_0, \beta N / (\alpha + \mu)$

epidemic phase selection for per-unit-time offspring production:
maximize $r, \beta N - (\alpha + \mu)$

Transient virulence

Selection differs between the **epidemic** and **endemic** phases of an outbreak (Frank, 1996; Day and Proulx, 2004)

endemic phase selection for per-generation offspring production:
maximize $\mathcal{R}_0, \beta N / (\alpha + \mu)$

epidemic phase selection for per-unit-time offspring production:
maximize $r, \beta N - (\alpha + \mu)$

Transient emerging virulence


When a parasite previously in eco-evolutionary equilibrium emerges in a new host population (at low density) it will show a transient peak in virulence as it spreads

How big is the peak? Does it matter?


Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 **Transient virulence and emerging diseases**
 - Overview
 - **Toy model**
 - Myxomatosis data
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions


Example



Response variables



Peak height




Estimates for emerging pathogens

Order of magnitude estimates for some emerging high-virulence pathogens:

Pathogen	\mathcal{R}_0^*	α^*	Reference
SARS	3	640	Anderson et al. (2004)
West Nile	1.61–3.24	639	Wonham et al. (2004)
HIV	1.43	6.36	Velasco-Hernandez et al. (2002)
myxomatosis	3	5	Dwyer et al. (1990)

Emerging pathogens: where are we?

$CV_g = 0.5$, $I(0) = 10^{-3}$ (middle panel):




Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 **Transient virulence and emerging diseases**
 - Overview
 - Toy model
 - **Myxomatosis data**
- 3 Transient virulence of HIV
 - abc
- 4 Conclusions


Overview

- Mosquito-borne viral disease of rabbits
- Benign in South American rabbits, quickly fatal in European rabbits
- Well characterized (Fenner et al., 1956; Dwyer et al., 1990)


Myxomatosis tradeoff curve




Myxomatosis grades vs. time




Myxomatosis variance vs. time




Myxomatosis virulence dynamics: power-law tradeoff




Myxomatosis virulence dynamics: realistic tradeoff




Myxo virulence: equilibrium start, power-law tradeoff



Myxo virulence: equilibrium start, realistic tradeoff



Phage dynamics (Berngruber et al., 2013)



Outline

1 Overview

- The evolution of host-pathogen theory
- Toy models

2 Transient virulence and emerging diseases


- Overview
- Toy model
- Myxomatosis data

3 Transient virulence of HIV

- abc

4 Conclusions


HIV tradeoff curve (Fraser et al., 2007)



HIV virulence dynamics (Shirreff et al., 2011)

- virulence/transmission from set-point viral load (SPVL)
- Fraser et al. (2007) : plausible tradeoff observed from Rakai data
 - SPVL heritable
 - correlated with rate of progression to AIDS
 - correlated with probability (rate) of within-couple transmission


Model results




Model variations

- 6 models:
 - extra-pair contact (yes/no) ×
 - instantaneous partnership formation (yes/no)
- implicit (Shirreff/Hollingsworth) model
- random-mixing model

Virulence trajectories (envelopes)



Univariate distributions



Conclusions Open questions

- Eco-evolutionary dynamics is still a reasonable framework (Alizon et al., 2009; Luo and Koelle, 2013)
- need to know: genetic variance, shapes of tradeoffs
- theory meets molecular biology: mutations of large effect vs. quantitative variability
- next steps in connecting with data? general theory?


Crome (1997) on theory

When we regard theories as tight, real entities and devote ourselves to their analysis, we can limit our horizons and, worse, attempt to make the world fit them. A lot of ecological discussion is not about nature, but about theories, generalizations, or models supposed to represent nature . . .

References

- Abrams, P.A., 2001. *Ecol Lett*, 4:166–175.
- Alizon, S., Hurford, A., et al., 2009. *J. Evol. Biol.*, 22:245–259. doi:10.1111/j.1420-9101.2008.01658.x.
- Anderson, R.M., Fraser, C., et al., 2004. *Phil Trans R Soc London B*, 359(1447):1091–1105.
- Berngruber, T.W., Froissart, R., et al., 2013. *PLoS Pathog*, 9(3):e1003209. doi:10.1371/journal.ppat.1003209.
- Blower, S.M., Hartel, D., et al., 1991. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 331(1260):171–187. ISSN 0962-8436, 1471-2970. doi:10.1098/rstb.1991.0006.
- Champredon, D., Bellan, S., and Dushoff, J., 2013. *PLoS ONE*, 8(12):e82906. doi:10.1371/journal.pone.0082906.
- Crome, F.H.J., 1997. In W.F. Laurance and J. Richard O. Bierregard, editors, *Tropical Forest Remnants: Ecology, Management and Conservation of Fragmented Communities*, chapter 31, pages 485–501. University of Chicago Press, Chicago.
- Day, T. and Proulx, S.R., 2004. *Amer Nat*, 163(4):E40–E63.
- Diekmann, O. and Heesterbeek, J.A.P., 2000. *Mathematical Epidemiology of Infectious Diseases: Model Building, Analysis and Interpretation*. Wiley. ISBN 978-0-471-49241-2.
- Dwyer, G., Levin, S., and Buttel, L., 1990. *Ecol Monogr*, 60:423–447.
- Ebert, D., 1998. *Science*, 282(5393):1432–1435.
- Fenner, F., Day, M.F., and Woodroffe, G.M., 1956. *J Hyg (London)*, 54(2):284–302.
- Frank, S.A., 1996. *Q Rev Biol*, 71(1):37–78.
- Fraser, C., Hollingsworth, T.D., et al., 2007. *PNAS*, 104:17441–17446.
- Herbeck, J.T., Mittler, J.E., et al., 2014. *PLoS Computational Biology*, 10(6):e1003673. doi:10.1371/journal.pcbi.1003673.
- Hollingsworth, T., Anderson, R., and Fraser, C., 2008. *Journal of Infectious Diseases*, 198(5):687–693. ISSN 0022-1899, 1537-6613. doi:10.1086/590501.
- Knell, R.J., 2004. *Proc R Soc London B*, 271:S174–S176.
- Lenski, R.E. and May, R.M., 1994. *J Theor Biol*, 169:253–265.
- Luo, S. and Koelle, K., 2013. *The American Naturalist*, 181(S1):S58–S75. ISSN 0003-0147. doi:10.1086/669952.
- Mackinnon, M.J. and Read, A.F., 1999. *Evolution*, 53(3):689–703.
- Shirreff, G., Pellis, L., et al., 2011. *PLoS Computational Biology*, 7(10). ISSN 1553-734X. doi:10.1371/journal.pcbi.1002185. WOS:000297262700019.
- Velasco-Hernandez, J.X., Gershgorin, H.B., and Blower, S.M., 2002. *Lancet*, 2:487–493.
- Wonham, M.J., de Camino-Beck, T., and Lewis, M.A., 2004. *Proc R Soc London B*, 271:501–507.

Baseline results



Overview
○○○○○○○○
○○○○○○


Emerging disease
○○○○
○○○○○○
○○○○○○○○○○○○

HIV eco-evo dynamics
○○○○○○○○○○○○○○

Conclusions
○○

References

Pairs plots



oooooooo
oooooooo

oooo
oooooooo
oooooooooooo

oooooooooooo

oo

Sensitivity

