


# Week 6 Report

Spring 2019  
Sam Frederick

# Ellipticity Calculations: Preparation

- Spent time finding sources with similar procedures
  - Gao et al. 2018 - Interesting paper which attempts to compute ellipticity for magnetars via Soft-Gamma Ray Burst sources.
  - May be able to compare results with their findings.
  - Also provides analytic equations for detector wave strain sensitivity for LIGO and the future Einstein Telescope.
  - Added raw, unsmoothed wave strain data of Advanced LIGO via <https://dcc.ligo.org/LIGO-G1801949/public>





# EOS Work

Ideally, I'd like to update the equation of state I use for the computational model in order to be more physically accurate

- Came across TOV solver code via <http://www.ictp-saifr.org/schoolgr/Lecture2Creighton.pdf>
  - Probably should check to make sure it's okay to use code?
  - Required a bit of modification to get working.
  - Basic premise: Use script to solve TOV equations for a piecewise polytrope and plot output (density and pressure) vs. radius.

# Verification of working code:



# Analytic Expression for SLy EOS

- Input:  $\log(\text{density})$
- Output:  $\log(\text{pressure})$
- Because analytic, could be used in place of piecewise polytrope for TOV code.
- Need to find inverse to compute pressure  $\longrightarrow$  density
  - Inverse of Taylor series expansion in Mathematica

$$\zeta = \frac{a_1 + a_2\xi + a_3\xi^3}{1 + a_4\xi} f_0(a_5(\xi - a_6)) \\ + (a_7 + a_8\xi) f_0(a_9(a_{10} - \xi)) \\ + (a_{11} + a_{12}\xi) f_0(a_{13}(a_{14} - \xi)) \\ + (a_{15} + a_{16}\xi) f_0(a_{17}(a_{18} - \xi)).$$