

Abstract

Title of Dissertation: γ -Ray Studies of Stellar Graveyards:
Fermi-LAT Observations of Supernova Remnants
and Spatially Extended Emission

Jamie Michael Cohen, Doctor of Philosophy, 2016

Dissertation directed by: Doctor Elizabeth Hays
Astroparticle Physics Laboratory, Code 661
NASA Goddard Space Flight Center

Professor M. Coleman Miller
Department of Astronomy
University of Maryland

Here I shall abstract!

**γ -Ray Studies of Stellar Graveyards:
Fermi-LAT Observations of Supernova Remnants
and Spatially Extended Emission**

by

Jamie Michael Cohen

Dissertation submitted to the Faculty of the Graduate School of the
University of Maryland at College Park in partial fulfillment
of the requirements for the degree of
Doctor of Philosophy
2016

Advisory Committee:

Doctor Elizabeth Hays, Advisor
Professor M. Coleman Miller, Chair/Advisor
Professor Christopher S. Reynolds
Professor Derek C. Richardson
Professor Jordan Goodman, Dean's Representative

© Jamie Michael Cohen 2016

Preface

This thesis consists of 8 chapters [JAM: 9 if I have time thrown in other work I’ve done like SNR-MC, above 10 GeV] including an introduction, conclusion and three background chapters on supernova remnants, γ -ray emission theory and detection methods, and a description of the relevant aspects of the *Fermi* Gamma Ray Space Telescope.

Chapters 5 and 6 are, respectively, taken in part from “The First *Fermi*-LAT Supernova Remnant Catalog” and “2FHL: The Second Catalog of Hard Fermi-LAT Sources”, both published in *The Astrophysical Journal Supplement* in 2016. Both papers are large, catalog studies involving the entire LAT collaboration. The parts of those papers included in this dissertation are the those in which I had direct contributed to via analysis, writing, discussion. The text in Chapters 5 and 6 also expands on the work I did for those papers, and provides further detail on analysis not included in the papers.

Chapter 7 is the contents of a paper currently in preparation and under LAT team internal review. The title of this paper is to be “*Fermi*-LAT Observations of Extended Gamma-Ray Emission in the Direction of SNR G150.3+4.5” (Cohen et al. 2016) [JAM: to be published in?]. The paper is included in entirety in this thesis, including additional supplementary material not to be included in the journal article.

To Vanessa ♡

Acknowledgements

I should probably thank someone because I'm not a degenerate.

Contents

List of Tables	viii
List of Figures	ix
1 Introduction	1
1.1 Goooo γ -rays go!	1
1.2 I Think I Hate Most of the Section Titles :(.	2
1.3 Maybe None of the Chapters Need Introductions?	2
1.4 Dissertation Overview	2
2 Supernova Remnants: Theory and Observation	3
2.1 Introduction	3
2.2 Formation and Evolution	3
2.3 Morphology and Classification	4
2.4 Cosmic Ray SNR connection	4
2.5 Summary	4
2.6 Scratch	5
3 Gamma-ray Astronomy	6
3.1 Introduction	6
3.2 γ -ray Emission Mechanisms	7
3.3 Sources of γ -ray's	7
3.4 γ -ray Detection	7
3.5 Scratch	8
4 The <i>Fermi</i> Gamma-Ray Space Telescope and γ-ray Data Analysis	9
4.1 Introduction	9
4.2 The Large Area Telescope	10
4.3 γ -ray Data Analysis	11
4.3.1 Do I need subsections?	11
4.4 Scratch	11

5	Revealing the GeV Supernova Remnant Population: The First <i>Fermi</i>-LAT Supernova Remnant Catalog	12
5.1	Supernova Remnants at γ -ray Energies	12
5.2	The <code>pointlike</code> Maximum-Likelihood Package and <code>addSrcs</code>	17
5.3	Galactic Supernova Remnants	24
5.4	Analysis Methods	24
5.5	Data Selection	25
5.6	Input Source Model Construction	27
5.7	Comparison of Source Models with 2FGL	36
5.8	Detection Method	40
5.8.1	Localization, Extension, and Spectral Curvature	43
5.8.2	Fluxes and Upper Limits	46
5.9	Catalog Results	47
5.10	GeV SNRs in a Multiwavelength Context: Discussion Summary . . .	48
5.11	Conclusions	59
6	Extended Source Detection above 50 GeV: The 2FHL Catalog	61
6.1	Introduction	62
6.2	Analysis	63
6.2.1	Data Selection	63
6.2.2	Source Detection	64
6.3	Search for Spatially-Extended Sources	67
6.3.1	Extended Sources Previously Detected by the LAT	67
6.3.2	Newly Detected Extended Sources	68
6.4	The 2FHL Catalog	70
6.4.1	General Characteristics of 2FHL Sources	72
6.4.2	The 2FHL Galactic Source Population	73
6.4.3	Comparison with the H.E.S.S. Galactic Plane Survey	77
6.4.4	Extended Source Results	79
6.5	Summary	85
6.6	Scratch	86
7	SNR G150.3+4.5	88
7.1	Introduction	89
7.2	<i>Fermi</i> -LAT Observations and Analysis	89
7.2.1	Data Set and Reduction	89
7.2.2	Morphological Analysis	91
7.2.3	Spectral Analysis	95
7.3	Multiwavelength Observations and Analysis	97
7.3.1	HI	97
7.3.2	CO?	97

7.3.3	X-ray	98
7.4	Discussion and Results	98
7.4.1	What is it?	98
7.4.2	Distance Considerations	99
7.4.3	Nonthermal Modeling	99
7.5	Conclusions	99
8	SNR-MC, 10 GeV, and anything else?	100
9	Conclusions	101
	List of Symbols and Acronyms	102
	Bibliography	104

List of Tables

5.1	Distances to SNRs	56
5.1	Distances to SNRs	58
6.1	2FHL extended sources previously detected by the <i>Fermi</i> -LAT	84
6.2	New 2FHL extended sources	84

List of Figures

5.1	Third EGRET catalog all-sky map.	13
5.2	Histogram of the number of significant sources remaining in each of the 8 test region of interest (RoI) for iterations in which $\Delta(\log \mathcal{L}) < 8$. .	32
5.3	SNR Gamma Cygni flux evolution for successive addSrcs iterations .	33
5.4	SNR Gamma Cygni index evolution for successive addSrcs iterations	34
5.5	SNR Gamma Cygni extension evolution for successive addSrcs iterations	35
5.6	1-100GeV residual TS map for supernova remnant (SNR) W44 before running addSrcs and with 2FGL sources removed from the inner 3° radius.	36
5.7	1-100GeV residual TS map for SNR W44 after addSrcs has completed.	37
5.8	Comparison of the number of 2FGL sources with with the number of newly added input model sources.	39
5.9	Same as Figure 5.8, including only input model sources lying within 0.2° of a 2FGL source.	40
5.10	Distribution of SNR radio diameters from Green's catalog	42
5.11	Radio diameter of Green's catalog SNRs plotted against the fitted GeV diameter	49
5.12	Comparison of γ -ray and radio spectral flux densities for all SNRs and candidates.	50
5.13	Comparison of radio spectral index, α , and GeV photon index, Γ . . .	51
5.14	GeV index compared to published index measurements from Imaging Air Cherenkov Telescopes (IACTs).	53
5.15	Age versus GeV spectral index	54
5.16	The 1 – 100 GeV luminosity vs. D^2 for SNRs with distance measurement.	59
6.1	Adaptively smoothed count map in the 50 GeV–2 TeV band represented in Galactic coordinates and Hammer-Aitoff projection. The image has been smoothed with a Gaussian kernel whose size was varied to achieve a minimum signal-to-noise ratio under the kernel of 2. The color scale is logarithmic and the units are counts per $(0.1 \text{ deg})^2$.	64

6.2	Sky map, in Galactic coordinates and Hammer-Aitoff projection, showing the sources in the 2FHL catalog classified by their most likely association.	71
6.3	Distribution of the spectral indices (<i>top panel</i>) and highest photon energy (<i>bottom panel</i>) of the Galactic sources (orange), extragalactic sources (green slash), and unassociated sources (brown dotted). The medians of the distributions are plotted with dashed, dash-dotted, and dotted vertical lines, respectively. Both plots show that a distinct population of hard-spectrum sources is of Galactic origin.	74
6.4	Adaptively smoothed count map showing the whole Galactic plane $0^\circ \leq l \leq 360^\circ$ at Galactic latitudes $-14^\circ \leq b \leq 14^\circ$ divided in four panels. The panels are centered at $l = 0^\circ, 90^\circ, 180^\circ$ and 270° , respectively. Detected point sources are marked with a cross whereas extended sources are indicated with their extensions. Only sources located at $-4^\circ \leq b \leq 4^\circ$ are explicitly named, plus the Crab Nebula.	75
6.5	Spectral energy distributions of four Galactic sources constructed by combining data from the 3FGL (green diamonds), 1FHL (blue circles), and 2FHL (red stars). We show the 3FGL extended source SNR IC 443 (<i>top left</i>), the new 2FHL extended source PSR J1420–6048 (<i>top right</i>), and two “dark accelerators” detected by H.E.S.S. at TeV energies (Carrigan et al. 2013, purple squares) without a previous LAT counterpart: HESS J1503–582 (<i>bottom left</i>) and HESS J1702–420 (<i>bottom right</i>).	80
6.6	Residual TS maps for the five new extended sources described in Chapter 6.4.4. Only the Galactic diffuse and isotropic emission are included in the model to highlight the location of emission not associated with the diffuse background. Circles indicate the extents of the fit disks. The x marker in the bottom panel (2FHL J0431.2+5553e) shows the location of a point source in the ROI.	87
7.1	Background subtracted residual TS map above 1 GeV with $0.1^\circ \times 0.1^\circ$ pixels for fixed index $\Gamma = 2$, centered on SNR SNR G150.3+4.5. The orange circle and translucent shading show the fit disk radius and 1σ errors, respectively, for the extended source, the orange cross shows the position of 3FGL J0426.7+5437 (included in the background model), and blue dashed circle is the extent of the radio SNR.	93
7.2	Same as Figure 7.1 but with disk in the background model [JAM: should this be a residual counts map instead?]	94

7.3	Spectral energy distribution for the extended source coincident with SNR SNR G150.3+4.5. [JAM: replace with gtlike SED when I have it]	96
7.4	Spectral energy distribution of 3FGL J0426.. [JAM: replace with gtlike SED when I have it]	97

Chapter 1

Introduction

“Maybe I’ll have a super relevant quote here!”

—by some awesome human, from *Some book*

1.1 Goooo γ -rays go!

In this thesis we...or should I start with the extreme environs line?

Overview of the entire thesis, why gamma-rays, why the Large Area Telescope (LAT), why SNR and pulsar wind nebula (PWN) and extended sources.

Higher energy studies with the LAT have been my focus since the beginning. Talk about what’s nice about staying above 1 GeV, 10 GeV, 50 GeV.

GeV TeV connection for 2FHL

Radio GeV for SNR cat (traces same particle population)

The advent of the LAT presents for the first time the capability to spectrally and spatially resolve SNR at GeV energies.

it is uniquely situated to address these issues

egret was mostly pointed observation instrumented that would sometimes dwell

on a spot for a couple of weeks, had a smaller field o view, didn't get as many photons (the LAT saw the entire EGRET sky in some short amount of time)

SNRsas sources of relativistic particles

Despite being the prime energy range to observe the effects of cosmic particle acceleration, the photon spectral energy distribution (SED) resulting from these overlapping emission channels are often difficult to spectrally distinguish from one another. [JAM: what's the point of this last sentence here? maybe no need to mention this now, i really just want to motivate GeV energies]

when talking about Energetic Gamma-Ray Experiment Telescope (EGRET)

With its unprecedented sensitivity and angular resolution above 1 GeV, the LAT provides for the first time the opportunity to distinguish SNR-emitted photons from their backgrounds, and to unambiguously detect and identify dozens of SNRs.

[JAM: maybe this is for the intro/abstract because it's a little vague]

The LAT is uniquely situated to address these goals and definitively detect and identify dozens SNRs

Thompson et al. (1993) gives the 68% containment radius as $\theta \leq 5.85^\circ (E_\gamma/100 \text{ MeV})^{-0.534}$

1.2 I Think I Hate Most of the Section Titles :(

1.3 Maybe None of the Chapters Need Introductions?

1.4 Dissertation Overview

Chapter 2

Supernova Remnants: Theory and Observation

2.1 Introduction

Why study SNRs, what they are, history of SNR, radio detections,

2.2 Formation and Evolution

-Stars die and explode, that energy is very quickly put into the surroundings

- snowplough, ST, radiative,

- what else?

How we detect gamma-rays from SNRs/PWNe in the Galaxy leads to and analysis section maybe?

2.3 Morphology and Classification

SNRs characterized by morphology and evolution properties

shell type, mixed morphology, filled center composite)

Since I eventually do these all plane surveys, what does the spatial distribution of them at radio look like?

Not sure how much to say about radio observations, x-ray, TeV

2.4 Cosmic Ray SNR connection

Give the whole, if 10% of energy of SN explosion goes into particle acceleration, we can explain cosmic rays

Particle acceleration and DSA

This leads to gamma-ray section

2.5 Summary

In this section we summarized the end phase of stellar evolution (just enough to motivate SNRs) and described the environs surrounding the supernova; development and phases of SNRs (and PWNe?). In particular we detailed the nonthermal emission mechanisms that produce γ -ray radiation, detection of young vs middle-aged(evolved, interacting with surroundings/dense medium), TeV detects younger typically, the troubles of detecting extension from them(?) something about different emission zones? Troubles disentangling hadronic from leptonic at γ -rays. γ -ray spectral and morphological features. Trends across the population wrt spectral shape/breaks, higher luminosity for interacting rems. Cosmic rays, using gamma-rays to probe CR population. So much of γ -ray astro is really about studying CRs,

how much to say about them?

2.6 Scratch

This chapter needs a different title. It's more focused on the specific sources being studied in this thesis. Galactic extended sources, SNRs, PWNe, but as in the SNRcat, not just extended SNRs, point-like SNRs as well.

Less focus on PWNe. Only give as much as I feel I need to support mentioning them a bit for 2FHL?

The focus of this section is supernova remnants in a gamma-ray context. Theory of evolution, what the gamma-ray emission is like, what we can learn from them individually. This leads to the 1st SNR cat section for what we can do with them ensemble

NOt sure I really need any PWN stuff yet

in 2FHL we detect some pwn. If including above 10gev work, they'll be there too. Much of the thesis is really about extended gamma-ray sources, but not sure how that fits into the title and chapters yet

Do I need to get into composite SNRs (composite means SNR + PWN) Maybe relevant for G150? Some things about interaction of reverse shock with PWN and crushing/reverberations of the PWN?

Montmerle (1979)

Chapter 3

Gamma-ray Astronomy

3.1 Introduction

Maybe this is not just gamma astro, but gamma astro of SNRS

The story of γ -ray's from astrophysical objects is a tale of the most extreme, energetic, and violent environments in our universe. Discovered by Paul Villard studying radiation from radium and named by Ernest Rutherford, who previously uncovered the nature of α and β radiation, γ -ray's are the highest named energy of light

more historical context instead of a separate section

what is a gamma-ray

why bother studying gamma-rays

probe of extreme environments

thermal means follows maxwellian (maxwell-Boltzman) distribution

[JAM: snrs as source of Galactic cosmic rays, and thus as drivers of Galactic evolution]

3.2 γ -ray Emission Mechanisms

Gamma-rays as a probe of cosmic rays and cosmic acceleration processes

gamma-ray astronomy as a proxy for studying cosmic rays and acceleration/diffusion processes. How much to get into CR.

Cosmic particle accelerators and γ -ray's accelerator plus target often

What's a CR, quick, general CR properties that are relevant to SNRs

why use gamma's to study CR

gamma-ray production mechanisms

myriad of gamma-ray generating mechanisms operate at this energy, so it's sometimes the only range that we can observe this emission in:

-Synch, -Bremss, -IC, - π^0 ,

3.3 Sources of γ -ray's

Maybe don't need this? The reason to is to say SNRs early on. Would I mention other sources to be complete?

SNRs as the primary source of Galactic CRs, order of mag (zwick?) energy from $0.1 \cdot E_{\text{SNR}}$ could account for energy in CRs in Galaxy

3.4 γ -ray Detection

Quick rehash of method of detecting γ -ray's? Or is this just about previous γ -ray detectors and the state of the γ -ray sky pre-Fermi? mention telescopes up to EGRET, bit of detail on EGRET and what the pre-Fermi γ -ray sky looked like, in particular in the context of SNRs, PWN, Galactic plane

gamma-ray telescopes leads into the LAT, Egret was predecessor , what it did and what were some relevant unanswered questions regarding supernova remnants

One of the primary goals of fermi was to identify these sources and the site of CR acceleration. But also to uniquely open this high energy window that where no other telescopes really operated Brief history of radio detection of SNRs. detection at other wavelengths. what we see at γ -rays?

Motivation for why to study them (Sturmer & Dermer 1995) (Esposito et al. 1996)

3.5 Scratch

Altho' many miles from bomb zero, Dr. Bruce Banner is bathed in the full force of the mysterious Gamma Rays!

Chapter 4

The *Fermi* Gamma-Ray Space Telescope and γ -ray Data Analysis

4.1 Introduction

The *Fermi* Gamma-Ray Space Telescope (*Fermi* hereafter), successor to the EGRET instrument on Compton Gamma-Ray Observatory (CGRO), was successfully launched into orbit around Earth on June 11 2008. *Fermi* consists of two instruments, the LAT and the Gamma-ray Burst Monitor (GBM).

The LAT, which is the primary instrument on *Fermi*, is a pair conversion telescope designed to detect photons from 20 MeV to greater than 1 TeV. Its standard mode of operation is a sky-survey mode in which it observes the entire sky every 3 hours. The GBM is designed to detect gamma-ray bursts (GRBs) in a waveband overlapping that of the LAT, and complementary in lowering that energy range. It is comprised of two types of scintillator detectors: two bismuth germanate crystals that operate from 150 keV to 30 MeV, and 12 sodium iodide crystals sensitive to photons between 8 keV and 1 MeV.

Combined the LAT and GBM make up a formidable observatory, spanning more than 8 decades in energy, and is currently the only instrument performing all-sky observation in this broad energy range. [JAM: maybe I don't need any gbm stuff? I mentioned it just to be complete about what fermi is, probably won't mention it again, and this last par doesn't really flow into the next]

4.2 The Large Area Telescope

The need for Fermi in the context of what EGRET did

What were open questions from EGRET era, state of γ -ray detection of SNRs, what question was Fermi deigned to answe

Description of the instrument [JAM: not sure this really goes here, separate section for what questions Fermi was designed to answer?]

track and reconstruct the path of Describe it's objectives and strengths over predecessors Details on the LAT and it's design, be sure to focus on things that particularly pertinent to the work I've done like what determines the PSF, thing about Pass 8 here maybe? Or maybe later on.

what science was it designed to answer

general capabilities

details about aspect of the LAT related to extended sources, what determines PSF

4.3 γ -ray Data Analysis

Why maximum likelihood, how it's formulated, implemented in the Science Tools, pointlike and the analysis for extended sources. Diffuse emission.

Four steps to going from observing the sky to final LAT analysis:

Instrument taking data: How we get to counts

Reconstruction : How we get photons

Likelihood: How to characterize sky using response functions, point source and diffuse modeling

Likelihood for ES: how to use likelihood methods to char and resolve sources measure extension

Section on diffuse emission

4.3.1 Do I need subsections?

4.4 Scratch

[JAM: Fermi goals: 1. Resolve the γ -ray sky: the origins of diffuse emission and the nature of unidentified sources: Source identification through good source localization, measurement of spectra across broad energy range, nearly continuous monitoring of the sky for temporal variability 2. Understand the mechanisms of particle acceleration in celestial sources] I'm not sure about this chapter yet. Maybe it's a general section on Analysis of Fermi data,

Chapter 5

Revealing the GeV Supernova Remnant Population: The First *Fermi*-LAT Supernova Remnant Catalog

5.1 Supernova Remnants at γ -ray Energies

[JAM: should I move much of this section to the SNR chapter?]By the end of the its science run, EGRET had detected 271 sources above 100 MeV, within a minimum detection significance of 4σ , 170 of which had no clear multiwavelength counterpart, with 81 of those unidentified lying within $|b| < 10^\circ$ of the Galactic plane (Hartman et al. 1999). The main hindrances to source identification were the numerous potential source counterparts (the EGRET point spread function (PSF) was energy dependent, with a 68% containment radius of $\sim 6^\circ$ at 100 MeV and smaller for higher energies) and the large EGRET error boxes. In addition to this, the primary

method for identifying a γ -ray source as an SNR is through a compatible angular extent with observations at some other wavelength, thus the ability to resolve emission from an SNR is vital to understanding the mechanisms therein giving rise to γ -rays. Figure 5.1 shows an EGRET all-sky map at $E > 100$ MeV where the preponderance of unidentified sources and locations thereof are made clear.




Figure 5.1: Third EGRET catalog all-sky map. Unidentified sources represented by triangles. Image courtesy of <https://heasarc.gsfc.nasa.gov/docs/cgro/images/epo/gallery/skymaps/>

In spite of the difficulties in EGRET source association, many studies have attempted correlating the unidentified EGRET sources with various Galactic populations. In particular, several authors found strong evidence for statistical correlation between SNRs and some of the low-latitude unidentified sources (Esposito et al.

1996; Romero et al. 1999; Sturmer & Dermer 1995). In a review of the state of potential SNR / EGRET associations, Torres et al. (2003) showed that there were 19 unidentified EGRET sources that had an SNR fall within its 95% error box. Performing Monte Carlo simulations of the population of EGRET sources, they determined that the chance probability for the 19 sources to be coincident with an SNR was 1.05×10^{-5} , implying a probability of 0.99998 that at least one of the associations is real. Despite the statistical correlation of EGRET sources with SNRs, there were no definitive associations of an SNR with any EGRET sources.

As the successor to EGRET, the LAT was designed to improve upon its predecessor in a multitude of areas relevant to detecting SNRs (Ackermann et al. 2012c; Atwood et al. 2009). The LAT has a much improved angular resolution (68% single-photon containment radius $\sim 0.4^\circ$ at 1 GeV for photons with the best quality direction reconstruction, PSF3 event type, compared to $\sim 1.7^\circ$ for EGRET at the same energy), necessary to resolve SNRs as extended objects. The LAT also benefits from a superior sensitivity due to a combination of the improved PSF, larger peak effective area ($> 9000 \text{ cm}^2$ vs. $\sim 1500 \text{ cm}^2$), wider field of view (FoV) (2.4 sr, which is nearly 5 times that of EGRET), and deeper, more-uniform sky exposure (afforded by the LAT's scanning observations as opposed to EGRET's pointing operation).

This bump in sensitivity results in the LAT detecting considerably more sources than EGRET. Remarkably, within its first three months of commission, the LAT detected 205 sources above 10σ significance (Abdo 2009), and by 11 months, 1451 sources above 4σ (Abdo et al. 2010a), compared to the aforementioned 271 over the entire EGRET mission. In fact, over its lifetime, EGRET detected a total of about 1.5×10^6 cosmic photons (Thompson et al. 1993), while as of March 2016, the LAT has detected $\sim 863 \times 10^6$ [JAM: change this number in June] source class photons. The LAT's point-source sensitivity peaks between 1 and 10 GeV, depending on

location on the sky. [JAM: plot comparing lat to egret sensitivity?]. With its increased sensitivity and higher energy range (up to ~ 2 TeV with the recent Pass 8 event reconstruction improvements, which is nearly an order of magnitude higher than EGRET), the LAT is uniquely situated to study the γ -ray morphology and spectra of SNRs.

Both energetic lepton interactions (i.e. inverse compton (IC) radiation of relativistic electrons interacting with ambient photon fields, and nonthermal bremsstrahlung) and hadronic processes (π_0 decay γ -rays from cosmic ray (CR) protons encountering surrounding nuclei) produce spectra observable at γ -ray energies (see Chapter 3 for details). While the IC generating electron population is also observable through emission of radio synchrotron photons, the proton-proton interaction solely emits γ -rays. Despite being the prime energy range to observe the effects of cosmic particle acceleration, complexities at the lower LAT energy range stymie SNR morphology studies.

The LAT detects a strong, soft band of diffuse emission in the Galactic plane due to the interactions of CRs with interstellar material. This bright diffuse radiation combined with the multiple potential emission scenarios, broadening PSF at decreasing energy, and a high source density in the plane can make it difficult to spatially disentangle sources observed by the LAT. To circumvent these difficulties, the majority of the analyses undertaken in this thesis are focused on the $E \geq 1$ GeV energy range. This energy band is ideal for probing the properties of the accelerated particle populations present in the SNR environment. Studies of SNRs above 1 GeV benefit from finer LAT PSF, striking a balance between minimizing the diffuse contribution, maximizing photon sensitivity, and retaining good photon statistics. Furthermore, evolved SNRs exhibit a spectral break between 1-10 GeV (Hewitt & Lemoine-Goumard 2015). Explanations for the break range from Alfvén wave

evanescence generated by collisions of partially ionized material in molecular clouds (MCs) overtaken by SNR shocks (Malkov et al. 2011), reflected shocks in clouds Inoue et al. (2010), and energy-dependent diffusion from shocks Ohira et al. (2011). Studying SNRs in this energy range hones our capability to tackle several goals set out by the *Fermi* team when the mission was conceived.

Two of the primary science goals of the LAT are to 1. resolve the γ -ray sky, uncovering the nature of the unidentified sources detected by EGRET, and 2. to understand the mechanisms of cosmic particle acceleration (Atwood et al. 2009). In this chapter, we describe our efforts towards addressing these questions by studying the γ -ray emission coincident with sources comprising the population of known radio emitting SNRs.

Prior to this work, several individual studies with the LAT had successfully resolved spatially extended emission from SNRs (Acero et al. 2015, and references therein), yet no systematic analysis leveraging the LAT’s full-sky coverage had thus been attempted. We performed for the first time a uniform study of the SNRs in aggregate to measure the properties common to these objects. An understanding of these common characteristics allows us to assess SNRs as a class of γ -ray and CR emitting objects and serves as the impetus for this uniform analysis of the known Galactic SNRs. We report here on the published results from the First *Fermi*-LAT Supernova Remnant Catalog (SNRcat) (Acero et al. 2016).

[JAM: 2FGL only had 7 ID’d SNR, 4 snr, 58 spp. 3FGL had 12 SNR, 11 snr, 49 spp. I’m not sure what made some snr vs. spp. They must have all been point sources right? No known radio pwn, psr?]

5.2 The pointlike Maximum-Likelihood Package and addSrcs

As described in Chapter 4, maximum-likelihood analysis is the ideal method for determining the properties of LAT-observed sources due to the “counting-experiment” nature of *Fermi*-LAT. The standard maximum-likelihood tools for analyzing LAT data are implemented via the *Fermi* Science Tools, and in particular `gtlike`. Despite being the optimum method, likelihood analysis of LAT data is complex due to the highly non-linear performance of the instrument and can be computationally expensive. It is necessary to manage the data and response of the telescope as well as the source and background models. Furthermore, due to the broadening of the PSF at low energies, even when studying a single source, it is necessary to include in the model descriptions of multiple surrounding sources. The `pointlike` binned maximum likelihood package was created to ameliorate some of these issues. Described in detail in Kerr (2010), `pointlike` is an alternate likelihood analysis framework (a collection of Python modules with additional wrappers for accessing C++ code), designed to be interactive and rapidly evaluate likelihoods.

There are several ways in which `pointlike` improves in efficiency compared to the Science Tools. It saves computational time, while sacrificing some accuracy, with several assumptions and approximations, such as the PSF not varying strongly with photon incidence angle (allowing a single PSF for each individual bin), and sources having a steady flux in individual short time bins. Most importantly though, `pointlike` varies the size of spatially binned `HEALPix` pixels (Górski et al. 2005) according to energy. The PSF at lower energies is large and each energy bin can contain multiple counts, while at higher energies, the PSF shrinks and many pixels will not contain even a single count. `pointlike` creates `HEALPix` bins that are

approximately the size of the PSF at a given energy, and disregards empty bins to speed up the likelihood calculation.

In addition to these computational, time saving efficiencies, tools to analyze spatially extended sources have also been built into the `pointlike` framework. Studying the position and extension of an extended source, while possible with the standard *Fermi* Science Tools, is a cumbersome process. `gtlike` is not capable of simultaneously maximizing the likelihood of a source’s spectral and spatial parameters, so to assess the morphology of a source, an iterative process of fitting a spatially fixed source’s spectrum and then varying the sources centroid and extension is required. To address the issues that arise when studying individual extended sources, Lande et al. (2012) developed and validated spatial likelihood fitting tools for `pointlike`, taking advantage of the time-saving properties built therein.

To fit the position and extension of a source, `pointlike` assumes that the spatial and spectral distribution of a source’s expected photon distribution are separable. The extended source’s shape is convolved with the LAT PSF (which is a function of energy) to determine the expected distribution. Then, the `minuit` numerical minimization library (James & Roos 1975) is used to maximize the likelihood of the model by simultaneously varying the spectrum, extension, and position of the source. Various geometric surface brightness models are built into `pointlike`, including, but not limited to a uniform intensity disk and ring, and a 2D Gaussian, with radially and non-radially symmetric versions of each. Akin to the speed optimizations mentioned previously, for radially symmetric sources, `pointlike` calculates the angular integral of a source’s expected photon distributions analytically to save computational time.

The significance of extension of a source is determined by using the likelihood ratio test (LRT). The LRT is a statistical method to assess the goodness-of-fit of two different models. The likelihood (as described in Chapter §4.3 [JAM: do this,

reference cash,mattox, fisher for first use of word likelihood?]) is calculated for two models, one of which can be reduced to the other hypothesis under certain conditions. If the more complex model can be reduced to the simpler model (called the null hypothesis), we say the simpler hypothesis is nested within the more complex. In the LRT, the test statistic (TS) is defined as:

$$\text{TS} \equiv 2 \log(\mathcal{L}(\text{H}_1) / \mathcal{L}(\text{H}_0)), \quad (5.1)$$

with H_1 being the more complex hypothesis and H_0 the null. Applying this to the hypothesis of a spatially extended source, we can calculate the significance of a source being extended compared to that of the source being modeled as point source as:

$$\text{TS}_{\text{ext}} \equiv 2 \log(\mathcal{L}_{\text{es}} / \mathcal{L}_{\text{ps}}) = \text{TS}_{\text{es}} - \text{TS}_{\text{ps}}, \quad (5.2)$$

Mattox et al. (1996) detail how by Wilk's theorem, the TS for detection of a point source (with the null hypothesis being that with no source present, or 0 flux) should be distributed as a chi-squared distribution in the null hypothesis for an increasing sample size, which for photon counting experiments, is the number of events relevant to the parameter being estimated. Specifically,

$$\text{PDF}(\text{TS}) = 1/2 \chi_1^2, \quad (5.3)$$

where $\text{PDF}(\text{TS})$ is the probability distribution function for obtaining a specific value of TS and χ_1^2 is the chi-squared distribution for one degrees of freedom. The factor of $1/2$ arises from the fact that the flux of a source is not permitted to be zero, and since negative and positive fluctuations in a parameter's value contribute equally to the TS, half of the distribution is lost with the positive flux restriction. The significance of detection is oft quoted as $\sigma \approx \sqrt{\text{TS}}$, which is strictly valid only

for χ_1^2 . More generally, when comparing the likelihood of two models with n degrees of freedom between them, equation 5.3 applies, but using χ_n^2 for n degrees of freedom versus one.

Lande et al. (2012), extended (and verified) this definition of TS to calculating the significance of extension, replacing the source flux with its radius. The uncertainty of the extension parameter is estimated by fixing a source’s position while varying the extension until the log likelihood decreases by 1/2 from the maximum value (i.e. 1σ errors). [JAM: use that TS vs extension figure I didn’t include in the G150 paper to demonstrate where the TS drops by half from the peak?] A similar procedure is used to estimate the errors on a source’s position, but rather, fixing the extension and spectrum (Nolan et al. 2012). While `pointlike` is an tool for the analyses described above, `gtlike` is still the go-to for estimating the best-fit spectral parameters since it is expected to be slightly more accurate than `pointlike` since it makes approximations. For the studies in this thesis, we used `pointlike` to calculate extension and source positions, and then use the `pointlike` results as a starting point for the likelihood parameter estimation of spectra with `gtlike`.

With its efficient likelihood calculations, and ability to simultaneously fit both the spectral and spatial parameters of a source, `pointlike` is ideally suited for large-scale studies (like the all-sky analyses performed for the LAT point source catalogs), and analyses requiring several iterations. Studying the γ -ray emission from the population of Galactic SNRs is precisely the sort of analysis that `pointlike` was designed to perform. To attain the best understanding of a source of interest, the best characterization of the corresponding RoI is necessary. In particular, to understand the GeV emission from a potentially extended SNR, it is important to quantify the surrounding emission because of the steep energy-dependence of the LAT PSF. This can be especially challenging in dense source and strong diffuse-

dominated regions, like the Galactic plane where the SNRs we are studying lie. We have developed an automated method for systematically locating and modeling all potential point and/or extended sources in an RoI using `pointlike`.

A typical LAT analysis starts by including all sources from the most recent LAT point source catalog and modifying the RoI to suit one's needs. Unmodeled emission can arise if using a dataset longer than that used in the most recent catalog or by focusing on a different energy range compared to that of the catalogs. We created a Python subclass of the primary `pointlike` analysis object (which works within that framework, inheriting all of the class' features, while adding new functionality) to systematically and uniformly characterize sources in an RoI by finding residual, unmodeled emission in the region and iteratively add sources to the RoI to account for this emission. The main module in the designed codebase was dubbed `addSrcs`.

The general work flow of `addSrcs` is to start with a model of the RoI, including some combination of the diffuse background components, point and extended sources. `addSrcs` reads in a residual TS map or creates one on the fly if none is passed in. Residual emission is detected by finding the peak emission in the TS map and adding a source to the existing RoI at the position of the peak pixel. Either all point or extended sources can be iteratively detected and added to the RoI. For the SNRcat, we exclusively ran `addSrcs` in point source mode. Chapter 6 provides an application of `addSrcs` for extended sources.

In point source mode, a point source with a power law (PL) spectrum is added to the model of the region, a likelihood fit of the RoI is performed, and subsequently, the source's position is localized. Similarly, in extended mode a PL extended source (of any morphological form included in `pointlike`) is added to the RoI with a small seed radius, and the spatial parameters of the newly added source are fit simultaneously with the spectra of the other sources already in the model. If the source has

$TS_{\text{ext}} < 16$ (equivalent to a 4σ extension significance and validated through simulations in Lande et al. (2012) as a reasonable extension detection significance), the extended source is replaced with a point source and the iteration continues as in point source mode. To extend the functionality of `addSrcs` and make it generally applicable to a multitude of LAT analyses, several optional methods were built in.

One such option is to test the newly added source for signs of spectral curvature (described further in Chapter 5.6). If the source is found to show significant spectral curvature, the appropriate curved spectral model is retained, otherwise, we revert to the best-fit PL model. Another option provided is to fix the new sources spectrum if it is within a given angular separation of the center of the RoI to limit the number of free parameters for the likelihood fit and aid in proper convergence. If the source of interest being studied is not central in the RoI it might be beneficial to free the spectral parameters of sources within a given distance of the newly added source rather than from the center of the RoI. This choice was also built into `addSrcs`. Further, we included an option to refit the extension of any extended source already in the model at each iteration if they are within a given distance of the new (point or extended) source. Due to the broad size of the PSF, nearby source spectra can be influenced by each other, (particularly for extended sources) so the iterative procedure allows the likelihood to relax to a preferred value when adding new sources.

Throughout the `addSrcs` process, various checks are performed to ensure that parameter values are reasonable, the likelihood fit converges, and the procedure is generally running as expected. The range of permissible fit values for a parameter can be limited, the values of the parameters themselves can be fixed, or a consistently poorly-fit source can be automatically removed from the model. During the source localization step, if the fit goes awry and the source wanders too far from its initial

position, the position of the source can be rolled back to its starting location and fixed. Checks were also included to keep track of the Galactic diffuse and isotropic emission models to ensure they were adequately fit.

The penultimate step of the iteration is to produce various diagnostic plots and output information about the fits, the spectral and spatial parameters of each source in the model, and other relevant information such as the TS of the source and loglikelihood of the fit. Finally, a new residual TS map of the region is created and the source addition procedure repeats until a given threshold in TS is reached. The peak pixel TS found in the residual TS map does not necessarily decrease monotonically, as is expected of the actual TS of successive sources as more of the emission is accounted for and the model improved. Since the peak pixel TS can fluctuate a bit, to ensure that we do not miss significant sources in the RoI, we continue adding sources until the TS threshold is reached for some number of successive sources (discussed further in 5.6). After sources are no longer being added to the region, we iteratively remove sources with TS less than a given threshold (typically $TS < 16$, again see Chapter 5.6) starting with the lowest TS sources first. As each source is removed, we refit the RoI, including any extended sources close to the removed source. When the TS of all sources in the RoI are above threshold, we deem the emission in the RoI to be sufficiently characterized.

In the following sections, we detail the application of `addSrcs` to studying the GeV Galactic SNR population and describe the analysis and results presented in Acero et al. (2016).

5.3 Galactic Supernova Remnants

In this work we focus on the 279 currently known Galactic SNRs. They are derived from the 274 SNRs noted in the catalog of Green (2009, hereafter Green’s catalog), plus five additional SNRs identified following its publication. All but 16 of these SNRs have been identified by their radio synchrotron emission, so their centroids and extensions are primarily determined from the radio. When the radio detection is not securely identified through the synchrotron emission, positional information is obtained from the optical, X-ray, or TeV observations that identified the SNR, as noted in Green’s catalog. The catalog is thought to be complete down to a 1 GHz radio surface brightness limit of $\approx 10^{-20} \text{ W m}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$ (i.e. 1 MJy sr $^{-1}$). However, selection effects are known to bias radio surveys against the identification of radio faint and small angular size remnants (Brogan et al. 2006; Green 2004). We note that as this work neared completion, a revised catalog of 294 SNRs was published (Green 2014), representing only a small increase ($< 10\%$) over the previous catalog.

5.4 Analysis Methods

To systematically analyze the *Fermi*-LAT γ -ray data, we apply a maximum likelihood (Mattox et al. 1996) framework to RoIs centered on known SNRs (Green 2009). For each SNR, we begin by constructing a model for the spectral and spatial dependence of the γ -ray emission which includes significant point sources in the RoI. We then test for the existence of a γ -ray source near the center. This includes determining the most likely position and extension of the candidate source and testing for spectral curvature, rather than assuming it follows a PL across the energy range studied. In cases where we find no significant source associated with the SNR,

we calculate upper limits on the flux. We calculate both statistical and systematic errors, where the latter are estimated from both the uncertainty in the effective area and the effects of changing the interstellar emission model (IEM), which accounts for γ -rays produced by CR interactions with interstellar gas and radiation fields in the Milky Way.

This analysis uses both the standard Science Tools (version 09-32-05), including `gtlike`, and the `pointlike` analysis package (Kerr 2010) which has been developed and verified for characterizing source extension for *Fermi*-LAT data (Lande et al. 2012). §5.5 describes our data selection; §5.6 details our new method for automatically finding point sources in the *Fermi*-LAT γ -ray emission; and §5.8 discusses the detection method.

5.5 Data Selection

This catalog was constructed using 3 years of LAT survey data from the Pass 7 (P7) “Source” class and the associated P7V6 instrument response functions (IRFs). This interval spans 36 months, from 2008 August 4 to 2011 August 4 (mission elapsed time 239557417 – 334108806). The Source event class is optimized for the analysis of persistent LAT sources, and balances effective area against suppression of background from residual misclassified charged particles. We selected only events within a maximum zenith angle of 100° and use the recommended filter string “DATA_QUAL==1 && LAT_CONFIG==1” in `gtmktime`¹. The P7 data and associated products are comparable to those used in the other γ -ray catalogs employed in this work. We used the first three years of science data for which the associated IEM is suitable for

¹See LAT data selection recommendations at: http://fermi.gsfc.nasa.gov/ssc/data/analysis/documentation/Cicerone/Cicerone_Data_Exploration/Data_preparation.html.

measuring sources with extensions $> 2^\circ$ ². A detailed discussion of the instrument and event classes can be found in Atwood et al. (2009) and at the *Fermi* Science Support Center¹.

For each of the 279 SNRs we modeled emission within a 10° radius of the SNR’s center. As a compromise between number of photons collected, spatial resolution, and the impact of the IEM, we chose 1 GeV as our minimum energy threshold. The limited statistics in source class above 100 GeV motivated using this as our upper energy limit.

To avoid times during which transient sources near SNRs were flaring, we removed periods with significant weekly variability detected by the *Fermi* All-sky Variability Analysis (FAVA) (Ackermann et al. 2013a). We conservatively defined a radius within which a flaring source may significantly affect the flux of a source at the center. We take this distance to be the radio radius of an SNR plus 2.8° , corresponding to the overall 95% containment radius for the *Fermi*-LAT point spread function (PSF) for a 1 GeV photon at normal incidence (Ackermann et al. 2012c). The time ranges of FAVA flares within this distance were removed in 23 RoIs, leaving $\geq 98.9\%$ of the total data in each RoI.

²See the LAT caveats, http://fermi.gsfc.nasa.gov/ssc/data/analysis/LAT_caveats.html, particularly those for the IEM developed for Pass 7 reprocessed data described in http://fermi.gsfc.nasa.gov/ssc/data/access/lat/Model_details/FSSC_model_diffus_reprocessed_v12.pdf.

5.6 Input Source Model Construction

To characterize each candidate SNR we constructed a model of γ -ray emission in the RoI which includes all significant sources of emission as well as the residual background from CRs misclassified as γ -rays. We implemented an analysis method, built upon the `addSrcs` method described in 5.2, to create and optimize the 279 models for each of the 279 RoIs. For each RoI, we initially included all sources within the 10° RoI listed in the Second Fermi LAT source catalog (2FGL) (Nolan et al. 2012), based on 2 years of Source class data. To this we added pulsars from the Second Fermi LAT catalog of Gamma-ray Pulsars (2PC) (Abdo et al. 2013), based on 3 years of source class data, with 2PC taking precedence for sources that exist in both. For the diffuse emission we combined the standard IEM corresponding to our P7 data set, `gal_2yearp7v6_v0.fits`, with the standard model for isotropic emission, which accounts for extragalactic diffuse γ -ray emission and residual charged particles misclassified as γ -rays. Both the corresponding isotropic model, `iso_p7v6source.txt`, and the IEM are the same as used for the 2FGL catalog analysis³.

Compared to 2FGL, we used an additional year of data and limited the energy range to $1 - 100$ GeV. This can result in different detection significances and localizations than previously reported in 2FGL. To account for these effects, we recreated the RoIs' inner 3° radius regions, which encompass the radio extents of all known SNRs, observed to be $\leq 2.6^\circ$ and allows a margin for the LAT PSF. The weighted average 68% containment radius of the LAT PSF for events at 1 GeV is $\sim 0.7^\circ$ (?). We note that this implicitly assumes that an SNR's GeV extent should not be more than about an order of magnitude larger than its radio extension and also note that

³Further details on the diffuse emission models are available at <http://fermi.gsfc.nasa.gov/ssc/data/access/lat/BackgroundModels.html> [JAM: and in Chapter blah]

the selection biases stated in Green’s catalog limit the range of known SNRs’ radio extensions.

To build the inner 3° radius model of each RoI, we first removed all sources except identified active galactic nuclei (AGN) and pulsars, whose positions on the sky are independently confirmed by precise timing measurements (Abdo et al. 2013). Retained AGN were assigned their 2FGL positions and spectral model forms. Pulsars’ positions and spectral forms were taken from 2PC. 2FGL sources identified or associated with SNRs are removed when they lie within the inner 3° .

Using `addSrcs`, we generated a TS map via `pointlike` on a square grid with $0.1^\circ \times 0.1^\circ$ spacing that covers the entire RoI. At the position of the maximum TS value, we added a new point source with a Power Law (PL) spectral model:

$$\frac{dN}{dE} = N \frac{(-\Gamma + 1)E^{-\Gamma}}{E_{\max}^{-\Gamma+1} - E_{\min}^{-\Gamma+1}} \quad (5.4)$$

where N is the integrated photon flux, Γ is the photon index, and E_{\min} and E_{\max} are the lower and upper limit of the energy range in the fit, set to 1 GeV and 100 GeV, respectively. We then performed a maximum likelihood fit of the RoI to determine N and Γ and localized the newly added source. The significance of a point source with a PL spectral model is determined by the χ_n^2 distribution for n additional degrees of freedom for the additional point source, which is typically slightly less than $\sqrt{\text{TS}}$

To promote consistent convergence of the likelihood fit, we limited the number of free parameters in the model. For sources remaining after the removal step, described above, we freed the normalization parameters for the sources within 5° of the RoI center, including identified AGN and pulsars. For 2FGL sources between 5° and 10° , we fixed all parameters. The spectrum of the IEM was scaled with a PL whose normalization and index were free, as done in 2FGL. For the isotropic emission model, we left the normalization fixed to the global fit value since the RoIs are too

small to allow fitting the isotropic and Galactic IEM components independently. The isotropic component's contribution to the total flux is small compared to the IEM's at low Galactic latitudes.

After localizing them, the new sources were tested for spectral curvature. In each of the four energy bands between 1 and 100 GeV, centered at 1.8, 5.6, 17.8 and 56.2 GeV, we calculated the TS value for a PL with spectral index fixed to 2 and then summed the TS values. We refer to this as $\text{TS}_{\text{bandfits}}$. A value for $\text{TS}_{\text{bandfits}}$ much greater than the TS calculated with a PL (TS_{PL}) suggests that, with a more rapid calculation, that the PL model may not accurately describe the source. Analogously to 2FGL, we allow for deviations of source spectra from a PL form by modeling sources with a log-normal model known colloquially as LogParabola or logP:

$$\frac{dN}{dE} = N_0 \left(\frac{E}{E_b} \right)^{-(\alpha + \beta \log(E/E_b))} \quad (5.5)$$

where N_0 is the normalization in units of photons/MeV, α and β define the curved spectrum, and E_b is fixed to 2 GeV^4 . If $\text{TS}_{\text{bandfits}} - \text{TS}_{\text{PL}} \geq 25$, we replaced the PL spectral model with a logP model and refit the RoI, including a new localization step for the source. We retained the logP model for the source if the global log likelihood across the full band improved sufficiently: $\text{TS}_{\text{curve}} \equiv 2(\log \mathcal{L}_{\text{logP}} - \log \mathcal{L}_{\text{PL}}) \geq 16$. Otherwise we returned the source to the PL model which provided the better global log likelihood. Across all RoIs, less than 2% of the newly added sources retained the logP model.

We continued iteratively generating TS maps and adding sources within the entire RoI until additional new sources did not significantly change the global likelihood of the fit. The threshold criterion was defined as obtaining $\text{TS} < 16$ for three consecutively added new sources, denoted as $N_{\text{TS} < 16} = 3$. Despite iteratively adding

⁴Note: E_b is a scale parameter which should be set near the lower energy range of the spectrum being fit and is usually fixed, see Massaro et al. (2004)

a source at the location of the peak position in the TS map, the TS values of new sources may not decrease monotonically with iteration for several reasons. First, source positions were localized after fitting the RoI and generating the TS map. Second, some added sources were fit with a more complex spectral model than a simple PL. Finally, when creating the TS map, we fixed the source’s spectral index to 2, whereas when adding the actual source to the model, we allowed its index to vary.

The specific value of $N_{\text{TS} < 16} = 3$ was chosen to avoid missing sources with $\text{TS} \geq 25$ (the threshold commonly used for source detection in LAT data), and to optimize computation time. We tested the threshold by selecting eight representative SNRs from both complex and relatively simple regions of the sky, with both hard and soft spectral indices. The eight chosen regions were:

SNR043.3-00.2 (W49B): A relatively simple region and test case that was previously detected as a point-source SNR (Abdo et al. 2010b)

SNR034.7-00.4 (W44): Previous LAT studies showed the SNR had a GeV extension slightly larger than the radio size as well as surrounding GeV emission from nearby extended sources associated with molecular clouds (Abdo et al. 2010c; Uchiyama et al. 2012)

SNR078.2+02.1 (gamma Cygni): A complex region containing the SNR an embedded pulsar, several nearby pulsars, and a large diffuse structure known as the Cygnus cocoon which is believed to be a bubble of hot gas acting as a source of freshly accelerated CRs (Ackermann et al. 2011, 2012b). The region serves as a test of how robust `addSrcs` is in one of the more extreme RoIs. Despite the complexity of the region, Lande et al. (2012) detected GeV emission co-spatial with the radio SNR.

SNR027.4+00.0 (Kes 73), SNR031.9+00.0 (3C391), SNR292.2-00.5,

SNR332.4-00.4 (MSH 16-51), SNR205.5+00.5 (Monoceros): These five sources were found to have large fitted extensions (greater than twice the radio radius of the SNR) in preliminary SNRcat pipeline runs so were included to understand this occurrence.

We applied the procedure detailed above to the test RoIs using a criterion of $N_{\text{TS} < 16} = 6$ and counted how many $\text{TS} \geq 25$ sources would be excluded if a smaller $N_{\text{TS} < 16}$ criterion was used. Figure 5.2 shows how reducing the threshold to $N_{\text{TS} < 16} = 3$ cut only one significant source in any of the regions. A further criteria to validate the value of $N_{\text{TS} < 16}$ used in this paper was that the spectrum of a source of interest (i.e. the central extended SNR in an RoI) or extension was robust to the addition of nearby sources. In Figures 5.3, 5.4, and 5.5, we show an example of the evolution of the flux, index, and extension of SNR gamma Cygni as subsequent sources are added. Sources were added to the ROI until $\Delta(\log \mathcal{L}) < 8$, 6 times in a row, and see that while a significant source added close to the SNR can affect the fit of the extended source, these fits stabilize before our threshold is reached. Since the maximum number of sources added in any test RoI was 38, the minimum 14, and the total number of sources added across all test regions was 221, we chose to use $N_{\text{TS} < 16} = 3$ for the full sample of 279 RoIs.

To allow for proper convergence of the likelihood fit, we reduced the number of free parameters prior to each new source addition. If the previously added source was between 3° and 5° of the center of the RoI, just its normalization was freed, and if greater than 5° all its source parameters were fixed. To avoid having newly added sources overlap with pulsars, we deleted new sources from the RoI if they were within 0.2° of a γ -ray pulsar and refit the pulsar in the $1 - 100$ GeV range following the 2PC conventions. 2PC modeled pulsar spectra as a PL with an exponential




Figure 5.2: Histogram of the number TS ≥ 25 sources remaining in each of the 8 test RoI for iterations in which $\Delta(\log \mathcal{L}) < 8$ (i.e. TS < 16). Points are offset for each SNR for clarity. The text boxes detail statistics for the values of TS of significant sources for the 8 studied SNRs for each corresponding value on the x-axis.

cutoff (PLEC),

$$\frac{dN}{dE} = N_0 \left(\frac{E}{E_0} \right)^{-\Gamma} \exp \left(-\frac{E}{E_c} \right)^b, \quad (5.6)$$

where N_0 is the normalization factor, Γ is the photon spectral index, E_c the cutoff energy, and b determines to the sharpness of the cutoff. 2PC assessed the validity of fixing b to 1 in Equation 5.6 (PLEC1) by repeating the analysis using a PL model, as well as the more general exponentially cut off PL form, allowing the parameter b in Equation 5.6 to vary. For the pulsar spectra in this analysis, we compared the maximum likelihood values for spectral models with and without a cutoff and with and without the value of b being free, via $\text{TS}_{\text{cut}} \equiv 2(\log \mathcal{L}_{\text{PLEC1}} - \log \mathcal{L}_{\text{PL}})$ and




Figure 5.3: Flux evolution of the extended source coincident with SNR gamma Cygni (labeled ES 1 in the figure) as successive sources are added to the ROI. Dotted line is first time $\Delta(\log \mathcal{L}) < 8$, dashed line shows the final threshold for this test study.

$\text{TS}_b \equiv 2(\log \mathcal{L}_{\text{PLEC}} - \log \mathcal{L}_{\text{PLEC1}})$ to determine which to use. If $\text{TS}_{\text{cut}} < 9$ is reported for the pulsar in 2PC then a PL model is used. If $\text{TS}_{\text{cut}} \geq 9$, we then check to see if the cutoff energy fit in 2PC lies within the restricted energy range of 1 – 100 GeV used in this work. For pulsars with cutoffs ≥ 1 GeV, we then use the PLEC model if $\text{TS}_b \geq 9$, and the PLEC model with cutoff freed otherwise. For those pulsars with cutoffs less than 1 GeV the spectral parameters are fixed to the 2PC values.

To complete the construction of our point source RoI model, we took the output




Figure 5.4: Same as Figure 5.3 but for the PL spectral index.

of the previous steps and removed all sources with $TS < 16$. This final model was then used as the starting model for analyzing candidate SNR emission. In Figure 5.6, we show a residual TS map of the region around SNR W44 as an example of the source configuration in an RoI prior to running `addSrcs`. Figure 5.7 is a residual TS map of the same region after running `addSrcs` to decompose the region into point sources.

We conservatively allow sources with TS down to 16 ($\sim 4\sigma$) in order to account for the effects of at least the brightest sub-threshold sources on the parameter fits for the other sources in the model. Furthermore, while the SNR analysis method




Figure 5.5: Same as Figure 5.3 but for the radius of a uniform, symmetric disk.

described in the chapter 5.8 is allowed to remove sources, it cannot add them. Thus we start from a set of sources designed to allow the final model to capture all significant emission within the central region. To corroborate our method of systematically adding sources to a region, we compare our RoI source models with those found by the 2FGL approach in Chapter 5.7.




Figure 5.6: 1-100GeV residual TS map for SNRW44 before running `addSrcs` and with 2FGL sources removed from the inner 3° radius (yellow circle). Bin size is 0.2/pixel. Magenta circle shows a 5° radius. 2FGL and newly added sources are shown as green crosses.

5.7 Comparison of Source Models with 2FGL

This SNR catalog was constructed using 3 years of P7 Source class data in the energy range 1 – 100 GeV, whereas 2FGL used 2 years of data over the larger energy range 0.1 – 100 GeV. The differences in observing time and energy range resulted in residual, unmodeled emission in some RoIs as well as changes to some 2FGL sources’ spectral model, position localization, and detection significance. Here we




Figure 5.7: 1-100GeV residual TS map for SNR W44 after `addSrcs` has completed. 2FGL sources have been removed from the inner 3° radius (yellow circle), and the bin size is 0.2/pixel. Magenta circle shows a 5° radius. 2FGL sources are shown as green crosses.

compare the input source models constructed for this catalog, described in Chapter 5.6, with 2FGL to better understand the `addSrcs` method’s ability to describe the regions studied. Since we rederive the input source model only within a 3° radius of the center of each RoI, we consider sources only inside that radius.

Given the data set differences, in each RoI we expect similar but not identical numbers of sources relative to those in 2FGL. Figures 5.8 and 5.9 show the numbers

of significant ($TS \geq 25$) 2FGL sources and derived input model sources (excluding 2FGL identified AGN and pulsars kept in the input model) in individual RoIs as 2D histograms. In Figure 5.8, the number of sources in the derived input model is typically greater than the number of 2FGL sources that are significant at $1 - 100$ GeV. 73 of the 279 RoIs studied contain at least one of the 12 extended 2FGL sources. Since 2FGL extended sources were removed from the inner 3° of each RoI, and this region was repopulated with point sources, we can detect multiple point sources inside the extent of any removed extended 2FGL sources. This decomposition of extended sources, combined with the longer data set and different energy range compared to 2FGL, contribute to the high ratio of input model to 2FGL sources in some RoI, which demonstrates the need to rederive the source model.

To more accurately represent the 2FGL sources being reproduced in the central 3° , in Figure 5.9 we limited the input model sources to those within 0.2° (approximately the width of the core of the 10 GeV PSF) of a 2FGL source, effectively excluding input sources that are not co-spatial with a 2FGL source. Here we see that the majority of 2FGL sources have counterparts in the rederived set. As a region's complexity increases, seen as an increase in numbers of 2FGL sources, up to about half of the 2FGL sources may not have counterparts within 0.2° . Given that in these same regions we have more new sources than 2FGL sources, as seen in Figure 5.8, we find as expected that the longer data set with improved statistics at higher energies, where the angular resolution of the LAT is the best, allows us to add new sources to account for newly significant excesses in these complex regions. Additionally, sources with low TS in 2FGL are particularly susceptible to having a newly added source which may start at a similar position but then localize further than 0.2° from the 2FGL source.

Thus, we find that the method developed and used here produces a model which




Figure 5.8: Comparison of the number of 2FGL sources with $TS_{1-100\text{ GeV}} \geq 25$ (excluding AGN and pulsars) with the number of newly added input model sources in the present analysis, for sources within 3° of the center of each RoI. The color scale shows the number of RoIs with a particular combination of numbers of 2FGL sources and new sources. White corresponds to no RoI with that combination of source counts.

reproduces the 2FGL sources as expected, including differences that trend as anticipated given the longer data set and modified energy range, yielding better spatial resolution. The new method thus provides reasonable representations of the regions being modeled as input for the final analysis.




Figure 5.9: Same as Figure 5.8, including only input model sources lying within 0.2° of a 2FGL source.

5.8 Detection Method

For each SNR, we characterize the morphology and spectrum of any γ -ray emission that may be coincident with the radio position reported in Green’s catalog. This was achieved by testing multiple hypotheses for the spatial distribution of γ -ray emission: a point source and two different algorithms for an extended disk. The best fit was selected based on the global likelihoods of the fitted hypotheses and their numbers of degrees of freedom. The hypothesis with the best global likelihood was then evaluated using a classification algorithm described in Acero et al. (2016) to determine whether the radio SNR could be associated with the detected γ -ray emission.

Spatial coincidence is a necessary but not sufficient criterion to identify a γ -ray source with a known SNR. The detection of spatially extended γ -ray emission increases confidence in an identification, especially if GeV and radio sizes are similar, as has been observed on an individual basis for several extended SNRs (e.g. Lande et al. 2012). The LAT has sufficient spatial resolution to detect many Galactic SNRs as extended. Figure 5.10 shows the distribution of radio diameters from Green’s catalog. Vertical dashed lines show the minimum detectable extension for sources with flux and index typical of those observed in this catalog, based on simulations using the P7V6 IRFs (Lande et al. 2012). The minimum detectable extension depends not only on the source’s flux and spectrum, but also the flux of the background, which was estimated by scaling the average isotropic background level by factors of 10 and 100 to be comparable to the Galactic plane. As figure 5.10 illustrates, roughly one third of the known Galactic SNRs may be resolved by the LAT if they are sufficiently bright GeV sources.

In order to determine the best representation for each SNR, we analyzed each SNR-centered RoI using multiple hypotheses for the spatial and spectral form. We used `pointlike` (Kerr 2010) to compare PL and logP spectral forms, to compare point source versus extended source hypotheses, and to analyze the robustness of sources near the extended source.

For each hypothesis, we started with the input model described in Chapters 5.5 and 5.6. We removed sources falling within the SNR’s radio disk unless they had been identified as an AGN or pulsar, as described in Chapter 5.6. We then proceeded to evaluate the following point and extended source hypotheses. For the point source hypothesis, a point source with a PL index initialized to 2.5 was placed at the radio centroid of the SNR. The positions, spectral index, and spectral normalization of the point source were then fit. As for the initial input model described in Chapter 5.6,




Figure 5.10: Distribution of SNR radio diameters from Green’s catalog. The vertical dashed lines indicate the minimum detectable extension for a source with a photon flux of $10^{-8} \text{ ph cm}^{-2} \text{ s}^{-1}$ in the 1 – 100 GeV energy range and a PL index of -2.5 , from simulations of 2 years of data and the P7V6 IRFs (Lande et al. 2012). In that work, simulations using 10x and 100x the isotropic background level (thin-dotted and thick-dashed lines) are used to estimate a reasonable background range for sources in the Galactic plane. [JAM: idk what plots are ok or not to include from papers I am an author on. I didnt make this plot, but I made the first version of it that inspired this one. Should I specifically include a comment for these plots that state they’re from the snr cat vs ones not in the paper?]

we tested the source for spectral curvature. To test the extended source hypothesis, we employed two separate procedures. Both employed a uniform disk model initially placed at the center of the RoI with a radius equal to that observed in the radio. In the first procedure, called the “disk” hypothesis, we fit both the position and extension of the disk, as well as tested for spectral curvature. A second procedure, which results in a model we call the “neardisk” hypothesis, additionally examines

the significance of sources nearby the disk, removing those which are not considered independently significant and refitting the disk position and radius. This procedure is described in Chapter 5.8.1.

Having evaluated these hypotheses, we compared the global likelihood values of the final extended hypothesis and of the point source hypothesis to determine which model had the largest maximum likelihood. If the source is significant in the best hypothesis, the model parameters are reported as Tables 1 and 2 in Acero et al. (2016) [JAM: should I include abridged tables?]. If no hypothesis had a significant γ -ray source coincident with the radio SNR, we calculated the upper limit on the flux from a region consistent with the radio SNR, described in Chapter 5.8.2, and report the results in Table 3 in Acero et al. (2016). [JAM: should I include the tables here? only if they're short, or abridged. I should def include the dist table because I did that work]

5.8.1 Localization, Extension, and Spectral Curvature

To test our hypotheses, we combined the initial model of point sources (Chapter 5.6) and the Galactic and isotropic diffuse contributions (Chapter 5.5 and 5.6) with a test source at the center of each RoI. All sources that fell within the radio SNR radius other than previously identified AGN or pulsars were removed, as was done for the input source model (Chapter 5.6). We note that multiple point sources removed within a single radio SNR radius may represent substructure within the source itself. This process conservatively assigns the majority of the flux to a single source, rather than decomposing it. We optimized the position of the test source with `pointlike`, iteratively allowing other model parameters to vary. For all hypotheses, the normalizations of all sources within 5° of the radio SNR center were fit while all other spectral parameters were fixed. The parameters for sources outside 5° were

also fixed.

For the point source hypothesis, a point source was placed at the radio centroid of the SNR. For the disk hypothesis, a uniform disk with radius equal to the radio radius was placed at the center. In both hypotheses, the normalization, index, and position of the candidate source were fit. For the disk hypothesis, the extension was also fit. Previous analyses of a range of possible Galactic SNR sources with similar data sets (e.g. Lande et al. 2012) typically showed no differences in global likelihood significant enough to justify choosing a Gaussian over a uniform disk template or vice versa. In addition, there was typically little difference in spectral parameters for the two spatial forms. For simplicity and clarity, we thus test only the uniform disk hypothesis. We allowed the localization to wander up to 5° in the fits as a reasonable upper limit on what might later be associated with the SNR. This is roughly twice the radius of largest radio SNR.

We included an additional disk hypothesis in which we recalculated the significance of each nearby point source. Because neighboring sources can influence the best fit disk parameters, we iteratively evaluated the significance of the neighboring source by calculating $\text{TS}_{\text{nearby}}$, defined as twice the difference between the model’s log-likelihood ($\log \mathcal{L}$) with the nearby point source and the model without the source, as determined by `pointlike`. Starting from the fitted disk model, for each neighboring point source we refit the position, extension, normalization, and spectrum of the uniform disk after removing the source. A nearby source was considered to be significant and thus kept if $\text{TS}_{\text{nearby}} \geq 9$. Each point source was evaluated individually, starting with the closest point source and extending radially outward to all sources within 1° of the furthest edge of the SNR’s radio disk. The final result of this iterative process is called the “neardisk” hypothesis which, for cases where neighboring source(s) were removed, can have different best fit disk parameters. As

a final step we refit the region with **gtlike**, using the neardisk model.

We chose the best extended source hypothesis by comparing the final disk and neardisk **gtlike** $\log \mathcal{L}$ values. Since the neardisk hypothesis can have fewer degrees of freedom, we chose the final disk hypothesis only if $2 \times (\log \mathcal{L}_{\text{disk}} - \log \mathcal{L}_{\text{neardisk}}) \geq 9$. Otherwise, we used the neardisk model as the final extended source hypothesis, hereafter referred to as the “disk hypothesis”.

In some cases a point source could not be localized starting at the SNR center. If the **pointlike** localization failed to converge when starting at the SNR center, we placed the candidate at the position of the most significant source removed from within the radio SNR radius and followed the procedure outlined above. For 69 RoIs there was either no source removed within the radio SNR or localization failed. For 31 RoIs, the candidate found had a $\text{TS} < 1$ and was removed from the model so as not to cause instabilities in the minimization. If the disk hypotheses converged and the final candidate was significant ($\text{TS} \geq 25$) in both the localization and spectral fits, the best extended hypothesis was selected.

Prior to the final fit of the region, sources were tested for spectral curvature using $\text{TS}_{\text{bandfits}} - \text{TS}_{\text{PL}} \geq 25$. If this criterion was satisfied then we replaced the PL spectral model with a logP model and refit the RoI. The final spectral model was selected, as for the input model, by comparing the $\log \mathcal{L}$ values, in this case $\text{TS}_{\text{curve}} \geq 16$, as defined in Chapter 5.6. Seven sources were found to be significantly better fit by a logP spectrum. To obtain final spectral parameters, we performed a final fit using the standard likelihood analysis tool **gtlike**. The normalization and index parameters were constrained to lie within a physically reasonable range.

We determined the final RoI model by selecting the most likely hypothesis based on a comparison of the **gtlike** global $\log \mathcal{L}$ of the point source hypothesis with the most likely extended source hypothesis. An extended hypothesis was considered

significantly more likely if TS_{ext} was ≥ 16 , where TS_{ext} is defined as twice the difference between the $\log \mathcal{L}$ of the final model from the disk hypothesis and that of the point source hypothesis, $\text{TS}_{\text{ext}} = 2(\log \mathcal{L}_{\text{disk}} - \log \mathcal{L}_{\text{point}})$, as in Lande et al. (2012). Otherwise, if the point source itself had $\text{TS} > 25$, we chose the point source hypothesis. In cases in which the optimization for the position of the point source did not converge but an extended disk was detected, we calculated the global $\log \mathcal{L}$ of the region without any source and with a point source at the center of the extended source. We then use the latter value to calculate TS_{ext} reported in Table 1 in Acero et al. (2016). For these candidates, if the source was significantly extended in both cases, we select the extended hypothesis. If none of the criteria were met, the candidate was considered undetected and we calculated an upper limit on the flux. Both the upper limits and flux calculation are described in the following subsection.

5.8.2 Fluxes and Upper Limits

Fluxes in the $1 - 100 \text{ GeV}$ band are determined using the standard analysis tool `gtlike` by a final fit of the model chosen to have the overall maximum likelihood characterization of the morphology and spectrum of the candidate source from the analysis detailed in Chapter 5.8 and 5.8.1. For those RoIs where no significant source was detected, we computed Bayesian upper limits on the flux using the method in described in Helene (1983) excluding any overlapping sources in the model that have not been identified as AGN or pulsars, as described in Chapter 5.6. As a spatial model we used a uniform disk equal in position and radius to that reported in Green’s catalog. We assumed the spectral model to be a PL and report upper limits for indices of 2.0 and 2.5 at 95% and 99% confidence levels. The choice of indices was motivated by the distribution of PL indices for classified sources. The results are reported in Acero et al. (2016).

5.9 Catalog Results

We detected 102 candidates with a final source $TS \geq 25$ in the 279 SNR RoIs (see Chapter 5.8). Of the 102 detected candidates, 36 passed the association probability threshold (see Acero et al. (2016)). Of these, 30 SNRs ($\sim 11\%$ of the total) show significant emission for all alternative IEMs and are classified as likely GeV SNRs. An additional four were identified as sources which are not SNRs. Two other candidates were demoted to marginal due to their dependence on the IEM, as described in the next paragraph. Of the sources likely to be GeV SNRs, 17 show evidence for extension ($TS_{\text{ext}} > 16$). Only sources associated with SNRs G34.7–0.4 and G189.1+3.0 show evidence of significant spectral curvature in the $1 - 100$ GeV range and are fit with logP spectra. Of the classified candidates, four extended and 10 point SNRs are new and published here for the first time. Descriptions of the new extended (G24.7+0.6, G205.5+0.5, G296.5+10.0, and G326.3–1.8) SNRs is given in Acero et al. (2016).

For those 245 SNRs that are either not detected by this analysis or which fail to meet the most stringent threshold for classification as a detected SNR, upper limits assuming the radio disk morphology of Green’s catalog with PL indices of 2.0 and 2.5 are reported in Table 3 in Acero et al. (2016). For those candidates which fail to meet the most stringent threshold, we replaced the source with the radio disk. We do not calculate upper limits for the four sources which are identified as not SNRs. A FITS version of the catalog is available through the *Fermi* Science Support Center, as described in Acero et al. (2016)⁵. [JAM: not including the detailed description of the new SNRs]

[JAM: all the multiwavelength CR stuff and I wasn’t so involved in, but it’s kind

⁵http://fermi.gsfc.nasa.gov/ssc/data/access/lat/1st_SNR_catalog/

of the crux of the paper. I don't know how much of it, if any to include here.]

5.10 GeV SNRs in a Multiwavelength Context:

Discussion Summary

Here, we summarize some of the findings detailed in the SNRcat that are most pertinent to the work performed for this thesis. [JAM: do I need a statement like this?]

As discussed in Chapter 3.2, the same population of radio, synchrotron-emitting CR electrons active in the shell of an SNR are expected to also produced γ -rays through the IC process and non-thermal bremsstrahlung radiation. If indeed the GeV and radio emission are produced in a single zone, it is reasonable to assume that the radio and γ -ray morphologies will correlate. We find that the best GeV diameter is within errors of the radio diameter for most of the candidates classified as being associated with an SNR, as shown in Figure 5.11. The same, co-spatial, electron population producing the GeV and radio emission is also suggestive of a potential correlation between the radio and γ -ray flux. Figure 5.12 shows the 1 GHz synchrotron flux versus the 1 GeV γ -ray flux for all SNRs. Despite suggestions of correlation, Kendalls τ rank correlation tests suggested no significant correlation exists between the radio and GeV flux or luminosities (not shown here). Various factors, such as the lack of detailed nonthermal emission modeling, distance measurement errors, and use of oversimplified γ -ray spectral models can skew these results, obscuring any inherent correlation.

We test for one further relationship between the radio and GeV emission and the underlying particle populations through the measured radio and GeV spectral indices. The energy of synchrotron-emitting leptons traced by 1 GHz observations




Figure 5.11: Radio diameters of Green’s catalog SNRs plotted against the fitted GeV diameters for those candidates with significant extension. The solid line represents equal radio and GeV diameters. All cases of detected extension have diameters greater than 0.2° . The ticks denote the radio extension of GeV point-like candidates, colored in order of their characteristics (young or interacting) and by their classifications (well defined or marginal). The small ‘x’s bracketing the points show the minimum and maximum GeV extensions allowed such that the source remains classified or marginally classified given the radio position and extension and best fit GeV position. Open circles indicate extended SNRs. All SNRs that passed classification are shown as black unless also classified as young, nonthermal X-ray SNRs (blue) or as interacting with MCs (red). Candidates that did not pass classification but that still had both fractional overlaps > 0.1 are gray. Statistical error bars have caps; error bars without caps represent the systematic error. [JAM: took out the filled circles and outlined part, If they are also young or interacting, they are outlined in blue or red, respectively (No extended marginally classified candidates were also identified as young or interacting.)]

depends on the magnetic field. If radio and GeV emission trace the same underlying particle population, then, at energies below the maximum energy reached by




Figure 5.12: Comparison of γ -ray and radio spectral flux densities for all SNRs and candidates. For all SNRs that were not detected or which failed classification, grey triangles indicate upper limits at 99% confidence, computed assuming the radio location and extension. Symbols, colors, and error bars are as in Figure 5.11. In addition, filled circles indicate point-like sources, and if grey markers are also young or interacting, they are outlined in blue or red, respectively (No extended marginally classified candidates were also identified as young or interacting).

the accelerated particles, the photon indices of radio and γ -ray emission should be correlated. For π^0 decay and e^\pm bremsstrahlung, the GeV and radio photon indices (Γ and α respectively) are related as $\Gamma = 2\alpha + 1$. For IC scattering leptons, the GeV and radio photon indices follow $\Gamma = \alpha + 1$, or in the case in which high-energy leptons have been cooled via synchrotron or IC radiation, $\Gamma = \alpha + 3/2$ (Reynolds 2008). Figure 5.13 compares the deduced radio spectral index α with the 1–100 GeV photon index Γ .

Nearly all candidates have γ -ray photon indices that are softer than predicted




Figure 5.13: Comparison of radio spectral index, α , and GeV photon index, Γ . The expected correlations are plotted for π^0 decay or e^\pm bremsstrahlung (solid) and IC emission from an electron population that is freshly accelerated (dashed) or cooled by radiative processes (dotted). Emission via a combination of processes would fall between the lines (e.g. between the solid and dashed for a combination of π^0 decay and IC emission). Symbols, colors, and error bars are as in Figure 5.11; ticks along the right hand side show the 1 – 100 GeV photon indices of those SNRs without reported radio spectral indices.

given their radio spectra, regardless of the GeV emission mechanism. The three young SNRs in blue are most consistent with a single underlying particle population, and it has been suggested they emit via IC (dashed line) at GeV energies. SNRs emitting via a combination of mechanisms under these simple assumptions would have indices falling between the two index relations, that is, they would lie in the region spanned by the π^0 /bremsstrahlung (solid) and IC (dashed) lines.

The lack of an observed correlation between the indices as expected under these simple assumptions suggests that more detailed physical models are required for the majority of SNR candidates. The observed soft GeV spectra relative to the radio has several potential explanations. The underlying leptonic and hadronic populations

may have different PL indices. The emitting particle populations may not follow a PL but may instead have breaks or even differing spectral shapes. Finally, there may be different zones with different properties dominating the emission at different wavelengths.

In the SNRcat, we also compared the GeV and TeV properties of SNRs to test the second common assumption in SNR models: that momentum distributions of the emitting particle populations do not follow simple PLs but have curvature or breaks. Such changes in spectral slope could also cause breaks in the γ -ray spectra. As TeV emission may originate via the same processes as the *Fermi*-LAT-observed GeV emission (e.g. Funk et al. 2008; Tam et al. 2010; Tibolla 2009), we might expect to see such a change reflected in a spectrum combining *Fermi*-LAT data with observations from iacts such as the High Energy Stereoscopic System (H.E.S.S.), the Very Energetic Radiation Imaging Telescope Array System (VERITAS), and the Major Atmospheric Gamma-ray Imaging Cherenkov telescopes (MAGIC). The converse is also true, where detection predictions in the GeV based on simple PL extrapolation from the TeV have been borne out in GeV studies, e.g. identifications of H.E.S.S. sources from Tibolla (2009) in 2FGL (Nolan et al. 2012) and Ackermann et al. (2012a).

In Figure 5.14 we plot the PL index in the GeV versus TeV range for all SNRs observed with both *Fermi*-LAT and an IACT detections. Six of the ten SNR candidates have TeV indices that are softer than their GeV indices, while three have GeV and TeV indices that are consistent with each other, within statistical and systematic errors. The remaining interacting candidate has a somewhat softer index at GeV energies than at TeV. Such a hardening of the index from GeV to TeV suggests that another particle population may dominate at higher energies or that the emission mechanism may change between the GeV and TeV regimes. Such curvature in

the spectrum may also explain the lack of a simple correlation between GeV and radio PL indices, as described above in this section. We also note that Figure 5.14 shows a distinct separation between young and interacting SNRs, which are often older. This suggests an evolution in index with age, from harder when younger to softer when older.




Figure 5.14: GeV index compared to published index measurements from iacts. The line corresponds to equal index values. The predominance of SNRs below the line suggests spectral curvature, potentially reflecting a change in spectral slope of the underlying particle population(s') index or indices. The ticks represent the GeV candidates with indices in the range of those with a TeV counterpart but with no TeV measurements themselves, demonstrating the limitations of the data set. Symbols, colors, and error bars are as in Figure 5.11.

In Figure 5.15, we take SNR ages from the literature and plot the 1 – 100 GeV photon index versus age. For our uniform sample of all GeV SNR candidates, young

SNRs tend to have harder GeV photon indices than interacting SNRs, which are likely middle aged, though the scatter in age for the two classes is one to two orders of magnitude. The general trend of younger SNRs having harder indices may be due to the decrease of the maximum acceleration energy as SNRs age and their shock speeds slow down. This would also result in fewer particles being swept up by the shock front, given a constant density, suggesting a corresponding decrease in luminosity with age.




Figure 5.15: Age versus GeV spectral index. For those with ages in the literature, the young (blue) SNR candidates are separated in this phase space from the identified interacting candidates (red). The ticks on the right show indices for GeV candidates without well-established ages. Symbols, colors, and error bars are as in Figure 5.11.

It is important to account for the distances of the SNRs when comparing physical quantities such as luminosity. Table 5.1 records distance from the literature, including the most recent and/or most certain distance estimates adopted in this

work. Of the 279 SNRs studied, only 112 have published distance estimates. Most often these distances are determined from observed line-of-sight velocities using an assumed Galactic rotation curve. Furthermore, kinematic distance estimates have largely been done on an individual basis, and are not uniformly determined for all SNRs. We do not consider distances derived using the “ Σ -D relation” because SNRs show a wide range of physical diameters (D) for a given surface brightness (Σ), limiting the utility of such a relationship for determining the distances to individual SNRs (Green 2012).

Table 5.1. Distances to SNRs

Name	d [kpc]	Method	Reference(s)
G000.0+00.0	8.5	IAU value	Kerr & Lynden-Bell (1986)
G000.3+00.0	$8.5^{+3.0}_{-3.0}$	H I	Lang et al. (2010)
G000.9+00.1	$8.5^{+7.5}_{-1.5}$	PSR	Camilo et al. (2009b)
G001.0-00.1	8.5	Maser	Yusef-Zadeh et al. (1999)
G001.4-00.1	$8.5^{+5.6}_{-0.0}$	Maser	Yusef-Zadeh et al. (1999)
G004.5+06.8	$7.0^{+2.0}_{-0.6}$	H I	Reynoso & Goss (1999), Sankrit et al. (2005), Aharonian et al. (2008b)
G005.4-01.2	$4.75^{+0.45}_{-0.45}$	Maser	Hewitt & Yusef-Zadeh (2009)
G005.7-00.0	$8.4^{+5.3}_{-5.3}$	Maser	Hewitt & Yusef-Zadeh (2009)
G006.4-00.1	$1.9^{+0.4}_{-0.4}$	Maser, CO	Velázquez et al. (2002)
G008.7-00.1	4.5	Maser	Kassim & Weiler (1990)
G009.7-00.0	4.7	Maser	Hewitt & Yusef-Zadeh (2009)
G011.2-00.3	$5^{+21}_{-0.5}$	H I	Radhakrishnan et al. (1972), Becker et al. (1985), Green et al. (1988)
G012.8-00.0	$4.7^{+1.3}_{-1.1}$	PSR	Halpern et al. (2012)
G013.3-01.3	$3.3^{+1.8}_{-1.7}$	CO	Seward et al. (1995), Koralesky et al. (1998)
G015.1-01.6	$5.7^{+1.3}_{-3.5}$	NH	Boumis et al. (2008)
G015.4+00.1	$4.8^{+1.0}_{-1.0}$	CO	Castelletti et al. (2013)
G016.7+00.1	$10.0^{+3.7}_{-7.4}$	Maser, CO	Hewitt et al. (2008), Reynoso & Mangum (2000)
G016.8-01.1	$5.1^{+4.6}_{-1.8}$	H I	Sun et al. (2011)
G018.1-00.1	$5.58^{+0.24}_{-0.27}$	H I	Leahy et al. (2014)
G018.6-00.2	$4.6^{+0.6}_{-0.6}$	H I	Johanson & Kerton (2009)
G018.8+00.3	$12.0^{+3.0}_{-5.1}$	H I	Tian et al. (2007b)
G021.5-00.9	$4.7^{+0.4}_{-0.4}$	PSR	Camilo et al. (2006), Tian & Leahy (2008b)
G021.8-00.6	$5.35^{+0.15}_{-0.15}$	CO, PSR	Tian & Leahy (2008b), Zhou et al. (2009)
G023.3-00.3	$4.2^{+0.3}_{-0.3}$	H I, CO	Leahy & Tian (2008b), Tian et al. (2007c)
G027.4+00.0	$8.5^{+0.6}_{-1.0}$	H I	Tian & Leahy (2008a)
G028.6-00.1	$7.0^{+1.5}_{-1.0}$	H I, NH	Bamba et al. (2001)
G028.8+01.5	4.0	NH	Schwentker (1994), Misanovic et al. (2010)
G029.7-00.3	$7.8^{+2.8}_{-2.7}$	H I	Leahy & Tian (2008a)
G031.9+00.0	7.2	Maser	Frail et al. (1996)
G032.4+00.1	17	NH	Yamaguchi et al. (2004)
G032.8-00.1	$5.2^{+1.5}_{-0.4}$	Maser	Zhou & Chen (2011)
G033.6+00.1	$7.0^{+1.0}_{-0.5}$	H I	Giacani et al. (2009), Frail & Clifton (1989)
G034.7-00.4	3.0	Maser	Paron et al. (2009)
G035.6-00.4	$3.6^{+0.4}_{-0.4}$	H I	Zhu et al. (2013)
G039.2-00.3	$6.5^{+6.0}_{-0.3}$	CO	Hewitt et al. (2009a), Su et al. (2011)
G041.1-00.3	$10.3^{+2.5}_{-3.9}$	CO	Jiang et al. (2010)
G043.3-00.2	10^{+2}_{-2}	H I	Brogan & Troland (2001)
G049.2-00.7	$4.3^{+1.7}_{-0.0}$	Maser, H I	Koo & Moon (1997), Hewitt et al. (2009b), Tian & Leahy (2013)
G054.1+00.3	$7^{+2.0}_{-2.5}$	H I	Leahy et al. (2008)
G054.4-00.3	$3.0^{+0.8}_{-0.8}$	CO	Junkes et al. (1992), Caswell (1985)
G069.0+02.7	$1.5^{+0.6}_{-0.4}$	H I, PSR	Leahy & Ranasinghe (2012)
G073.9+00.9	$1.3^{+0.7}_{-0.8}$	NH	Lozinskaya et al. (1993)
G074.0-08.5	$0.58^{+0.06}_{-0.06}$	PM	Blair et al. (2009)
G074.9+01.2	$6.1^{+0.9}_{-0.9}$	H I	Kothes et al. (2003)
G076.9+01.0	$10.0^{+5.0}_{-4.0}$	NH	Arzoumanian et al. (2011)
G078.2+02.1	$2^{+2.0}_{-1.5}$	H I	Leahy et al. (2013), Ladouceur & Pineault (2008)
G089.0+04.7	$1.7^{+1.3}_{-1.0}$	CO	Byun et al. (2006)
G106.3+02.7	$0.8^{+1.2}_{-0.1}$	H I	Kothes et al. (2001)
G109.1-01.0	$3.2^{+0.2}_{-0.2}$	H I, CO	Kothes & Foster (2012)
G111.7-02.1	$3.4^{+0.3}_{-0.1}$	PM	Reed et al. (1995)
G114.3+00.3	$1.0^{+1.5}_{-0.3}$	H I	Yar-Uyaniker et al. (2004)
G116.5+01.1	1.6	H I	Yar-Uyaniker et al. (2004)
G116.9+00.2	$1.6^{+1.9}_{-0.0}$	H I	Yar-Uyaniker et al. (2004), Hailey & Craig (1994)
G119.5+10.2	$1.4^{+0.3}_{-0.3}$	H I	Pineault et al. (1993)
G120.1+01.4	$3.0^{+2.0}_{-0.6}$	H I	Tian & Leahy (2011), Hayato et al. (2010), Krause et al. (2008)
G127.1+00.5	$1.15^{+0.35}_{-0.25}$	H I	Pauls (1977), Xilouris et al. (1993), Leahy & Tian (2006)
G132.7+01.3	$2.2^{+0.2}_{-0.2}$	H I	Routledge et al. (1991)

To investigate the role of environment in the trends for the young and interacting SNRs, we examined the GeV luminosity versus radio diameter in Figure 5.16. The square of the physical diameter (D) can be regarded as a reasonable indicator for SNR age and environment (see 2.2), as its evolution during the Sedov-Taylor phase follows

$$D \propto n_0^{-1/5} E_{\text{SN}}^{1/5} t^{2/5} \quad (5.7)$$

where n_0 is the ambient density of the surrounding medium, E_{SN} is the supernova energy, and t is the age of the SNR (Sedov 1959; Taylor 1950). We can thus use the physical diameter as an age proxy: “effective age”. Any apparent correlation between the luminosity and D^2 may be due to their inherent dependence on distance (squared). As observed in earlier works, e.g. Thompson et al. (2012), Figure 5.16 shows that, for the detected candidates, interacting SNRs are generally more luminous for a given physical diameter than young SNRs, though there is large scatter. This suggests that SNRs at the same effective age may be more luminous because they have encountered denser gas (n_0). It should also be noted that there is an explicit correlation between the luminosity and physical diameter plotted in Figure 5.16 as both are proportional to distance (squared), which is only reliably measured for a subset of our sample. Observational biases, including that young, often smaller and fainter SNRs tend to be more difficult to detect in the radio as well as in γ -ray, may also affect the observed trends.

[JAM: didn’t add anything about maximal CR energy content of all SNRs assuming purely hadronic emission, maybe don’t need to]

Table 5.1 (cont'd)

Name	d [kpc]	Method	Reference(s)
G156.2+05.7	$1.1^{+1.9}_{-0.8}$	NH	Pfeffermann et al. (1991), Gerardy & Fesen (2007)
G160.9+02.6	$0.8^{+3.2}_{-0.4}$	H I	Leahy & Tian (2007), Leahy & Roger (1991)
G166.0+04.3	$4.5^{+1.5}_{-1.5}$	H I	Landecker et al. (1989)
G180.0-01.7	$1.3^{+0.22}_{-0.16}$	PSR	Sallmen & Welsh (2004), Ng et al. (2007), Chatterjee et al. (2009)
G184.6-05.8	$1.93^{+0.57}_{-0.43}$	PM	Trimble (1973)
G189.1+03.0	1.5	Maser	Hewitt et al. (2006)
G205.5+00.5	$1.5^{+0.1}_{-0.7}$	H I	Odegard (1986), Fesen et al. (1985), Xiao & Zhu (2012)
G260.4-03.4	$2.2^{+0.3}_{-0.2}$	H I	Dubner & Arnal (1988), Paron et al. (2008)
G263.9-03.3	$0.287^{+0.017}_{-0.021}$	PSR	Moriguchi et al. (2001), Caraveo et al. (2001), Dodson et al. (2003)
G266.2-01.2	$0.75^{+0.15}_{-0.25}$	PM	Katsuda et al. (2008)
G272.2-03.2	$4.0^{+1.0}_{-2.2}$	NH	Lopez et al. (2011)
G284.3-01.8	3	CO	Ruiz & May (1986)
G290.1-00.8	$7^{+4.0}_{-3.5}$	H I	Rosado et al. (1996a), Slane et al. (2002), Reynoso et al. (2006)
G291.0-00.1	$5^{+1}_{-1.5}$	NH	Harrus et al. (1998)
G292.0+01.8	$6.2^{+0.9}_{-0.9}$	H I, PSR	Gaensler & Wallace (2003)
G292.2-00.5	$8.4^{+0.4}_{-0.4}$	PSR	Caswell et al. (2004), Camilo et al. (2000)
G296.5+10.0	$2.1^{+1.8}_{-0.9}$	H I	Giacani et al. (2000)
G304.6+00.1	$9.7^{+4.3}_{-1.7}$	H I	Caswell et al. (1975)
G308.4-01.4	$9.8^{+0.0}_{-3.9}$	NH	Prinz & Becker (2012)
G309.2-00.6	$4.0^{+1.4}_{-2.0}$	NH	Rakowski et al. (2001)
G315.1+02.7	$1.7^{+3.7}_{-0.3}$	PM	Stupar et al. (2007)
G315.4-02.3	$2.5^{+0.3}_{-0.2}$	PM	Rosado et al. (1996a), Sollerman et al. (2003a)
G315.9-00.0	8^{+2}_{-2}	PSR	Camilo et al. (2009a)
G316.3-00.0	$7.2^{+22.8}_{-2.5}$	H I	Caswell et al. (1975)
G318.2+00.1	$4.0^{+5.4}_{-0.7}$	H I	Hofverberg et al. (2010)
G320.4-01.2	$5.2^{+1.4}_{-1.4}$	H I, NH	Gaensler et al. (1999)
G321.9-00.3	$6^{+4.0}_{-0.5}$	H I	Stewart et al. (1993)
G326.3-01.8	$4.1^{+0.7}_{-0.7}$	NH	Rosado et al. (1996a), Kassim et al. (1993)
G327.1-01.1	$6.5^{+6.5}_{-1.5}$	NH	Sun et al. (1999)
G327.4+00.4	4.3	H I	McClure-Griffiths et al. (2001)
G327.6+14.6	$2^{+0.2}_{-0.4}$	PM	Nikolić et al. (2013)
G328.4+00.2	$17.4^{+2.6}_{-5.4}$	H I	McClure-Griffiths et al. (2001)
G330.2+01.0	4.9	H I	McClure-Griffiths et al. (2001)
G332.4-00.4	3.3	H I, CO	Paron et al. (2006), Reynoso et al. (2004)
G332.4+00.1	$7.5^{+3.5}_{-4.2}$	NH	Vink (2004)
G335.2+00.1	1.8	CO	Eger et al. (2011)
G337.0-00.1	11.0	Maser	Frail et al. (1996)
G337.2+00.1	$14.0^{+16.0}_{-0.5}$	H I, NH	Combi et al. (2005), Combi et al. (2006)
G337.2-00.7	$5.8^{+3.8}_{-3.8}$	H I	Rakowski et al. (2006), Lopez et al. (2011)
G337.8-00.1	12.3	Maser	Frail et al. (1996)
G338.3-00.0	$10.0^{+3.0}_{-2.0}$	H I	Lemiere et al. (2009)
G343.0-06.0	$1.0^{+0.5}_{-0.5}$	H I, NH	Kim et al. (2010), Welsh et al. (2003), Walker & Zealey (2001)
G346.6-00.2	11.0	Maser	Frail et al. (1996)
G347.3-00.5	$1.0^{+0.3}_{-0.2}$	H I, CO	Moriguchi et al. (2005)
G348.5+00.1	$9^{+0.5}_{-2.7}$	H I	Tian & Leahy (2012)
G348.5-00.0	$6.3^{+7.4}_{-3.3}$	Maser	Tian & Leahy (2012)
G348.7+00.3	13.2	H I	Tian & Leahy (2012)
G349.7+00.2	$11.5^{+0.7}_{-0.7}$	Maser	Frail et al. (1996), Tian & Leahy (2014)
G350.1-00.3	$4.5^{+6.2}_{-0.5}$	H I	Gaensler et al. (2008)
G351.7+00.8	$13.2^{+0.5}_{-11.1}$	H I	Tian et al. (2007a)
G352.7-00.1	$7.5^{+0.9}_{-0.7}$	H I, CO	Giacani et al. (2009)
G353.6-00.7	$3.2^{+0.8}_{-0.8}$	H I, CO	Tian et al. (2008)
G357.7+00.3	6.9	Maser	Frail et al. (1996)
G357.7-00.1	12	Maser	Frail et al. (1996), Gaensler et al. (2003), Lazendic et al. (2004)
G359.1-00.5	4.6	Maser	Yusef-Zadeh et al. (2007), Hewitt et al. (2008)

Note. — Table of SNR distances drawn from the literature. The method for determining the distance is noted as: CO = line-of-sight velocity from molecular CO lines; H I = kinematic distance from H I absorption; NH = extinction estimate from optical or X-rays; Maser = kinematic distance from OH maser velocity; PM = Proper motions; PSR = association with pulsar. The d_{error} values indicate the range of uncertainties from the quoted distance values as assessed in the cited publications. The distance uncertainties are often asymmetric.




Figure 5.16: The 1 – 100 GeV luminosity is plotted against the square of the radio diameters in pc of those SNRs with known distances. Symbols, colors, and error bars are as in Figure 5.11.

5.11 Conclusions

In this Chapter, we have discussed the state of γ -ray observations of SNRs prior to the launch of *Fermi*, and the unique role that the LAT plays in identifying SNRs and exploring γ -ray production mechanisms therein. We presented the new automated source addition and analysis method, **addSrcs**, and its application to studying the population of SNRs emitting GeV γ -rays, published in Acero et al. (2016). With this first *Fermi*-LAT SNR Catalog we have systematically characterized GeV emission in regions containing known radio SNRs, creating new methods to address issues associated with these typically complex regions. These include methods for system-

atically adding sources to a region and better estimating the systematic error due to choice of interstellar emission model (discussed in detail in)Acero et al. (2016)). From this, we have determined characteristics of the GeV SNR population, down to our measurement limit, finding 30 classified and 14 marginal candidates with a false identification limit of <22%(Acero et al. 2016). This GeV data provide a crucial context for the detailed modeling of individual SNRs. In combination with multiwavelength measurements, the GeV data now challenge simple, previously sufficient SNR emission models. Within the limits of existing multiwavelength data, our observations generally support previous findings of changes in spectral slope at or near TeV energies and a softening and brightening in the GeV range with age and effective age, yet we see indications that new candidates and new multiwavelength data may provide evidence of exceptions to this trend.

[JAM: EGRET point source sensitivity is $\sim 1 \times 10^{-7} \text{ cm}^{-2} \text{ s}^{-1}$ <http://fermi.gsfc.nasa.gov/science/instruments/table1-1.html> get this number from some paper instead?]

[JAM: Another pointlike assumption to speed things up is that the PSF doesn't vary too much with event incidence angle in individual bins. To ensure this even more, events with a reconstructed angle $\geq 66.4^\circ$ ($\cos \theta = 0.4$) are removed (idk why this angle)]

[JAM: I did work for mock catalog, but it was really just running addsrcs centered on the mock positions]

Chapter 6

Extended Source Detection above 50 GeV: The 2FHL Catalog

[JAM: chage ROI to RoI]

[JAM: Application of addSrcs searching for extended (and really point too) sources in the sky. 2FHL results on all Galactic sources. Index histograms for entire Second Catalog of Hard Fermi-LAT Sources (2FHL)population showing harder index for Galactic sources.]

[JAM: we don't just detect extended though! The pipeline was built to try extended and if that's not likely, revert to point source. Simple check with Alberto's pipeline to show we didn't detect any glaring discrepancies. Quick check of Alberto's results for clusters of point source at $|b| < 10^\circ$.]

6.1 Introduction

[JAM: give this a different title] The LAT (Atwood et al. 2009) on board the *Fermi* γ -ray space telescope has been surveying the whole sky since August 2008. Its unprecedented sensitivity and localization accuracy allowed the detection of over 3,000 point-like sources in 4 years of data (see the third catalog of *Fermi*-LAT sources, 3FGL, Acero et al. 2015). Typically, *Fermi*-LAT catalog studies are based on source detection and characterization in the whole 0.1 GeV–100 GeV energy band. The larger photon statistics present at low energy, counterbalanced by the LAT point-spread function (PSF) whose size decreases with energy, yields an optimum sensitivity at few-GeV energies. The *Fermi*-LAT catalogs are thus representative of the GeV sky more than they are of the MeV or the sub-TeV sky.

The first *Fermi*-LAT catalog of hard sources, named 1FHL (Ackermann et al. 2013b), provided an unbiased census of the sky at energies from 10 GeV up to 500 GeV. All-sky surveys at γ -ray energies are instrumental for ground-based imaging atmospheric Cherenkov telescopes (IACTs) such as H.E.S.S., MAGIC, and VERITAS (Hinton 2004; Holder et al. 2009; Lorenz 2004, respectively) in order to find new sources because of their limited fields of view (FoV).

Recently, a new event-level analysis (known as Pass 8) has been developed by the *Fermi*-LAT collaboration (Atwood et al. 2013a,b). Pass 8 significantly improves the LAT’s background rejection, PSF, and effective area. All these enhancements lead to a significant increase of the LAT sensitivity and its effective energy range, from below 100 MeV to beyond a few hundred GeV (Atwood et al. 2013a,b). These improvements are particularly significant above 50 GeV, yielding an enhancement in the acceptance and PSF by a factor between 1.2 and 2. It is interesting to note that, above 50 GeV, both the PSF (governed mostly by the pitch of the tracker silicon

strips and the spacing of the tracker planes, see Chapter 4.2) and the effective area of the LAT are only weakly dependent on energy and that the LAT operates, due to the (almost complete) absence of background, in the photon-limited regime.

We use 80 months of Pass 8 data to produce a catalog of sources detected by the LAT at energies¹ between 50 GeV and 2 TeV. This constitutes the second catalog of hard LATsources, named 2FHL, which allows a thorough study of the properties of the whole sky in the sub-TeV domain. In this thesis, we present results published in Ackermann et al. (2016), exclusively focusing on the Galactic science analysis and results and eschew the extragalactic results to the published 2FHL paper.

6.2 Analysis

6.2.1 Data Selection

We use 80 months (from August 2008 to April 2015) of P8_SOURCE photons with reconstructed energy in the 50 GeV–2 TeV range. At these energies the LAT has an energy resolution of around 10–15 % (1σ). Photons detected at zenith angles larger than 105° were excised to limit the contamination from γ -rays generated by cosmic-ray interactions in the upper layers of the atmosphere. Moreover, data were filtered removing time periods when the instrument was not in sky-survey mode². This leaves approximately 61,000 photons detected all over the sky. The count map reported in Figure 6.1 shows that the LAT observes many point-like sources and large scale diffuse emission in the direction of our Galaxy, some of which appears coincident with the so-called *Fermi* bubbles (Ackermann et al. 2014; Su et al. 2010).

¹Note the different energy range with respect to the 1FHL.

²This was achieved using the expression ‘(DATA_QUAL>0)&&(LAT_CONFIG==1)’ in `gtmkttime`.




Figure 6.1: Adaptively smoothed count map in the 50 GeV–2 TeV band represented in Galactic coordinates and Hammer-Aitoff projection. The image has been smoothed with a Gaussian kernel whose size was varied to achieve a minimum signal-to-noise ratio under the kernel of 2. The color scale is logarithmic and the units are counts per $(0.1 \text{ deg})^2$.

6.2.2 Source Detection

The first step of the source detection stage comprises the identification of source seeds, which are locations of potential sources whose significance is later tested through a maximum likelihood (ML) analysis. The seed detection method, described further in Ackermann et al. (2016), includes all the point sources detected in the 1FHL catalog. We note that this seed list may include statistical fluctuations as well as real sources with a non-optimal position.

A full ML analysis is then performed in order to verify which, among the seeds, are the reliable sources. The analysis is performed in 154 RoIs, varying between 10° and 20° in radius, whose sizes and positions in the sky are optimized to cover all

the seeds, ensuring that no more than 45 seeds are contained in a single RoI. For each RoI, we build a sky model that includes all the potential sources in the region as well as the Galactic and isotropic diffuse emissions³. These models, which are defined only up to ~ 600 GeV and ~ 900 GeV respectively, where extrapolated up to 2 TeV. The RoI models include also the extended sources present in the region (see Chapter ??). The model is fit to the data via the unbinned ML algorithm provided within the *Fermi* Science Tools⁴ (version v9r34p3).

The spectrum of each source is modeled with a power law because none of the sources is expected to show statistically significant spectral curvature detectable by the LAT in this energy band. Indeed, this was the case for the sources in the 1FHL catalog (Ackermann et al. 2013b).

The fit is performed iteratively in order to ensure convergence and to produce an optimal solution. It proceeds as follows:

1. Complex ML fits require approximate knowledge of the starting values of the parameters. For this reason the first step aims to find those values by fitting each single source separately to determine approximate spectral parameters. Throughout the entire process, the parameters of the diffuse emission models are left free to vary. The significance of each source is evaluated using the test statistic $TS = 2(\ln \mathcal{L}_1 - \ln \mathcal{L}_0)$, where \mathcal{L}_0 and \mathcal{L}_1 are the likelihoods of the background (null hypothesis) and the hypothesis being tested (e.g. source plus background). At each step in the procedure, marginal sources, those with $TS < 10$, are removed from the model. Once the spectral parameters and significance of each source have been evaluated, a global fit for which all the

³We use the `gll_iem_v06.fits` and `iso_P8R2_SOURCE_V6_v06.txt` templates available at <http://fermi.gsfc.nasa.gov/ssc/data/access/lat/BackgroundModels.html>.

⁴Available at <http://fermi.gsfc.nasa.gov/ssc/data/analysis/software/>.

parameters of the sources with a $TS \geq 10$ are allowed to vary is performed. Then one more global fit is performed after removing all the sources that had $TS < 10$ at the previous global fit. This step, as well as all the others, includes sources that are spatially extended (see Chapter ??);

2. In this second step, the positions of point-like sources, using the best-fit sky model derived at step 1, are optimized using the `gtfindsrc` tool. This step is done iteratively as well by optimizing first the positions of the most significant sources found at step 1 and later those of the fainter ones;
3. The parameters and significances of sources are estimated again (as in step 1) using the best-fit source positions. This step produces the best-fit sky model for any given RoI. Seeds with $10 \leq TS < 25$ are included in the model, but not reported in the final catalog;
4. For each source we estimate the energy of the highest-energy photon (HEP) that the fit attributes robustly to the source model. This is done using the tool `gtsrcprob` and selecting the HEP that has a probability $> 85\%$ to belong to the source;
5. A spectrum with three logarithmically spaced bins (boundaries of 50 GeV, 171 GeV, 585 GeV, 2 TeV) is generated for each source in the RoI that is detected with $TS \geq 25$ and with the number of detected γ rays (estimated by the likelihood, N_{pred}) to be ≥ 3 .

The procedure described above achieves the detection of 360 sources (including the extended sources discussed next at Chapter ??) with $TS \geq 25$ and $N_{\text{pred}} \geq 3$ across the entire sky. The number of seeds kept in the ROI models with $10 \leq TS < 25$ is 453, while 7 are the seeds with $TS \geq 25$, but $N_{\text{pred}} < 3$. [JAM: took out the monte carlo sims reasoning for Npred]

6.3 Search for Spatially-Extended Sources

Preliminary runs of the source detection method described in Chapter 6.2.2 detected clusters of point sources in the Galactic plane, which were suggestive of spatially extended sources. It is also possible that clusters of seed sources, each with sub-detection-threshold significance, could be detected as a significant extended source. Not modeling extended γ -ray emission as such can lead to inaccurate measurements of spectral and spatial properties of both the extended source and neighboring point sources, particularly in the Galactic plane (Lande et al. 2012). Most of the TeV sources in the Galactic plane are spatially extended (Carrigan et al. 2013; Ong 2013), so to clearly connect LAT detections spectrally to these sources, extension detection and characterization is important. In the following, we distinguish between sources whose extension have been previously determined with *Fermi*-LAT and new extended sources that are reported for the first time in a LAT catalog. The details of all significantly detected extended sources are reported in Chapter 6.4.4.

6.3.1 Extended Sources Previously Detected by the LAT

We explicitly modeled sources as spatially extended when a previous, dedicated, analysis found the source to be resolved by the LAT. The 25 extended sources reported in 3FGL were included in our model using the spatial templates derived in the individual source studies (see references in Acero et al. 2015). Refitting the positions and extensions of the 3FGL extended sources in this energy range is beyond the scope of this work.

Of the 25 3FGL extended sources, 19 are significantly detected here above the detection threshold ($TS \geq 25$). Only 6 sources are not detected and, since all have $TS < 10$, are removed from the sky model (see Chapter 6.4.4 for details).

One extended LAT source has had a dedicated analysis published since the release of the 3FGL catalog. Abramowski et al. (2015a) reported joint H.E.S.S. and LAT observations of the very high energy (VHE) source HESS J1834-087. This source is coincident with supernova remnant (SNR) W41 and was detected as spatially extended in a wide energy range spanning 1.8 GeV to 30 TeV. In this paper, we employ the spatial model for the GeV emission determined in Abramowski et al. (2015a), leading to a significant detection of this source.

6.3.2 Newly Detected Extended Sources

In addition to modeling the extended sources mentioned in Chapter 6.3.1, we performed a blind search of the Galactic plane ($|b| < 10^\circ$) to identify potential extended sources not included in previously published works. Our analysis pipeline is similar to that used in Acero et al. (2016), with some modifications tailored to searching for multiple extended sources in an ROI. The pipeline employed the `pointlike` binned maximum likelihood package (Kerr 2010), in particular utilizing the extended source fitting tools validated by Lande et al. (2012) to simultaneously fit the position, extension, and spectra of sources in our ROI. We used the `addSrcs` method (developed for the SNRcat and described in Chapter 5.6) to characterize potentially extended sources across the Galactic plane. We detail how it was applied to the 2FHL catalog study below.

We created 72 ROIs of radius 10° , centered on $b = 0^\circ$ with neighboring ROIs overlapping and separated by 5° in Galactic longitude. Our initial model of the γ -ray emission in each ROI consisted solely of the Galactic diffuse (allowing just the normalization to be fit) and isotropic emission models (fixing the normalization), with no other sources in the ROI. Emission in the ROIs was further characterized by adding sources and fitting their spectral parameters (normalization and spectral

index) in a $14^\circ \times 14^\circ$ region.

A TS map, that included all significant sources found previously, made up of $0.1^\circ \times 0.1^\circ$ bins across the ROI, was created at each iteration and a small radius (0.1°) uniform disk, with a power-law spectrum was placed at the position of the peak TS pixel. The spectra of any newly added sources, as well as the position, extension, and spectral parameters of the disk were then fit. If $\text{TS}_{\text{ext}} \geq 16$, where $\text{TS}_{\text{ext}} = 2 \log(\mathcal{L}_{\text{ext}}/\mathcal{L}_{\text{ps}})$ (i.e. twice the log-likelihood ratio of an extended to a point source, Lande et al. 2012), then the disk was kept in the model. For $\text{TS}_{\text{ext}} < 16$, the extended source was replaced by a point source with a power-law spectral model. For the point-source replacement case, spectral parameters of sources in the ROI were fit and the position of the new point source was optimized. Finally, the spatial parameters of any previously added extended sources were refit iteratively before creating a new TS map and repeating the process. We stopped adding sources when the peak TS was less than 16 for two successive sources.

To assess the impact of fitting extended sources when starting with an ROI devoid of sources, a crosscheck analysis (also using `pointlike`) was performed across the Galactic plane. We included 3FGL point and extended sources, the Galactic diffuse and isotropic emission, and pulsars from the 2PC catalog (Abdo et al. 2013) (as well as from 3FGL) in the preliminary source model for each region. Sources were iteratively added to account for residual emission and both these residual sources and 3FGL sources were tested for extension. Remarkably, this alternative analysis converges (i.e. spectral and spatial parameters for the detected extended sources are compatible in both analyses) to the initially source-devoid analysis for nearly all detected extended sources.

Extended sources detected in the analysis described in this Chapter for which the position and extension were compatible with those found by the crosscheck were

included in the ROI model at step 1 of the full ML analysis detailed in Chapter 6.2.2. Seed point sources interior to the extended sources were removed prior to the ML fit. To address the ambiguity between detecting a source as spatially extended as opposed to a combination of point sources, we utilized the algorithm detailed in Lande et al. (2012) to simultaneously fit the spectra and positions of two nearby point sources.

Add more details about ts2pts We only consider a source to be extended if $TS_{\text{ext}} > TS_{2\text{pts}}$ (improvement when adding a second point source). Our blind search of the Galactic plane allowed us to find 5 sources not previously detected as extended by *Fermi*-LAT. Further details on these sources are presented in Chapter 6.4.4.

[JAM: say something more about comparing the point sources we detected with the ML analysis] [JAM: something quick also about testing the egal sky for nearest neighbors to see if there were any egal ES. do I have numbers or a plot for this somewhere?]

6.4 The 2FHL Catalog

The 2FHL catalog includes 360 sources detected over the whole sky, each with a likelihood test statistic of $TS \geq 25$ and number of associated photons, $N_{\text{pred}} \geq 3$. The source association procedure (detailed in Ackermann et al. (2016)) finds that 75% of the sources in the catalog (274 sources) are extragalactic⁵, 11% (38 sources) are of Galactic nature, and 13% (48 sources) are unassociated (or associated with a TeV source of unknown nature). The unassociated sources are divided between 23 sources located at $|b| < 10^\circ$, and 25 sources at $|b| \geq 10^\circ$. Therefore the fraction

⁵This includes N 157B, an extragalactic pulsar wind nebula (PWN).




Figure 6.2: Sky map, in Galactic coordinates and Hammer-Aitoff projection, showing the sources in the 2FHL catalog classified by their most likely association.

of extragalactic sources in the sample is likely larger than 80%. The number of 2FHL sources that have not been reported in 3FGL is 57, 47 of which have not been previously reported in any LAT catalog nor in the TeVCat⁶ catalog of TeV detected sources, and are thus new γ -ray sources. Figure 6.2 shows the location of 2FHL sources color-coded according to their source class. [JAM: don't include assoc table since I didn't do that work or include because it summarizes what we found?]

⁶<http://tevcat.uchicago.edu/>

6.4.1 General Characteristics of 2FHL Sources

The 2FHL sources have > 50 GeV fluxes ranging from $\sim 8 \times 10^{-12}$ ph cm $^{-2}$ s $^{-1}$ to $\sim 1.3 \times 10^{-9}$ ph cm $^{-2}$ s $^{-1}$ with a median flux of 2.0×10^{-11} ph cm $^{-2}$ s $^{-1}$ and a median spectral index of 2.83. The index uncertainty increases rapidly with the spectral index (e.g. the uncertainty is about ± 0.5 for sources with $\Gamma = 2$ whereas it is ± 2 for sources with $\Gamma = 5$). Half of the sources are localized to better than 1.7' radius at 68 % confidence. [JAM: don't include flux vs index stuff] The distributions of spectral indices and the highest photon energy reported in Figure 6.3 show that extragalactic sources tend to have larger photon indices (median of 3.13) than Galactic sources (median of 2.10). Because of the harder spectra, Galactic sources tend to have higher-energy HEPs than those of extragalactic sources as shown as well in Figure 6.3. It is interesting to note that unassociated sources have a median index of 2.22 (2.00 for sources at $|b| < 10^\circ$ and 2.96 for those at $|b| \geq 10^\circ$), showing that a fraction (see later) of unassociated sources is likely of Galactic origin.

[JAM: keep this?] Building a spectral energy distribution (SED) represents a powerful way to discriminate or infer the nature of a source. By combining the spectral data from the 3FGL, 1FHL, and 2FHL catalogs, it becomes possible to measure the SEDs of sources over four decades in energy. Although these catalogs rely on different exposures and most γ -ray sources are variable, these data allow us to characterize the high-energy peak of their broadband SEDs. The SEDs of a few notable sources will be shown in the next sections.

6.4.2 The 2FHL Galactic Source Population

The narrow PSF core (about 0.1°) and moderate Galactic diffuse emission (in comparison with the > 100 MeV band) allows the LAT to characterize and study well the emission of sources in the plane of our Galaxy above 50 GeV. Within $|b| < 10^\circ$, LAT has detected 103 sources. Of those, 38 sources are associated with Galactic sources, 42 with blazars, 14 are unassociated and 9 are associated with other γ -ray sources whose origin is not known (see below). Figure 6.4 shows cut-outs of the Galactic plane with all detected sources labeled.

Among the 38 Galactic sources, 16 are spatially coincident with SNRs, 13 are coincident with PWNe, 4 are associated with PWN/SNR complexes and the other 5 sources are X-ray binaries (3), one pulsar (PSR J0835–4510) and the Cygnus Cocoon. It is clear that the majority of Galactic sources detected above 50 GeV are associated with objects at the final stage of stellar evolution.

Galactic sources display on average hard spectra, which is a sign of efficient particle acceleration. Roughly 55% of all Galactic sources have a spectral index lower than 2.2. For comparison, only 14% of the 2FHL blazars display such hard spectra. A sizable fraction (approximately 25%, see Figure 6.3, upper panel) of Galactic sources has a photon index harder than 2, implying a high-energy SED peak in the TeV band. Indeed, as the lower panel of Figure 6.3 shows, LAT detects emission from many Galactic sources well beyond 500 GeV. All PWNe detected by *Fermi* are found to be powered by young and energetic pulsars (age $\lesssim 30$ kyr, Acero et al. 2013). While it is common for PWNe to show hard spectra, this is less so for SNRs whose majority (about 85%) display softer spectra (Hewitt et al. 2013). Hard-spectrum SNRs are typically young or mid-aged ($\lesssim 3$ – 5 kyr) and might be difficult to find in radio surveys. Thus, Galactic surveys at above 50 GeV have




Figure 6.3: Distribution of the spectral indices (*top panel*) and highest photon energy (*bottom panel*) of the Galactic sources (orange), extragalactic sources (green slash), and unassociated sources (brown dotted). The medians of the distributions are plotted with dashed, dash-dotted, and dotted vertical lines, respectively. Both plots show that a distinct population of hard-spectrum sources is of Galactic origin.




Fig. 6.4.— continued

the capability to detect new SNRs that might have been previously missed. Such an example is represented by the extended source 2FHL J0431.2+5553e which is spatially coincident with a new SNR (SNR G150.3+4.5) recently reported by Gao & Han (2014).

Of the 14 sources at $|b| < 10^\circ$ that do not have an association, 7 have power-law indices harder than 2 which renders them likely Galactic objects. It is interesting to note that 6 of these 7 objects are offset from the plane of the Galaxy by more than 4° . This is in marked contrast with the associated portion of the sample where only the Crab Nebula and the newly discovered SNR G150.3+4.5 (out of 34 SNR/PWN systems) have such a large offset. Thus it seems unlikely that all these unassociated sources are SNR/PWN systems.

6.4.3 Comparison with the H.E.S.S. Galactic Plane Survey

The H.E.S.S. array, with a field of view of about 5° and an angular resolution of approximately 0.12° , has invested 2800 hrs of exposure to survey part⁷ of the Galactic plane, reaching an average sensitivity of 2 % of the Crab Nebula flux (i.e. $4.5 \times 10^{-13} \text{ ph cm}^{-2} \text{ s}^{-1}$) at $\geq 1 \text{ TeV}$ (Aharonian et al. 2006b; Carrigan et al. 2013). Considering that the Crab Nebula spectrum is harder in the 2FHL band than in the $>1 \text{ TeV}$ band, we estimate that the average sensitivity of 2FHL in the same region of the H.E.S.S. survey is $\sim 3\text{--}4\%$ of the $50 \text{ GeV--}2 \text{ TeV}$ Crab Nebula flux. The slightly better sensitivity allows H.E.S.S. to detect 69 sources (as reported in the TeVCat), while the LAT finds 36 objects in the same area. However, the comparable sensitivities of the two surveys allow the study of the properties of the high-energy Galactic population. In the 2FHL catalog there is almost an equal number of SNRs and PWNe in contrast to what is found in the H.E.S.S. survey where the ratio of PWNe to SNRs is 1.5 to 1. This might be because the hardest PWNe and softest SNRs are difficult to detect respectively in the $>50 \text{ GeV}$ and $>1 \text{ TeV}$ bands.

Of the 36 2FHL sources that fall within the footprint of the H.E.S.S. survey, 23

⁷The H.E.S.S. Galactic plane survey extends between $283^\circ < l < 59^\circ$ and Galactic latitudes of $|b| < 3.5^\circ$.

have already been detected at TeV energies and are associated with known counterparts, while 7 are undetected. The remaining 6 objects (2FHL J1022.0–5750, 2FHL J1505.1–5808, 2FHL J1507.4–6213, 2FHL J1703.4–4145, 2FHL J1745.1–3035 and 2FHL J1856.8+0256) are spatially coincident with TeV sources whose origin is not known. All of them have hard spectral indices ($\Gamma < 2.2$), but it is interesting to note that 4 of them (2FHL J1022.0–5750, 2FHL J1505.1–5808, 2FHL J1703.4–4145, and 2FHL J1745.1–3035) have $\Gamma < 1.7$ (see also Figure 6.5).

We find that 2FHL J1022.0–5750 is spatially compatible with HESS J1023–575, an extended TeV source (Abramowski et al. 2011b), whose emission might be due to a PWN powered by PSR J1023–5746 (Acero et al. 2013). 2FHL J1505.1–5808 is spatially coincident with the unidentified object HESS J1503–582, which has a size of 0.26° and a flux above 1 TeV (Renaud et al. 2008) compatible with the extrapolation of the 2FHL J1505.1–5808 spectrum. Its spectrum, reminiscent of that of a PWN (e.g., HESS J1825–137, Grondin et al. 2011), is reported in Figure 6.5.

2FHL J1507.4–6213 is spatially coincident with HESS J1507–622, an extended source with a radius of 0.15° located 3.5° from the plane (Acero et al. 2011). The analysis of multiwavelength data showed that it is not possible to discriminate between a hadronic and leptonic origin of the emission, but that the latter scenario, if the emission is powered by a PWN, would require a pulsar generated in the explosion of a hyper-velocity star in order to reach the required distance from the plane (Domainko & Ohm 2012).

The sources 2FHL J1703.4–4145 and 2FHL J1745.1–3035 are the hardest sources ($\Gamma < 1.3$) among the six objects. 2FHL J1703.4–4145 is spatially coincident with the bright radio emission observed from the western side of the shell of SNR G344.7–001, a nearby mid-aged shell-type (age ~ 3000 yr and $8'$ diameter) SNR (Giacani et al. 2011). Both the 2FHL source and the SNR are spatially coincident with the larger,

elongated and unidentified HESS J1702–420 (Aharonian et al. 2008c). It thus seems likely that SNR G344.7–001 is the counterpart of 2FHL J1703.4–4145 and perhaps also of HESS J1702–420. The combined *Fermi*-H.E.S.S. spectrum of this source is reported in Figure 6.5.

2FHL J1745.1–3035 is found to be spatially coincident with the extended source HESS J1745–303, which may be comprised of up to three different sources (Aharonian et al. 2008a). Indeed, the position of 2FHL J1745.1–3035 is compatible with the ‘C’ emission region (the second brightest region in the complex, Aharonian et al. 2008a). However, the nature of this source is more complex, because the 2FHL source is marginally brighter at 1 TeV than the entire H.E.S.S. region and has also a harder spectrum (spectral index of 1.25 ± 0.38 in 2FHL versus 2.17 ± 0.11 as measured by H.E.S.S.).

Finally, 2FHL J1856.8+0256 is coincident with HESS J1857+026, an almost radially symmetric extended source (Aharonian et al. 2008d), whose emission likely originates from a PWN powered by PSR J1856+0245 (Rousseau et al. 2012).

6.4.4 Extended Source Results

In total, 31 sources are modeled as spatially extended and input into the ML analysis: 25 listed in 3FGL, 5 sources detected in the `pointlike` analysis (described in § 6.3.1) that were not detected as extended at the time of 3FGL, and one, SNR W41, reported recently by both the H.E.S.S. and LAT teams (Abramowski et al. 2015a). Names and properties of the extended sources are provided in Tables 6.1 and 6.2. Six extended sources, detected in 3FGL, were not detected in 2FHL: the SMC, S 147 (the point source 2FHL J0534.1+2753 was detected inside it), the lobes of Centaurus A (although we detect its core as a point source, 2FHL J1325.6–4301), W 44, HB 21 and the Cygnus Loop.




Figure 6.5: Spectral energy distributions of four Galactic sources constructed by combining data from the 3FGL (green diamonds), 1FHL (blue circles), and 2FHL (red stars). We show the 3FGL extended source SNR IC 443 (*top left*), the new 2FHL extended source PSR J1420–6048 (*top right*), and two “dark accelerators” detected by H.E.S.S. at TeV energies (Carrigan et al. 2013, purple squares) without a previous LAT counterpart: HESS J1503–582 (*bottom left*) and HESS J1702–420 (*bottom right*).

We detect a weak source, 2FHL J1714.1–4012 (TS = 27), just outside the southwestern edge of the 3FGL spatial template used to model the emission from SNR RX J1713.7–3946 (2FHL J1713.5–3945e). 2FHL J1714.1–4012 has a hard spectral index $\Gamma = 1.63 \pm 0.38$, that is within errors of the spectral index derived for the SNR, $\Gamma = 2.03 \pm 0.20$ (Abdo et al. 2011). It is unclear whether 2FHL J1714.1–4012 is a distinct source separated from the SNR, or the result of un-modeled residual emission due to an imperfection in the spatial template adopted for the extended

source.

2FHL J1836.5–0655e is associated with the PWN HESS J1837–069. The 3FGL catalog contains several point sources in the vicinity of the PWN. We detect three sources in the vicinity, 2FHL J1834.5–0701, 2FHL J1837.4–0717 and 2FHL J1839.5–0705, the first two of which are coincident with 3FGL sources (3FGL J1834.6–0659, 3FGL J1837.6–0717 respectively). The power-law spectral indices of the three 2FHL point sources and 2FHL J1836.5–0655e are all consistent with each other. The concentration of sources around HESS J1837–069 combined with the spectral compatibility of the sources is suggestive of a common origin to the γ -ray emission in this region. However, the surrounding γ rays could arise from other sources in the region (Gotthelf & Halpern 2008); further analysis is necessary to determine the nature of the sources in this region.

A brief description of the five new 2FHL extended sources is given below with residual TS maps for the region surrounding each source shown in Figure 6.6. Detailed analyses of these new extended sources will be reported in separate papers.

2FHL J1443.2–6221e overlaps with the young, radio-detected SNR RCW 86 (G315.42.3). RCW 86 is a 42' diameter SNR that lies at a distance of 2.3-2.8 kpc and is likely associated with the first recorded supernova, SN 185 AD (Rosado et al. 1996b; Sollerman et al. 2003b). With more than 40 months of data and using the P7SOURCE dataset, the LAT did not significantly detect the SNR, but upper limits on detection at GeV energies combined with detection of significant extension in the TeV (Aharonian et al. 2009) were sufficient to strongly favor a leptonic origin for the emission (Lemoine-Goumard et al. 2012).

An updated LAT analysis of RCW 86 using 76 months of data, as well as the Pass 8 event-level analysis, resulted in detection of the SNR by the LAT as well as significant extension measurement (Ajello et al. 2016; Hewitt & Fermi-LAT Collab-

oration 2015, the former published after Ackermann et al. (2016)). In this paper, we report the results derived for 2FHL J1443.2–6221e from the `pointlike` analysis described in § 6.3.1.

2FHL J1419.2–6048e is a newly detected extended sources with size $\sigma_{\text{disk}} = 0.36^\circ \pm 0.03^\circ$, that overlaps two nearby PWN/PSR complexes in the Kookaburra region. In the southwest of Kookaburra, HESS J1418–609 (Aharonian et al. 2006a) is coincident with both the extended non-thermal X-ray “Rabbit” PWN (G313.3+0.1, Roberts et al. 1999), and the γ -ray detected pulsar PSR J1418–6058 (Abdo et al. 2009). The northeast region, called “K3”, contains HESS J1420–607, coincident with PWN G313.5+0.3 and PSR J1420–6048. Acero et al. (2013) detected, with LAT, emission from both HESS J1418–609 (with a soft spectral index, pulsar-like spectrum) and HESS J1420–607 (with a hard power-law index) above 10 GeV, but only HESS J1420–607 was significantly detected above 30 GeV. Neither showed significant extension. Our result for the fitted power-law spectral index of 2FHL J1419.2–6048e is in agreement with the previous GeV and TeV results, yet our measured radius is considerably larger than the TeV extension. To compare the extensions of the uniform disk model used for 2FHL J1419.2–6048e in this paper to the Gaussian model of Aharonian et al. (2006a), we defined the radius which contains 68% of the source’s intensity as r_{68} , with $r_{68,\text{Gaussian}} = 1.51\sigma$, and $r_{68,\text{disk}} = 0.82\sigma$ (Lande et al. 2012). We find that $r_{68} \simeq 0.30^\circ$ for 2FHL J1419.2–6048e, and $r_{68} \simeq 0.09^\circ$ for HESS J1420–607. [\[JAM: Include details on the small kookaburra study if I have time\]](#)

2FHL J1355.2–6430e, coincident with the VHE source HESS J1356–645, is detected as extended ($\sigma_{\text{disk}} = 0.57^\circ \pm 0.02^\circ$) for the first time by the LAT in this work. The source HESS J1356–645 (Abramowski et al. 2011a) is associated with the pulsar PSR J1357–6429, which was determined to be powering a surround-

ing extended radio and X-ray PWN (Lemoine-Goumard et al. 2011). Acero et al. (2013) detected faint emission from the nebula, and derived a 99% confidence limit, Bayesian upper limit on extension ($\sigma_{\text{Gauss}} < 0.39^\circ$) in the absence of significant extension. The fitted spectral index for 2FHL J1355.2–6430e is compatible with the GeV and TeV results (Abramowski et al. 2011a; Acero et al. 2013), however, the fitted disk extension is larger than that of the TeV detection, with $r_{68} \simeq 0.47^\circ$ for 2FHL J1355.2–6430e and $r_{68} \simeq 0.30^\circ$ for HESS J1356–645.

2FHL J1112.4–6059e is an extended source ($\sigma_{\text{disk}} = 0.53^\circ \pm 0.03^\circ$) newly detected by the LAT that encircles two 3FGL sources, 3FGL J1111.9–6058 and 3FGL J1111.9–6038, and has another, 3FGL J1112.0–6135, just outside its boundary (Acero et al. 2015). The extended source also partially overlaps the massive star forming region NGC 3603.

Finally, **2FHL J0431.2+5553e** is a large extended source ($\sigma_{\text{disk}} = 1.27^\circ \pm 0.04^\circ$), with a hard spectrum, that has not been previously detected at γ -ray energies. It overlaps the recently discovered radio SNR G150.3+4.5 (Gao & Han 2014). G150.3+4.5 is a $2.5^\circ \times 3^\circ$ (Galactic coordinates) elliptical shell type SNR that has a steep radio synchrotron spectrum ($\alpha = -0.6$), indicative of radio SNRs. An in depth LAT analysis of this source extending then energy down to $E > 1$ GeV presented in Chapter 7

[JAM: where to add in things about comparing my point sources with alberto's]

[JAM: I also did this quick check looking to see if any extragal sources had close neighbors and might be extended.]

Table 6.1. 2FHL extended sources previously detected by the *Fermi*-LAT

2FHL Name	l [deg]	b [deg]	TS	Association	Class	Spatial model	Radius [deg]
J0526.6–6825e	278.843	-32.850	49.80	LMC	gal	2D Gaussian	1.87
J0617.2+2234e	189.048	3.033	398.64	IC 443	snr	2D Gaussian	0.27
J0822.6–4250e	260.317	-3.277	63.87	Puppis A	snr	Disk	0.37
J0833.1–4511e	263.333	-3.104	49.70	Vela X	pwn	Disk	0.91
J0852.8–4631e	266.491	-1.233	437.21	Vela Jr	snr	Disk	1.12
J1303.4–6312e	304.235	-0.358	56.06	HESS J1303–631	pwn	2D Gaussian	0.24
J1514.0–5915e	320.269	-1.276	165.51	MSH 15–52	pwn	Disk	0.25
J1615.3–5146e	331.659	-0.659	128.15	HESS J1614–518	spp	Disk	0.42
J1616.2–5054e	332.365	-0.131	87.18	HESS J1616–508	pwn	Disk	0.32
J1633.5–4746e	336.517	0.121	114.17	HESS J1632–478	pwn	Disk	0.35
J1713.5–3945e	347.336	-0.473	60.98	RX J1713.7–3946	snr	Map	0.56
J1801.3–2326e	6.527	-0.251	50.20	W 28	snr	Disk	0.39
J1805.6–2136e	8.606	-0.211	160.43	W 30	snr	Disk	0.37
J1824.5–1350e	17.569	-0.452	266.09	HESS J1825–137	pwn	2D Gaussian	0.75
J1834.9–0848e	23.216	-0.373	67.30	W 41	spp	2D Gaussian	0.23
J1836.5–0655e	25.081	0.136	62.72	HESS J1837–069	pwn	Disk	0.33
J1840.9–0532e	26.796	-0.198	163.15	HESS J1841–055	pwn	Elliptical 2D Gaussian	0.62, 0.38, 39
J1923.2+1408e	49.112	-0.466	44.60	W 51C	snr	Elliptical Disk	0.38, 0.26, 90
J2021.0+4031e	78.241	2.197	115.97	Gamma Cygni	snr	Disk	0.63
J2028.6+4110e	79.601	1.396	28.09	Cygnus Cocoon	sfr	2D Gaussian	3.0

Note. — List of the 20 extended sources in 2FHL that were previously detected as extended by the *Fermi*-LAT. All these sources are in 3FGL except W41, which is studied by Abramowski et al. (2015b). The Galactic coordinates l and b are given in degrees. The extension of the disk templates is given by the radius. The extension of the 2D Gaussian templates is given by the 1σ radius, and the elliptical templates are given by the semi-major axis, semi-minor axis, and position angle (East of North). Association, Class, and Spatial model are as given in Third Fermi LAT source catalog (3FGL).

Table 6.2. New 2FHL extended sources

2FHL Name	l [deg]	b [deg]	TS	TS_{ext}	TS_{2pts}	F_{50}	ΔF_{50}	Γ	$\Delta\Gamma$	Association	Class	Radius [deg]
J0431.2+5553e	150.384	5.216	87.9	83.4	26.2	11.70	2.11	1.66	0.20	G 150.3+4.5	snr	1.27 ± 0.04
J1112.4–6059e	291.222	-0.388	80.9	68.3	22.5	12.80	2.36	2.15	0.28	PSR J1112–6103	pwn	0.53 ± 0.03
J1355.2–6430e	309.730	-2.484	82.3	31.8	12.9	9.59	1.95	1.56	0.22	PSR J1357–6429	pwn	0.57 ± 0.02
J1419.2–6048e	313.432	0.260	109.3	49.1	15.6	17.60	2.80	1.87	0.19	PSR J1420–6048	pwn	0.36 ± 0.03
J1443.2–6221e	315.505	-2.239	75.6	29.9	19.2	7.23	1.70	2.07	0.30	SNR G315.4–2.3	snr	0.27 ± 0.03

Note. — List of the 5 new extended sources in 2FHL. All sources are characterized by a uniform disk template whose radius and uncertainty there in is given in the last column. l and b are Galactic coordinates. All coordinates are shown in degrees. TS is the test statistic. TS_{ext} is the significance of extension (6.3.2). TS_{2pts} is the TS of two simultaneously fit point sources (6.3.2). F_{50} and ΔF_{50} are the integrated photon flux between 50 GeV and 2 TeV and its uncertainty in units of 10^{-11} photon $\text{cm}^{-2} \text{s}^{-1}$. Γ and $\Delta\Gamma$ are the photon index and its uncertainty from a power-law fit. Association lists the primary overlapping source and Class the suspected source type. All uncertainties are 1σ uncertainties.

6.5 Summary

We have presented an all-sky analysis at ≥ 50 GeV of 80 months of LAT data relying on the new Pass 8 event-level analysis. Pass 8 delivers improvements in the acceptance and the PSF, reduces background of misclassified charged particles and extends the energy range at which the LAT is sensitive. All this allowed LAT to detect 360 sources in the 50 GeV–2 TeV range, performing an unbiased census of the >50 GeV sky for the first time. This catalog of sources (dubbed 2FHL) provides a bridge between the traditional 0.1–100 GeV band of LAT catalogs (Acero et al. 2015) and the $\gtrsim 100$ GeV band probed by IACTs from the ground. The 2FHL catalog has the potential to improve the efficiency with which new sources are detected at TeV energies since only about 25 % of the 2FHL sources were previously detected by IACTs.

The majority ($\gtrsim 80$ %) of sources detected in the 2FHL catalog are likely extragalactic because they are either located at high Galactic latitude or are associated with blazars. BL Lacs represent the largest population of sources detected by the LAT (54 %), followed by blazars of uncertain classification (16 %) and unassociated sources (13 %). Most BL Lacs in 2FHL belong to the HSP class and display the hardest (among blazars) γ -ray spectra and substantial emission in the 50 GeV–2 TeV band. These sources are powerful probes of the EBL and this work has shown that LAT has detected emission from many blazars at optical depths >1 .

The 2FHL includes 103 sources in the direction of the Galactic plane ($|b| < 10^\circ$). While a fraction of the sources (~ 39 %) is associated with blazars, the rest are Galactic and unassociated sources. Galactic sources generally display much harder photon indices than blazars (median of ~ 2 versus ~ 3) and copious TeV emission, both signs of efficient particle acceleration. Most Galactic sources are associated

with PWNe and SNRs, systems at the end of the stellar evolution cycle, and are detected as spatially extended. All the hard (spectral index < 2) unassociated sources within the plane of our Galaxy are likely of Galactic origin, since very few blazars have spectra as hard.

A comparison with a similarly long dataset of Pass 7 photons shows that Pass 8 allows the LAT to detect 35 % more sources. It is thus clear that Pass 8 and the accumulated exposure allow the LAT to extend its reach to higher energy and to open a new window on the sub-TeV sky. Sensitivity improves linearly with time in the photon-limited regime, thus further observations by the LAT in the coming years will probe the > 50 GeV sky even more deeply, providing important targets for current and future Cherenkov telescopes.

6.6 Scratch

Galactic seem to have harder indices than extra gal, suggests which unid'd in the plane are gal vs. egal. All extended harder than diffuse. Hints of different morphology between 2FHL and 2FGL, didn't study here, but this is one of the motivations for > 10 GeV study.




Figure 6.6: Residual TS maps for the five new extended sources described in Chapter 6.4.4. Only the Galactic diffuse and isotropic emission are included in the model to highlight the location of emission not associated with the diffuse background. Circles indicate the extents of the fit disks. The x marker in the bottom panel (2FHL J0431.2+5553e) shows the location of a point source in the ROI.

Chapter 7

SNR G150.3+4.5

Dedicated analysis of one interesting 2FHLresult. First blindly detected extended γ -ray source.

- Direct application of addSrcs to an interesting newly detected GeV SNR
- dedicated analysis of the one source to try to understand its nature
- It's interesting because :
- it was blindly detected
- might be the closest rem
- spectrum + index seems like dynamically young, but age and distance are hard to constrain, so it could be older
- Understanding this remnant contributes to the connection between Fermi and TeV telescopes

Take this all straight from the paper I write. Not sure how much more I'd need to add From the SNR cat, "We also note that one classified extended candidate may be consistent with an IC origin, though no error was reported on the radio spectral index measurement by ?". What sources is this? is G150 one of the only extended young SNRs? can't be true right? vela JR is young and extended

7.1 Introduction

Supernova remnants have long been thought to be the primary accelerators of cosmic rays up to the knee of the cosmic ray energy spectrum.

what to say about radio SNRs? Connect CRs to nonthermal emission and the LAT and Something about SNRs, cosmic ray accelerators, radio detections, connection between radio-LAT observations, G150 detection, 2FHL blind detection and SNRs at TeV (all young?), this paper extends the energy down to

Focus more on the hadronic vs leptonic since that's what's interesting? We describe the LAT and analysis results in §7.2, detail multiwavelength observations in §7.3, and discuss various emission origin scenarios in §7.4.

7.2 *Fermi*-LAT Observations and Analysis

7.2.1 Data Set and Reduction

Fermi-LAT is a pair conversion telescope sensitive to high energy γ -rays from 20 MeV to greater than 1 TeV (Ackermann et al. 2016), operating primarily in a sky-survey mode which views the entire sky every 3 hours. The LAT has wide field of view (~ 2.4 sr), a large effective area of ~ 8200 cm² above 1 GeV for on axis events and a 68% containment radius angular resolution of $\sim 0.8^\circ$ at 1 GeV. For further details on the instrument and its performance see Atwood et al. (2009) and Ackermann et al. (2012c).

In this analysis, we analyzed 7 years of Pass 8 data, from August 2nd 2008 to August 2nd 2015. The Pass 8 event reconstruction provides a significantly improved angular resolution [JAM: this is sadly unimportant unless I'm at higher energy or using the PSF types. The P8 total PSF at 1 GeV is about the same as for P7REP.

It's the acceptance/effective area that are considerably better at this energy], acceptance, and background event rejection (Atwood et al. 2013a,b), all of which lead to an increase in the effective energy range and sensitivity of the LAT. Source class events were analyzed within a $14^\circ \times 14^\circ$ region centered on SNR SNR G150.3+4.5 using the P8R2_SOURCE_V6 instrument response functions, with a pixel size of 0.1° . To reduce contamination from earth limb γ -rays, only events with zenith angle less than 100° were included.

For spectral and spatial analysis we utilized both the standard *Fermi*Science Tools (version 10-01-01)¹, and the binned maximum likelihood package **pointlike** (Kerr 2010). **pointlike** provides methods for simultaneously fitting the spectrum, position, and spatial extension of a source, and was extensively validated in Lande et al. (2012). Both packages fit a source model, the Galactic diffuse emission, and an isotropic component (which accounts for the background of misclassified charged particles and the extragalactic diffuse γ -ray background)² to the observations. In this analysis, we used the standard Galactic diffuse ring-hybrid model scaled for Pass 8 analysis, `gll_iem_v06.fits` (modulated by a power law function with free index and normalization), and for the isotropic emission, we used `iso_P8R2_SOURCE_V6_v06.txt`, extrapolated to 2 TeV as in Ackermann et al. (2016).

In our source model for the region, we included sources from the third *Fermi*-LAT catalog (Acero et al. 2015, 3FGL) within 15° of the center of our region of interest (RoI). We replaced the position and spectrum of any 3FGL pulsars in the region with their corresponding counterpart from the LAT 2nd pulsar catalog (Abdo et al. 2013). Residual emission unaccounted for by 3FGL sources is present in the RoI due to the increased time range and different energy selection with respect to that

¹<http://fermi.gsfc.nasa.gov/ssc/>

²<http://fermi.gsfc.nasa.gov/ssc/data/access/lat/BackgroundModels.html>

in 3FGL. We added to the RoI several point sources to account for this unmodeled emission and minimize the global residuals.[JAM: do I need to say more about these sources? should I mention adding them automatically and iteratively based on TS maps and reference SNRcat/2FHL? How close is the closest source? Mention this and use as an argument for not saying much more about them]. The normalization and spectral index of sources within 5° of the center of the RoI were free to vary, whereas all other source parameters were fixed. A preliminary maximum likelihood fit of the RoI was performed, and sources with a test statistic (TS) < 9 (TS is defined as, $TS = 2 \text{Log}(\mathcal{L}_1/\mathcal{L}_0)$ where \mathcal{L}_1 is the likelihood of source plus background and \mathcal{L}_0 that of just the background) were removed from the model.

7.2.2 Morphological Analysis

Studying the spatial extension of sources with the LAT is non-trivial due to the energy-dependent point spread function (PSF) and strong diffuse emission present in the Galactic plane. Soft spectrum point sources and uncertainties in the diffuse model can be a source of systematic error when not accurately modeling extended emission as such, particularly at low energies where the PSF is broad. To strike a balance between the best angular resolution and minimal source and diffuse contamination, we restrict our morphological analysis to energies between 1 GeV and 1 TeV. We divide this energy range into 12[JAM: 4bpd] logarithmically spaced bins for both `pointlike` and `gtlike` binned likelihood analyses.

Three unidentified 3FGL sources are located within the extent of SNR G150.3+4.5. 3FGL J0425.8+ 5600, located approximately 0.6° from the center of the SNR, is the closest of the three sources and is described with a power law spectrum of index $\Gamma = 2.35 \pm 0.17$ in the 3FGL catalog. The closest radio source to 3FGL J0425.8+5600 is NVSS J042719+560823, at 0.25 away (Ref?). 3FGL J0423.5+5442,

exhibits a power law spectral index, $\Gamma = 2.63 \pm 0.15$, with no clear multiwavelength source association. Finally, 3FGL J0426.7+5437 has a pulsar-like spectrum, yet in a timing survey performed with the 100-m Effelsberg radio telescope, Barr et al. (2013) were unable to detect pulsations from the source down to a limiting flux density of ~ 0.1 mJy. The source is located about 0.8° from the center of the SNR. We discuss this source and potential association with SNR G150.3+4.5 further in §7.4.2).

In our analysis, we removed 3FGL J0425.8+5600 and 3FGL J0423.5+544 from the RoI, but kept 3FGL J0426.7+5437 in the model since preliminary analyses showed clear positive residual emission at the position of the source if it was removed from the RoI. Figure 7.1 shows a residual TS map for the region around SNR G150.3+4.5. This point source detection-significance map was created by placing a point source modeled with a power law of photon index, $\Gamma = 2$ at each pixel and gives the significance of detecting a point source at each location above the background.

We modeled the excess emission in the direction of SNR G150.3+4.5 with a uniform intensity, radially-symmetric disk, simultaneously fitting the spatial and spectral components of the model via `pointlike`. The extension of the disk was initialized with a seed radius of $\sigma = 0.1^\circ$ and position centered on the radio position of SNR G150.3+4.5. We define the significance of extension as in Lande et al. (2012); $\text{TS}_{\text{ext}} = 2 \log(\mathcal{L}_{\text{ext}}/\mathcal{L}_{\text{ps}})$, with \mathcal{L}_{ext} being the likelihood of the model with the extended source and \mathcal{L}_{ps} that with of a point source located at the peak of emission interior to the extended source. For the disk model, $\text{TS}_{\text{ext}} = 298$, with a best fit radius, $\sigma = 1.40^\circ \pm 0.03^\circ$ [JAM: I should just put this all in a table and reference it], which is in excellent agreement with the radio size of the SNR determined in Gao & Han (2014). We tried adding back in to our model the two removed 3FGL




Figure 7.1: Background subtracted residual TS map above 1 GeV with $0.1^\circ \times 0.1^\circ$ pixels for fixed index $\Gamma = 2$, centered on SNR SNR G150.3+4.5. The orange circle and translucent shading show the fit disk radius and 1σ errors, respectively, for the extended source, the orange cross shows the position of 3FGL J0426.7+5437 (included in the background model), and blue dashed circle is the extent of the radio SNR.

sources but both were insignificant when fit on top of the best fit disk. Figure 7.2 is a Residual TS map of the same region as Figure 7.1, but with the disk source included in the background showing that the disk can account well for the emission in the region.

The morphology of the radio emission is suggestive of an elliptical or ring morphology, so an elliptical disk and ring spatial model were tested as well. For the ring model, the fit reduced to a disk with parameters matching those stated above. Using the elliptical model showed a weak improvement over the radially symmetric




Figure 7.2: Same as Figure 7.1 but with disk in the background model [JAM: should this be a residual counts map instead?]

model at the 2.6σ level ($\Delta\text{TS} = 9$ with two additional degrees of freedom), which we did not consider significant enough to say the GeV emission had an elliptical morphology (see Table ??). For the remainder of this study, we only considered the disk spatial model.[JAM: put Edisk in table too and reference i.] [JAM: I should double check this for 1GeV- 1TeV. I was done for 1-562 GeV, wait till addSrcs is done]

Other things we tried

fitting an extended source (starting with the 2FHL result) on top of the one currently there. Insignif. idk what to say about 2FHL yet.

another starting at the position of G149. Insignif

Say something about why we don't just go with the 3 3FGL sources. In the table I shouldn't just compare the 3 to the disk though because I also keep J0426 in the model. So the base comparison is really 2 sources vs the disk. Maybe it's enough to just say of course we keep the disk, we find one at GeV that matches really well with the radio. What did Josh's paper say about how modelling the spectrum of an intrinsically extended source as point sources skews the PS spectrum to softer energies?

He said, "Specifically, modeling a spatially extended source as point-like will systematically soften measured spectra", but idk if I get why. We see it with the 2 3FGL sources being softer than what the disk winds up being

Another thing to point out is how modeling as point vs extended, if it's really extended can affect the fit of other point sources nearby, like J0426, so I should show the spectrum of this source too? I fit both the norm and index of the source.

7.2.3 Spectral Analysis

After determining the best fit morphology with `pointlike` for the GeV emission coincident with SNR SNR G150.3+4.5, we used those results as a starting point for our `gtlike` maximum likelihood fit of the region to estimate the best spectral parameters for our model. The LAT data is well described by a power law across the entire energy range with a photon index, $\Gamma = 1.80 \pm 0.04$, and energy flux above 1 GeV of $(7.17 \pm 0.73 \times 10^{-11}) \text{ erg cm}^{-2} \text{ s}^{-1}$ and $\text{TS} = 373$ [JAM: these are the point-like results, change them when I get the gtlike res]. We tested the γ -ray spectrum of the extended disk for spectral curvature using a log-normal model (Log Parabola), and find no significant deviation from a power law ($\Delta\text{TS} \sim 1$).

Still to do

gtlike




Figure 7.3: Spectral energy distribution for the extended source coincident with SNR SNR G150.3+4.5. [JAM: replace with gtlike SED when I have it]

Systematics. Bracketing IRFs, alt iem, try varying the extension? still need to be done. Should probably just move this into the spectral section




Figure 7.4: Spectral energy distribution of 3FGL J0426.. [JAM: replace with gtlike SED when I have it]

7.3 Multiwavelength Observations and Analysis

7.3.1 HI

7.3.2 CO?

Do the CO maps add anything?

7.3.3 X-ray

No diffuse nonthermal X-ray emission observed by ROSAT. No point sources near the center? Should a pulsar even be near the center? How to quantify this? Can we place a limit on ambient density with an upper limit on thermal X-ray emission? Magnetic field with nonthermal?

7.4 Discussion and Results

7.4.1 What is it?

Size + HI suggest that near distance corresponding to different HI velocities suggest it's aged, spectrum looks more like young SNR (hard + no GeV break). Is it a weird young remnant or weird aged one? Leptonic dominated if young, hadronic dominated if older? Something about nearby dense clouds masking hadronic emission? Maybe this is only true for MeV cosmic rays that are screened out though and it would only mask the pion bump, but not this higher energy emission?

PWN or SNR. Can we rule out PWN? See W41 paper, MSH 11-61A, Fabios recent G326 work (no, he just tries to use the PSF types and testing different model templates to try to disentangle SNR from PWN)?

No PSR candidate near center (should it be near the center? Depends on age) Is there some limit we can place on the PWN based on not seeing the pulsar? Like on \dot{E} ? OR something like Mattana et al. 2009 correlation between $\text{flux}_x/\text{flux}_g \propto \dot{E}$?

Assume it's in Sedov phase based on size + near distance, and calculate age, upper limit on \dot{E} based on lack of x-ray flux? Or maybe if I assume the source is the PWN and GeV radius is PWN radius, then can I estimate \dot{E} based on size

and evolution inside SNR?

If we assume close distance, age is only $\approx 5\text{kyr}$, maybe this is a transitional SNR? What do others like this look like? Puppis A? Gamma Cygni is a similar age too.something

7.4.2 Distance Considerations

probably doesn't need to be a different section.

7.4.3 Nonthermal Modeling

I think I could get a working model with naima running pretty quickly, is it worth it?

7.5 Conclusions

[JAM: most of this should be the conclusions from the G150 paper]In this chapter, we have presented the publication on the dedicated analysis of the extended γ -ray emission detected in the direction of SNR G150.3+4.5. SNR G150.3+4.5 was first detected in radio by Gao & Han (2014), and subsequently detected in γ -rays in 2FHLabove 50 GeV. We discussed our LATmorphological analysis at energies $E \geq x$ GeV and spectral analysis down to $E \geq 750$ MeV, demonstrating a change in extension and centroid position compared to the 2FHLresult. Discuss potential source origin scenarios. Is it SNR or PWN? Is 2FHLsource same as $> \text{few GeV}$?

[JAM: for diss, not paper]. The way this figures into the whole is that this is a follow up analysis of one of the most interesting sources detected with addSrcs, and, (hopefully!), we're able to say something about the source and nature of the γ -ray emission, relation to SNR and 2FHLsource1

Chapter 8

SNR-MC, 10 GeV, and anything else?

Not sure how this is going to factor in yet. supernova remnant molecular cloud system (SNR-MC) should fit in somehow since a good deal of work was done? Less certain about 10gev.

Maybe this should say something about how the work I've done has contributed to the field of knowledge but also can lead to future work?

Chapter 9

Conclusions

Finally!

List of Symbols and Acronyms

2FGL Second Fermi LAT source catalog.

2FHL Second Catalog of Hard Fermi-LAT Sources.

2PC Second Fermi LAT catalog of Gamma-ray Pulsars.

AGN active galactic nuclei.

CGRO Compton Gamma-Ray Observatory.

CR cosmic ray.

EGRET Energetic Gamma-Ray Experiment Telescope.

FoV field of view.

GBM Gamma-ray Burst Monitor.

GRB gamma-ray burst.

IC inverse compton.

IEM interstellar emission model.

IRFs instrument response functions.

LAT Large Area Telescope.

LRT likelihood ratio test.

MC molecular cloud.

PL power law.

PSF point spread function.

PWN pulsar wind nebula.

RoI region of interest.

SED spectral energy distribution.

SNR supernova remnant.

SNR-MC supernova remnant molecular cloud system.

SNRcat First *Fermi*-LAT Supernova Remnant Catalog.

TS test statistic.

Bibliography

- Abdo, A. A. 2009, ArXiv:0902.1340, arXiv:0902.1340:0902.1340
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2009, Science, 325, 840
- . 2010a, ApJS, 188, 405
- . 2010b, ApJ, 722, 1303
- . 2010c, Science, 327, 1103
- . 2011, ApJ, 734, 28
- Abdo, A. A., Ajello, M., Allafort, A., et al. 2013, ApJS, 208, 17
- Abramowski, A., Acero, F., Aharonian, F., et al. 2011a, A&A, 533, A103
- . 2011b, A&A, 525, A46
- Abramowski, A., Aharonian, F., Ait Benkhali, F., et al. 2015a, A&A, 574, A27
- . 2015b, A&A, 574, A27
- Acero, F., Aharonian, F., Akhperjanian, A. G., et al. 2011, A&A, 525, A45
- Acero, F., Ackermann, M., Ajello, M., et al. 2013, ApJ, 773, 77
- . 2015, ApJS, 218, 23
- . 2016, ApJS, 224, 8
- Acharya, B. S., Actis, M., Aghajani, T., et al. 2013, Astroparticle Physics, 43, 3
- Ackermann, M., Ajello, M., Allafort, A., et al. 2011, Science, 334, 1103
- . 2012a, ApJ, 753, 83
- . 2012b, A&A, 538, A71

- Ackermann, M., Ajello, M., Albert, A., et al. 2012c, *ApJS*, 203, 4
- . 2013a, *ApJ*, 771, 57
- Ackermann, M., Ajello, M., Allafort, A., et al. 2013b, *ApJS*, 209, 34
- Ackermann, M., Albert, A., Atwood, W. B., et al. 2014, *ApJ*, 793, 64
- Ackermann, M., Ajello, M., Atwood, W. B., et al. 2016, *ApJS*, 222, 5
- Aharonian, F., Akhperjanian, A. G., Bazer-Bachi, A. R., et al. 2006a, *A&A*, 456, 245
- . 2006b, *ApJ*, 636, 777
- Aharonian, F., Akhperjanian, A. G., Barres de Almeida, U., et al. 2008a, *A&A*, 483, 509
- . 2008b, *A&A*, 488, 219
- . 2008c, *A&A*, 477, 353
- . 2008d, *A&A*, 477, 353
- Aharonian, F., Akhperjanian, A. G., de Almeida, U. B., et al. 2009, *ApJ*, 692, 1500
- Ajello, M., Baldini, L., Barbiellini, G., et al. 2016, *ApJ*, 819, 98
- Arzoumanian, Z., Gotthelf, E. V., Ransom, S. M., et al. 2011, *ApJ*, 739, 39
- Atwood, W., Albert, A., Baldini, L., et al. 2013a, *ArXiv:1303.3514*, *arXiv:1303.3514*
- Atwood, W. B., Abdo, A. A., Ackermann, M., et al. 2009, *ApJ*, 697, 1071
- Atwood, W. B., Baldini, L., Bregeon, J., et al. 2013b, *ApJ*, 774, 76
- Bamba, A., Ueno, M., Koyama, K., & Yamauchi, S. 2001, *PASJ*, 53, L21
- Barr, E. D., Guillemot, L., Champion, D. J., et al. 2013, *MNRAS*, 429, 1633
- Becker, R. H., Markert, T., & Donahue, M. 1985, *ApJ*, 296, 461
- Blair, W. P., Sankrit, R., Torres, S. I., Chayer, P., & Danforth, C. W. 2009, *ApJ*, 692, 335
- Boumis, P., Alikakos, J., Christopoulou, P. E., et al. 2008, *A&A*, 481, 705
- Brogan, C. L., Gelfand, J. D., Gaensler, B. M., Kassim, N. E., & Lazio, T. J. W.

- 2006, ApJ, 639, L25
- Brogan, C. L., & Troland, T. H. 2001, ApJ, 550, 799
- Byun, D.-Y., Koo, B.-C., Tatematsu, K., & Sunada, K. 2006, ApJ, 637, 283
- Camilo, F., Kaspi, V. M., Lyne, A. G., et al. 2000, ApJ, 541, 367
- Camilo, F., Ng, C.-Y., Gaensler, B. M., et al. 2009a, ApJ, 703, L55
- Camilo, F., Ransom, S. M., Gaensler, B. M., & Lorimer, D. R. 2009b, ApJ, 700, L34
- Camilo, F., Ransom, S. M., Gaensler, B. M., et al. 2006, ApJ, 637, 456
- Caraveo, P. A., De Luca, A., Mignani, R. P., & Bignami, G. F. 2001, ApJ, 561, 930
- Carrigan, S., Brun, F., Chaves, R. C. G., et al. 2013, ArXiv:1307.4868, arXiv:1307.4868
- Castelletti, G., Supan, L., Dubner, G., Joshi, B. C., & Surnis, M. P. 2013, A&A, 557, L15
- Caswell, J. L. 1985, AJ, 90, 1224
- Caswell, J. L., McClure-Griffiths, N. M., & Cheung, M. C. M. 2004, MNRAS, 352, 1405
- Caswell, J. L., Murray, J. D., Roger, R. S., Cole, D. J., & Cooke, D. J. 1975, A&A, 45, 239
- Chatterjee, S., Briske, W. F., Vlemmings, W. H. T., et al. 2009, ApJ, 698, 250
- Combi, J. A., Albacete Colombo, J. F., Romero, G. E., & Benaglia, P. 2006, ApJ, 653, L41
- Combi, J. A., Benaglia, P., Romero, G. E., & Sugizaki, M. 2005, A&A, 431, L9
- Dodson, R., Legge, D., Reynolds, J. E., & McCulloch, P. M. 2003, ApJ, 596, 1137
- Domainko, W., & Ohm, S. 2012, A&A, 545, A94
- Dubner, G. M., & Arnal, E. M. 1988, A&AS, 75, 363
- Eger, P., Rowell, G., Kawamura, A., et al. 2011, A&A, 526, A82

- Esposito, J. A., Hunter, S. D., Kanbach, G., & Sreekumar, P. 1996, *ApJ*, 461, 820
- Fesen, R. A., Blair, W. P., & Kirshner, R. P. 1985, *ApJ*, 292, 29
- Frail, D. A., & Clifton, T. R. 1989, *ApJ*, 336, 854
- Frail, D. A., Goss, W. M., Reynoso, E. M., et al. 1996, *AJ*, 111, 1651
- Funk, S., Reimer, O., Torres, D. F., & Hinton, J. A. 2008, *ApJ*, 679, 1299
- Gaensler, B. M., Brazier, K. T. S., Manchester, R. N., Johnston, S., & Green, A. J. 1999, *MNRAS*, 305, 724
- Gaensler, B. M., Fogel, J. K. J., Slane, P. O., et al. 2003, *ApJ*, 594, L35
- Gaensler, B. M., & Wallace, B. J. 2003, *ApJ*, 594, 326
- Gaensler, B. M., Tanna, A., Slane, P. O., et al. 2008, *ApJ*, 680, L37
- Gao, X. Y., & Han, J. L. 2014, *A&A*, 567, A59
- Gerardy, C. L., & Fesen, R. A. 2007, *MNRAS*, 376, 929
- Giacani, E., Smith, M. J. S., Dubner, G., & Loiseau, N. 2011, *A&A*, 531, A138
- Giacani, E., Smith, M. J. S., Dubner, G., et al. 2009, *A&A*, 507, 841
- Giacani, E. B., Dubner, G. M., Green, A. J., Goss, W. M., & Gaensler, B. M. 2000, *AJ*, 119, 281
- Górski, K. M., Hivon, E., Banday, A. J., et al. 2005, *ApJ*, 622, 759
- Gotthelf, E. V., & Halpern, J. P. 2008, *ApJ*, 681, 515
- Green, D. A. 2004, *Bulletin of the Astronomical Society of India*, 32, 335
- . 2009, *Bulletin of the Astronomical Society of India*, 37, 45
- Green, D. A. 2012, in *American Institute of Physics Conference Series*, Vol. 1505, American Institute of Physics Conference Series, ed. F. A. Aharonian, W. Hofmann, & F. M. Rieger, 5–12
- . 2014, *Bulletin of the Astronomical Society of India*, 42, 47
- Green, D. A., Gull, S. F., Tan, S. M., & Simon, A. J. B. 1988, *MNRAS*, 231, 735
- Grondin, M.-H., Funk, S., Lemoine-Goumard, M., et al. 2011, *ApJ*, 738, 42

- Hailey, C. J., & Craig, W. W. 1994, *ApJ*, 434, 635
- Halpern, J. P., Gotthelf, E. V., & Camilo, F. 2012, *ApJ*, 753, L14
- Harrus, I. M., Hughes, J. P., & Slane, P. O. 1998, *ApJ*, 499, 273
- Hartman, R. C., Bertsch, D. L., Bloom, S. D., et al. 1999, *ApJS*, 123, 79
- Hayato, A., Yamaguchi, H., Tamagawa, T., et al. 2010, *ApJ*, 725, 894
- Helene, O. 1983, *Nuclear Instruments and Methods in Physics Research*, 212, 319
- Hewitt, J. W., Acero, F., Brandt, T. J., et al. 2013, *ArXiv e-prints*:1307.6570, [arXiv:1307.6570](https://arxiv.org/abs/1307.6570)
- Hewitt, J. W., & Fermi-LAT Collaboration. 2015, in *American Astronomical Society Meeting Abstracts*, Vol. 225, *American Astronomical Society Meeting Abstracts*, 140.31
- Hewitt, J. W., & Lemoine-Goumard, M. 2015, *Comptes Rendus Physique*, 16, 674
- Hewitt, J. W., Rho, J., Andersen, M., & Reach, W. T. 2009a, *ApJ*, 694, 1266
- Hewitt, J. W., & Yusef-Zadeh, F. 2009, *ApJ*, 694, L16
- Hewitt, J. W., Yusef-Zadeh, F., & Wardle, M. 2008, *ApJ*, 683, 189
- . 2009b, *ApJ*, 706, L270
- Hewitt, J. W., Yusef-Zadeh, F., Wardle, M., Roberts, D. A., & Kassim, N. E. 2006, *ApJ*, 652, 1288
- Hinton, J. 2004, *New Astron.Rev.*, 48, 331
- Hofverberg, P., Chaves, R. C. G., Fiasson, A., et al. 2010, in *25th Texas Symposium on Relativistic Astrophysics*, 196
- Holder, J., Acciari, V., Aliu, E., et al. 2009, *AIP Conf.Proc.*, 1085, 657
- Inoue, T., Yamazaki, R., & Inutsuka, S.-i. 2010, *ApJ*, 723, L108
- James, F., & Roos, M. 1975, *Computer Physics Communications*, 10, 343
- Jiang, B., Chen, Y., Wang, J., et al. 2010, *ApJ*, 712, 1147
- Johanson, A. K., & Kerton, C. R. 2009, *AJ*, 138, 1615

- Junkes, N., Fuerst, E., & Reich, W. 1992, A&AS, 96, 1
- Kassim, N. E., Hertz, P., & Weiler, K. W. 1993, ApJ, 419, 733
- Kassim, N. E., & Weiler, K. W. 1990, Nature, 343, 146
- Katsuda, S., Tsunemi, H., & Mori, K. 2008, ApJ, 678, L35
- Kerr, F. J., & Lynden-Bell, D. 1986, MNRAS, 221, 1023
- Kerr, M. 2010, PhD thesis, University of Washington, arXiv:1101.6072
- Kim, I.-J., Min, K.-W., Seon, K.-I., Han, W., & Edelstein, J. 2010, ApJ, 709, 823
- Koo, B.-C., & Moon, D.-S. 1997, ApJ, 475, 194
- Koralesky, B., Frail, D. A., Goss, W. M., Claussen, M. J., & Green, A. J. 1998, AJ, 116, 1323
- Kothes, R., & Foster, T. 2012, ApJ, 746, L4
- Kothes, R., Reich, W., Foster, T., & Byun, D.-Y. 2003, ApJ, 588, 852
- Kothes, R., Uyaniker, B., & Pineault, S. 2001, ApJ, 560, 236
- Krause, O., Tanaka, M., Usuda, T., et al. 2008, Nature, 456, 617
- Ladouceur, Y., & Pineault, S. 2008, A&A, 490, 197
- Lande, J., Ackermann, M., Allafort, A., et al. 2012, ApJ, 756, 5
- Landecker, T. L., Pineault, S., Routledge, D., & Vaneldik, J. F. 1989, MNRAS, 237, 277
- Lang, C. C., Goss, W. M., Cyganowski, C., & Clubb, K. I. 2010, ApJS, 191, 275
- Lazendic, J. S., Wardle, M., Burton, M. G., et al. 2004, MNRAS, 354, 393
- Leahy, D., Green, K., & Tian, W. 2014, MNRAS, 438, 1813
- Leahy, D., & Tian, W. 2006, A&A, 451, 251
- Leahy, D. A., Green, K., & Ranasinghe, S. 2013, MNRAS, 436, 968
- Leahy, D. A., & Ranasinghe, S. 2012, MNRAS, 423, 718
- Leahy, D. A., & Roger, R. S. 1991, AJ, 101, 1033
- Leahy, D. A., Tian, W., & Wang, Q. D. 2008, AJ, 136, 1477

- Leahy, D. A., & Tian, W. W. 2007, *A&A*, 461, 1013
- . 2008a, *A&A*, 480, L25
- . 2008b, *AJ*, 135, 167
- Lemiere, A., Slane, P., Gaensler, B. M., & Murray, S. 2009, *ApJ*, 706, 1269
- Lemoine-Goumard, M., Renaud, M., Vink, J., et al. 2012, *A&A*, 545, A28
- Lemoine-Goumard, M., Zavlin, V. E., Grondin, M.-H., et al. 2011, *A&A*, 533, A102
- Lopez, L. A., Ramirez-Ruiz, E., Huppenkothen, D., Badenes, C., & Pooley, D. A. 2011, *ApJ*, 732, 114
- Lorenz, E. 2004, *New Astron.Rev.*, 48, 339
- Lozinskaya, T. A., Sitnik, T. G., & Pravdikova, V. V. 1993, *Astronomy Reports*, 37, 240
- Malkov, M. A., Diamond, P. H., & Sagdeev, R. Z. 2011, *Nature Communications*, 2, 194
- Massaro, E., Perri, M., Giommi, P., & Nesci, R. 2004, *A&A*, 413, 489
- Mattox, J. R., Bertsch, D. L., Chiang, J., et al. 1996, *ApJ*, 461, 396
- McClure-Griffiths, N. M., Green, A. J., Dickey, J. M., et al. 2001, *ApJ*, 551, 394
- Misanovic, Z., Kargaltsev, O., & Pavlov, G. G. 2010, *ApJ*, 725, 931
- Montmerle, T. 1979, *ApJ*, 231, 95
- Moriguchi, Y., Tamura, K., Tawara, Y., et al. 2005, *ApJ*, 631, 947
- Moriguchi, Y., Yamaguchi, N., Onishi, T., Mizuno, A., & Fukui, Y. 2001, *PASJ*, 53, 1025
- Ng, C.-Y., Romani, R. W., Briskin, W. F., Chatterjee, S., & Kramer, M. 2007, *ApJ*, 654, 487
- Nikolić, S., van de Ven, G., Heng, K., et al. 2013, *Science*, 340, 45
- Nolan, P. L., Abdo, A. A., Ackermann, M., et al. 2012, *ApJS*, 199, 31
- Odegard, N. 1986, *ApJ*, 301, 813

- Ohira, Y., Murase, K., & Yamazaki, R. 2011, MNRAS, 410, 1577
- Ong, R. A. 2013, ArXiv:1307.5003, arXiv:1307.5003
- Paron, S., Dubner, G., Reynoso, E., & Rubio, M. 2008, A&A, 480, 439
- Paron, S., Ortega, M. E., Rubio, M., & Dubner, G. 2009, A&A, 498, 445
- Paron, S. A., Reynoso, E. M., Purcell, C., Dubner, G. M., & Green, A. 2006, PASA, 23, 69
- Pauls, T. 1977, A&A, 59, L13
- Pfeffermann, E., Aschenbach, B., & Predehl, P. 1991, A&A, 246, L28
- Pineault, S., Landecker, T. L., Madore, B., & Gaumont-Guay, S. 1993, AJ, 105, 1060
- Prinz, T., & Becker, W. 2012, A&A, 544, A7
- Radhakrishnan, V., Goss, W. M., Murray, J. D., & Brooks, J. W. 1972, ApJS, 24, 49
- Rakowski, C. E., Badenes, C., Gaensler, B. M., et al. 2006, ApJ, 646, 982
- Rakowski, C. E., Hughes, J. P., & Slane, P. 2001, ApJ, 548, 258
- Reed, J. E., Hester, J. J., Fabian, A. C., & Winkler, P. F. 1995, ApJ, 440, 706
- Renaud, M., Goret, P., & Chaves, R. C. G. 2008, in American Institute of Physics Conference Series, Vol. 1085, American Institute of Physics Conference Series, ed. F. A. Aharonian, W. Hofmann, & F. Rieger, 281–284
- Reynolds, S. P. 2008, ARA&A, 46, 89
- Reynoso, E. M., & Goss, W. M. 1999, AJ, 118, 926
- Reynoso, E. M., Green, A. J., Johnston, S., et al. 2004, PASA, 21, 82
- Reynoso, E. M., Johnston, S., Green, A. J., & Koribalski, B. S. 2006, MNRAS, 369, 416
- Reynoso, E. M., & Mangum, J. G. 2000, ApJ, 545, 874
- Roberts, M. S. E., Romani, R. W., Johnston, S., & Green, A. J. 1999, ApJ, 515,

- Romero, G. E., Benaglia, P., & Torres, D. F. 1999, *A&A*, 348, 868
- Rosado, M., Ambrocio-Cruz, P., Le Coarer, E., & Marcelin, M. 1996a, *A&A*, 315, 243
- . 1996b, *A&A*, 315, 243
- Rousseau, R., Grondin, M.-H., Van Etten, A., et al. 2012, *A&A*, 544, A3
- Routledge, D., Dewdney, P. E., Landecker, T. L., & Vaneldik, J. F. 1991, *A&A*, 247, 529
- Ruiz, M. T., & May, J. 1986, *ApJ*, 309, 667
- Sallmen, S., & Welsh, B. Y. 2004, *A&A*, 426, 555
- Sankrit, R., Blair, W. P., Delaney, T., et al. 2005, *Advances in Space Research*, 35, 1027
- Schwentker, O. 1994, *A&A*, 286, L47
- Sedov, L. I. 1959, *Similarity and Dimensional Methods in Mechanics*
- Seward, F. D., Dame, T. M., Fesen, R. A., & Aschenbach, B. 1995, *ApJ*, 449, 681
- Slane, P., Smith, R. K., Hughes, J. P., & Petre, R. 2002, *ApJ*, 564, 284
- Sollerman, J., Ghavamian, P., Lundqvist, P., & Smith, R. C. 2003a, *A&A*, 407, 249
- . 2003b, *A&A*, 407, 249
- Stewart, R. T., Caswell, J. L., Haynes, R. F., & Nelson, G. J. 1993, *MNRAS*, 261, 593
- Stupar, M., Parker, Q. A., & Filipović, M. D. 2007, *MNRAS*, 374, 1441
- Sturmer, S. J., & Dermer, C. D. 1995, *A&A*, 293, astro-ph/9409047
- Su, M., Slatyer, T. R., & Finkbeiner, D. P. 2010, *ApJ*, 724, 1044
- Su, Y., Chen, Y., Yang, J., et al. 2011, *ApJ*, 727, 43
- Sun, M., Wang, Z.-r., & Chen, Y. 1999, *ApJ*, 511, 274
- Sun, X. H., Reich, P., Reich, W., et al. 2011, *A&A*, 536, A83

- Tam, P. H. T., Wagner, S. J., Tibolla, O., & Chaves, R. C. G. 2010, *A&A*, 518, A8
- Taylor, G. 1950, *Royal Society of London Proceedings Series A*, 201, 159
- Thompson, D. J., Baldini, L., & Uchiyama, Y. 2012, *Astroparticle Physics*, 39, 22
- Thompson, D. J., Bertsch, D. L., Fichtel, C. E., et al. 1993, *ApJS*, 86, 629
- Tian, W. W., Haverkorn, M., & Zhang, H. Y. 2007a, *MNRAS*, 378, 1283
- Tian, W. W., & Leahy, D. A. 2008a, *ApJ*, 677, 292
- . 2008b, *MNRAS*, 391, L54
- . 2011, *ApJ*, 729, L15
- . 2012, *MNRAS*, 421, 2593
- . 2013, *ApJ*, 769, L17
- . 2014, *ApJ*, 783, L2
- Tian, W. W., Leahy, D. A., Haverkorn, M., & Jiang, B. 2008, *ApJ*, 679, L85
- Tian, W. W., Leahy, D. A., & Wang, Q. D. 2007b, *A&A*, 474, 541
- Tian, W. W., Li, Z., Leahy, D. A., & Wang, Q. D. 2007c, *ApJ*, 657, L25
- Tibolla, O. 2009, in *American Institute of Physics Conference Series*, Vol. 1112, American Institute of Physics Conference Series, ed. D. Bastieri & R. Rando, 211–222
- Torres, D. F., Romero, G. E., Dame, T. M., Combi, J. A., & Butt, Y. M. 2003, *Phys. Rep.*, 382, 303
- Trimble, V. 1973, *PASP*, 85, 579
- Uchiyama, Y., Funk, S., Katagiri, H., et al. 2012, *ApJ*, 749, L35
- Velázquez, P. F., Dubner, G. M., Goss, W. M., & Green, A. J. 2002, *AJ*, 124, 2145
- Vink, J. 2004, *ApJ*, 604, 693
- Walker, A. J., & Zealey, W. J. 2001, *MNRAS*, 325, 287
- Welsh, B. Y., Sallmen, S., Jelinsky, S., & Lallement, R. 2003, *A&A*, 403, 605
- Xiao, L., & Zhu, M. 2012, *A&A*, 545, A86

- Xilouris, K. M., Papamastorakis, J., Paleologou, E. V., Andredakis, Y., & Haerendel, G. 1993, *A&A*, 270, 393
- Yamaguchi, H., Ueno, M., Koyama, K., Bamba, A., & Yamauchi, S. 2004, *PASJ*, 56, 1059
- Yar-Uyaniker, A., Uyaniker, B., & Kothes, R. 2004, *ApJ*, 616, 247
- Yusef-Zadeh, F., Goss, W. M., Roberts, D. A., Robinson, B., & Frail, D. A. 1999, *ApJ*, 527, 172
- Yusef-Zadeh, F., Arendt, R. G., Heinke, C. O., et al. 2007, in *IAU Symposium*, Vol. 242, *IAU Symposium*, ed. J. M. Chapman & W. A. Baan, 366–373
- Zhou, P., & Chen, Y. 2011, *ApJ*, 743, 4
- Zhou, X., Chen, Y., Su, Y., & Yang, J. 2009, *ApJ*, 691, 516
- Zhu, H., Tian, W. W., Torres, D. F., Pedalletti, G., & Su, H. Q. 2013, *ApJ*, 775, 95