


Kinematically similar basketball free throws have surprisingly different muscle contraction velocity profiles



¹Daniel A. Hagen, ¹Steven Caja, ²Suraj Chakravarthi Raja, ^{1,3}Francisco J. Valero-Cuevas



¹Department of Biomedical Engineering, ²Department of Electrical Engineering, ³Division of Biokinesiology & Physical Therapy, USC


Question

Is there a difference between a **good shot** and a **good looking shot** in basketball?


- Recent work re-emphasizes that neural control of limb movements is in fact **overdetermined**, with the rotation of a **few joints** determining the length changes in **many muscles** [1, 2].
- As Sherrington pointed out, if even one eccentrically contracting muscle fails to silence its stretch reflex appropriately, the movement is disrupted [3].
- Throws requiring **larger eccentric contractions** require **larger alpha-gamma control** and are therefore more prone to error, while **large concentric contractions reduce power output**.
- Therefore we investigated whether kinematically similar throws could exhibit large differences in eccentric and concentric muscle fiber contraction velocities.

Methods

We used an 18-muscle planar arm model to calculate a family of 100,000 random, feasible shoulder, elbow and wrist joint rotations that produced stereotyped basketball shots with different hand trajectories but identical starting and ending hand positions and velocities.




Utilizing a **posture specific moment arm matrix** it was possible to estimate fiber velocities for each of the 18 muscles from the time derivatives of the generated joint rotations (angular velocities) [1, 4, 5, 6].




Then for each throw, **muscle velocity profiles** were generated. The maximal eccentric and concentric contraction velocities were recorded for each muscle during the "upstroke" phase and the **Euclidean norms** of these values produced measures for **eccentric and concentric contraction levels**, respectively.

Results



Illustrated here are **three kinematically similar hand trajectories** that demonstrate **different levels of eccentric and concentric contractions** (top panel, bottom right), different configuration space trajectories (top panel, top right and bottom panel, bottom traces) as well as **different muscle fiber velocity profiles** (bottom panel, middle traces)



Discussion

- If there exist viable solutions to the motor task that exhibit different neuromuscular costs then this may help to explain the difference between a good shot and a good looking shot as a player searches the solution space.
- Alpha-gamma coactivation will severely limit kinematic redundancy as the time sensitive requirements of muscle afferentation will reduce the dimensionality of the feasible solution space of all desireable, smooth movement strategies.
- Additionally, these results may inform novel rehabilitation strategies that aim to treat neuromuscular disorders characterized by rigid movements (i.e. cerebral palsy, stroke, etc.) by avoiding these higher cost strategies.

References:

- [1] Valero-Cuevas, FJ. *Fundamentals of Neuromechanics*. Springer-Verlag London, 2016. [2] Valero-Cuevas, F, Cohn, B, Yingvason, H, & Lawrence, E. *J Biomech*, **48**(11), 2887–2896, 2015. [3] Sherrington, C.S. *Exp. Physiol.*, **6**(3) 252–310, 1913. [4] Winter, DA. *Biomechanics and motor control of human movement: Fourth edition*. Elsevier BV, 2013. [5] Ramsay, JW, Hunter, BV, & Gonzalez, RV. *J Biomech*, **42**(4), 463–473, 2009. [6] Holzbaur, KR, Murray, WM, & Delp, SL. *Annals of Biomedical Engineering*, **33**(6), 829–840, 2005.

(Additional references available upon request.)

Acknowledgements:

We thank the University of Southern California for facilities provided during the course BME/BKN 504, and Dr. Emily Lawrence who laid the foundation for this research and shared her prior work.

