
View Submission

CONTROL ID: 3604856

SUBMITTER (NAME ONLY): Brian Cohn

SUBMISSION TYPE: Scientific Oral or Poster

CURRENT CATEGORY: Neurogastroenterology/Functional/Motility

Abstract

TITLE: INTER-RATER VARIABILITY OF THE BRISTOL SCALE ON DIAPERED STOOL

AUTHORS (LAST NAME, FIRST NAME): Trieu, Melissa¹; Jamshidbaigi, Nikki¹; Punati, Jaya³; Danialifar, Tanaz³; Patel, Raza Ali²; Naon, Hillel³; Dallabrida, Susan^{1, 4}; Cohn, Brian Alexander¹

INSTITUTIONS (ALL): 1. Obvio Health USA, Inc, Orlando, FL, United States.

2. Department of Pediatrics, Intermountain Healthcare, Salt Lake City, UT, United States.

3. Division of Gastroenterology, Hepatology, and Nutrition, Children's Hospital of Los Angeles, Los Angeles, CA, United States.

4. SPRIM, Orlando, FL, United States.

ABSTRACT BODY:

Abstract Body : Background: The Bristol Stool Scale (BSS) is used in pediatric research trials as an outcome metric, and prior work has measured inter-rater reliability and intra-rater reliability across raters with various backgrounds—However, it is unclear whether the level of stool consistency of a given photo is more prone to inter-rater variability (SD).

Methods: 216 diapered stool photos from 9 children ages 2-4 years were collected. Per photo, we collected ratings from the careprovider (the untrained nonclinician), a central rater (a trained nonclinician) and 4 pediatric gastroenterologists. Group A (n=4 clinicians per photo, 215 photos) used the BSS ratings from the untrained nonclinicians as the reference rating, Group B references the trained nonclinician (n=4 clinicians, 212 photos), and Group C (n=3 clinicians, 214 photos) references 1 clinician (clinician “α”, a pediatric gastroenterologist with a focus on motility). Unreadable photos were removed.

Results: Certain BSS scores have higher inter-rater variability (ANOVA $F_A=9.867$, $p_A=1.74E-8$, $F_B=25.918$,

$p_B=7.66E-23$, $F_C=13.6$, $p_C=5.34E-13$, for A, B, and C respectively, which are shown as boxplots in Figure 1a-c, with Tukey HSD post hoc analyses in Table 1). In A, BSS-2 was the lowest score given by the untrained nonclinicians and also had the highest inter-rater variability ($std_{A2}=1.068$). For both B and C, BSS-1 had the highest inter-rater variability ($std_{B1}=1.703$, $std_{C1}=0.900$). For all 3 groups, BSS-7 had the lowest inter-rater variability ($std_{A7}=0.421$, $std_{B7}=0.163$, $std_{C7}=0.072$). Generally, stool ratings from BSS1-2 have the highest inter-rater variability, and BSS-7 has the lowest variability (Table 1, Figure 1a-c).

Conclusion: We provide evidence that inter-rater variability is significantly different across the Bristol stool scale when using a reference BSS rating from a) untrained nonclinicians, b) a trained nonclinician, and c) a pediatric gastroenterologist. This work motivates research into standardized training or automated scoring with a special emphasis on BSS 1-3, where we observe higher disagreement. The typical BSS transitions from constipation, normal and loose stools are 2-3 and 5-6, which had the same variability across A, B, and C (Table 1); although possibly useful in BSS assessments—this work may not hold clinical relevance for stool scales with fewer categories. This variance imbalance would represent a small difference in practice when stool is evaluated on a 3-point or 4-point scale. However, these measures of across-scale variability can provide valuable context when the full BSS1-7 is used.

TABLE:

PostHoc Tukey HSD Results

BSS-BSS comparison	A) Untrained nonclinician p-value	B) Trained Nonclinician p-value	C) Clinician p-value
1-2	NA	0.042	0.900
1-3	NA	0.001	0.662
1-4	NA	0.001	0.081
1-5	NA	0.001	0.327
1-6	NA	0.001	0.001
1-7	NA	0.001	0.001
2-3	0.768	0.632	0.900
2-4	0.900	0.449	0.232
2-5	0.583	0.690	0.670
2-6	0.764	0.188	0.003
2-7	0.004	0.001	0.001
3-4	0.900	0.900	0.744
3-5	0.900	0.900	0.900
3-6	0.900	0.900	0.025

3-7	0.001	0.001	0.001
4-5	0.868	0.900	0.900
4-6	0.900	0.900	0.229
4-7	0.001	0.001	0.001
5-6	0.900	0.865	0.227
5-7	0.091	0.001	0.001
6-7	0.001	0.001	0.131








Figure 1. Bristol Stool Scale (BSS) clinician inter-rater variability (SD), with respect to a given reference rating provided by **a)** untrained nonclinicians, **b)** a trained nonclinician, and **c)** clinician α . No untrained nonclinicians reported their child's stool as BSS-1.

© Clarivate Analytics | © ScholarOne, Inc., 2021. All Rights Reserved.

ScholarOne Abstracts and ScholarOne are registered trademarks of ScholarOne, Inc.

ScholarOne Abstracts Patents #7,257,767 and #7,263,655.

 [@ScholarOneNews](#) |  [System Requirements](#) |  [Privacy Statement](#) |  [Terms of Use](#)

Product version number 4.17.4 (Build 116). Build date Mon May 24 07:56:34 EDT 2021. Server ip-10-236-27-8