

# Mathematical Biostatistics Bootcamp: Lecture 10, T Confidence Intervals

Brian Caffo

Department of Biostatistics  
Johns Hopkins Bloomberg School of Public Health  
Johns Hopkins University

April 28, 2015



## Independent group $t$ confidence intervals

- Suppose that we want to compare the mean blood pressure between two groups in a randomized trial; those who received the treatment to those who received a placebo
- We cannot use the paired  $t$  test because the groups are independent and may have different sample sizes
- We now present methods for comparing independent groups

## Notation

- Let  $X_1, \dots, X_{n_x}$  be iid  $N(\mu_x, \sigma^2)$
- Let  $Y_1, \dots, Y_{n_y}$  be iid  $N(\mu_y, \sigma^2)$
- Let  $\bar{X}$ ,  $\bar{Y}$ ,  $S_x$ ,  $S_y$  be the means and standard deviations
- Using the fact that linear combinations of normals are again normal, we know that  $\bar{Y} - \bar{X}$  is also normal with mean  $\mu_y - \mu_x$  and variance  $\sigma^2(\frac{1}{n_x} + \frac{1}{n_y})$
- The pooled variance estimator

$$S_p^2 = \{(n_x - 1)S_x^2 + (n_y - 1)S_y^2\} / (n_x + n_y - 2)$$

is a good estimator of  $\sigma^2$

## Note

- The pooled estimator is a mixture of the group variances, placing greater weight on whichever has a larger sample size
- If the sample sizes are the same the pooled variance estimate is the average of the group variances
- The pooled estimator is unbiased

$$\begin{aligned} E[S_p^2] &= \frac{(n_x - 1)E[S_x^2] + (n_y - 1)E[S_y^2]}{n_x + n_y - 2} \\ &= \frac{(n_x - 1)\sigma^2 + (n_y - 1)\sigma^2}{n_x + n_y - 2} \end{aligned}$$

- The pooled variance estimate is independent of  $\bar{Y} - \bar{X}$  since  $S_x$  is independent of  $\bar{X}$  and  $S_y$  is independent of  $\bar{Y}$  and the groups are independent

## Result

- The sum of two independent Chi-squared random variables is Chi-squared with degrees of freedom equal to the sum of the degrees of freedom of the summands
- Therefore

$$\begin{aligned}(n_x + n_y - 2)S_p^2/\sigma^2 &= (n_x - 1)S_x^2/\sigma^2 + (n_y - 1)S_y^2/\sigma^2 \\&= \chi_{n_x-1}^2 + \chi_{n_y-1}^2 \\&= \chi_{n_x+n_y-2}^2\end{aligned}$$

## Putting this all together

- The statistic

$$\frac{\frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{\sigma \left( \frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}}}{\sqrt{\frac{(n_x + n_y - 2) S_p^2}{(n_x + n_y - 2) \sigma^2}}} = \frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{S_p \left( \frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}}$$

is a standard normal divided by the square root of an independent Chi-squared divided by its degrees of freedom

- Therefore this statistic follows Gosset's  $t$  distribution with  $n_x + n_y - 2$  degrees of freedom
- Notice the form is (estimator - true value) / SE

## Confidence interval

- Therefore a  $(1 - \alpha) \times 100\%$  confidence interval for  $\mu_y - \mu_x$  is

$$\bar{Y} - \bar{X} \pm t_{n_x+n_y-2, 1-\alpha/2} S_p \left( \frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}$$

- Remember this interval is assuming a constant variance across the two groups
- If there is some doubt, assume a different variance per group, which we will discuss later



## Likelihood method

- Exactly as before,

$$\frac{\bar{Y} - \bar{X}}{S_p \left( \frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}}$$

follows a non-central  $t$  distribution with non-centrality parameter  $\frac{\mu_y - \mu_x}{\sigma \left( \frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}}$

- Therefore, we can use this statistic to create a likelihood for  $(\mu_y - \mu_x)/\sigma$ , a standardized measure of the change in group means

## Example

Example from Rosner Fundamentals of Biostatistics, Page 304

- Comparing SBP for 8 oral contraceptive users versus 21 controls
- $\bar{X}_{OC} = 132.86$  mmHg with  $s_{OC} = 15.34$  mmHg
- $\bar{X}_C = 127.44$  mmHg with  $s_C = 18.23$  mmHg
- Pooled variance estimate

$$s_p^2 = \frac{7(15.34)^2 + 20(18.23)^2}{8 + 21 - 2} = 307.8$$


- $t_{27,.975} = 2.052$  (in R, `qt(.975, df = 27)`)
- Interval

$$132.86 - 127.44 \pm 2.052 \left\{ 307.8 \left( \frac{1}{8} + \frac{1}{21} \right) \right\}^{1/2} = [-9.52, 20.36]$$

## Likelihood plot for the effect size

Reasonable values for the effect size from the confidence interval

$$[-9.52, 20.36]/sp = [-.54, 1.16]$$



## Unequal variances

- Note that under unequal variances

$$\bar{Y} - \bar{X} \sim N\left(\mu_y - \mu_x, \frac{\sigma_x^2}{n_x} + \frac{\sigma_y^2}{n_y}\right)$$

- The statistic

$$\frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{\left(\frac{\sigma_x^2}{n_x} + \frac{\sigma_y^2}{n_y}\right)^{1/2}}$$

approximately follows Gosset's  $t$  distribution with degrees of freedom equal to

$$\frac{(S_x^2/n_x + S_y^2/n_y)^2}{\left(\frac{S_x^2}{n_x}\right)^2 / (n_x - 1) + \left(\frac{S_y^2}{n_y}\right)^2 / (n_y - 1)}$$

## Example

- Comparing SBP for 8 oral contraceptive users versus 21 controls
- $\bar{X}_{OC} = 132.86$  mmHg with  $s_{OC} = 15.34$  mmHg
- $\bar{X}_C = 127.44$  mmHg with  $s_C = 18.23$  mmHg
- $df = 15.04$ ,  $t_{15.04,.975} = 2.13$
- Interval

$$132.86 - 127.44 \pm 2.13 \left( \frac{15.34^2}{8} + \frac{18.23^2}{21} \right)^{1/2} = [-8.91, 19.75]$$