Table 2. Model comparison for proportions of carnivorous and omnivorous zooplankton along the Chl-a gradient. All models have the same terms of random effects (Longhurst Province intercept and Tow within Survey intercept and slope). The final model (indicated in bold) was selected based on the lowest BIC and highest marginal pseudo R^2.

Model	df residuals	Log-likelihood	BIC	Marginal Pseudo R^2	Conditional Pseudo R^2
Relative abundance of carnivorous zooplankton (RCZ)					
1. RCZ ~ Chl-a	138255	11292.2	-22489.6	0.0003	0.470
2. RCZ ~ Chl-a_sqrt	138255	11333.2	-22571.6	0.0018	0.472
3. RCZ ~ Chl-a_withCeiling	138255	11316.3	-22537.8	0.0011	0.471
4. RCZ ~ Chl-a_sqrt + Survey	138252	11513.5	-22896.9	0.2083	0.575
5. RCZ ~ Chl-a_sqrt + Latitude	138254	11336.3	-22566.1	0.0022	0.474
6. RCZ ~ Chl-a_sqrt + Survey + Latitude	138251	11518.3	-22894.5	0.2058	0.576
Relative abundance of omnivorous zooplankton (ROZ)					
1. ROZ ~ Chl-a	138255	340323.6	-680552.5	0.0002	0.449
2. ROZ ~ Chl-a_sqrt	138255	340319.0	-680543.2	0.0002	0.449
3. ROZ ~ Chl-a_withCeiling	138255	340326.9	-680559.1	0.0004	0.449
4. ROZ ~ Chl-a_sqrt + Survey	138252	340599.8	-681069.3	0.3171	0.680
5. ROZ ~ Chl-a_sqrt + Latitude	138254	340346.9	-680587.3	0.0045	0.466
6. ROZ ~ Chl-a_sqrt + Survey + Latitude	138251	340649.8	-681157.6	0.3055	0.683