Table S4. Summary statistics of surface chlorophyll-a and zooplankton trophic groups.

			SSP1-2.6	SSP2-4.5	SSP3-7.0	SSP5-8.5
Ocean climate variable	Mass concentration of phytoplankton expressed as chlorophyll in seawater (chlos kg/m^3)	Median	0.51843	0.51682	0.51347	0.39023
		Q1	0.44335	0.44380	0.45553	0.31936
		Q3	0.60039	0.59768	0.59416	0.47202
		Min	0	0	0	0
		Max	62.35185	70.35530	76.65367	4.18130
Proportion of zooplankton trophic groups	Omnivorous zooplankton	Median	0.56610	0.56610	0.56605	0.56410
		Q1	0.56494	0.56495	0.56513	0.56298
		Q3	0.56742	0.56738	0.56732	0.56539
		Min	0.55791	0.55791	0.55791	0.55791
		Max	0.94312	0.94695	0.94880	0.62237
	Carnivorous zooplankton	Median	0.49713	0.49719	0.49732	0.50195
		Q1	0.49403	0.49413	0.49427	0.49888
		Q3	0.49996	0.49994	0.49950	0.50459
		Min	1.03293e-06	1.645729e-07	3.877863e-08	0.34625
		Max	0.51628	0.51628	0.51628	0.51628
	Gelatinous filter-feeders	Median	0.14402	0.14410	0.14427	0.15059
		Q1	0.13993	0.14007	0.14024	0.14637
		Q3	0.14784	0.1478179	0.1472165	0.15430
		Min	2.11058e-13	6.014533e-15	3.65761e-16	0.03410
		Max	0.17187	0.17187	0.17187	0.17187