


Pathway and Network Analysis of Neoadjuvant Chemotherapy Nonresponders

Julia Gilley
BCB420

Presentation Roadmap



Background

What changes occur in breast cancer tumors that fail to respond to neoadjuvant chemotherapy (NAC)?

Experimental Design (Guevara-Nieto et al., 2025)

- Breast cancer patients
- Paired (before and after chemotherapy)
- Chemotherapy *nonresponders*

Dataset

- GEO accession GSE309004
- 27 patients, paired samples
- 19,736 genes x 54 samples


Normalization and Scoring

Normalization


- Authors already normalized
 - Median-of-ratios (MoR)
- I chose TMM normalization
 - Better for noisy data than MoR
 - Produced lower coefficient of variation

Scoring

- Quasi-likelihood generalized linear model
 - No info on true noise -> quasi-likelihood
- Significance filtering
 - logFC and FDR-corrected p-value
 - $FDR < 0.05$, and $|\log_2 \text{fold change}| > 1$



Presentation Roadmap



Thresholded ORA

- Used **clusterProfiler** (Yu et al., 2012; Wu et al., 2021) with GO Biological Process database (Ashburner et al., 2000; org.Hs.eg.db, Carlson 2024)
 - clusterProfiler: well-documented, uses BH correction
 - GO:BP: broad and granular

ORA thresholds: Adjusted p-value < 0.05, min gene set size = 10, max = 500 (Subramanian et al., 2005)

```
# ORA
ora_up <- enrichGO(gene = as.character(up_entrez),
                   OrgDb = org.Hs.eg.db, ont = "BP",
                   pAdjustMethod = "BH", universe = as.character(bg_entrez))
```

Non-Thresholded GSEA

Used **fgsea** (Korotkevich et al., 2021) with MSigDB Hallmark collection (v2023.2.Hs) gene sets


- Fgsea: computationally efficient
- Hallmark gene sets: very well annotated, non-redundant, and clean, easy to view

GSEA parameters: minSize = 15, maxSize = 500, nperm = 1000; significance threshold $FDR < 0.25$ (Subramanian et al., 2005)


```
# GSEA ranking metric
results_table$rank_metric <- -log10(results_table$PValue) * sign(results_table$logFC)

# fgsea
fgsea_results <- fgsea(pathways = hallmark_list, stats = ranked_genes,
                      minSize = 15, maxSize = 500, nperm = 1000)
```

Presentation Roadmap



Enrichment Results



ORA (upregulated genes): Stress response and AP-1/MAPK signaling

- Top genes: FOS, JUN, FOSB, ATF3, NR4A1

ORA (downregulated genes): Cell adhesion and tissue organization

- Top genes: CDH18, KISS1, SPANXA2

GSEA: 38 significantly enriched pathways, **all in FollowUp**


Enrichment Map

Enrichment map network from GSEA results

- **38 nodes** (pathways) and **23 edges** (all nodes enriched in the same direction (post-NAC))
- Edge similarity: Jaccard+Overlap
Combined, cutoff = 0.1

AutoAnnotate

- **6 clusters** and **12 singletons**



Major Themes

TNFA Signaling via NF- κ B: Major inflammatory pathway

Epithelial-Mesenchymal Transition (EMT): Process by which tumour cells lose cell adhesion


MYC Targets V2: MYC is a transcription factor that drives cell proliferation

Fatty Acid Metabolism: *Reprogramming of how cells can generate energy*

Coagulation/Complement: Immune activation and recruitment

Estrogen Response (Early/Late): Estrogen-driven gene expression

Presentation Roadmap



ORA vs GSEA Comparison

Design differences:

- Different inputs (binary gene list vs continuous ranking) and databases (GO:BP vs Hallmark)
- **Direct term-by-term comparison is not meaningful**

Shared:

- **Inflammatory/stress response** post-NAC
- **Cell adhesion decrease** post-NAC

GSEA only:

- **MYC Targets**
- **Fatty Acid Metabolism**

These signals are likely consistent across many genes but too moderate for individual genes to clear ORA's threshold

Interpretation and Insights

Fatty Acid Metabolism:

- ORA found this cluster enriched post-NAC
- Suggests surviving tumor cells switched from glucose to fatty acid oxidation as an alternative/ additional fuel source
- Lin et al. (2022) showed that chemotherapy can hurt glucose metabolism

Cell adhesion changes:

- ORA: cell adhesion genes (CDH18, KISS1) significantly downregulated, GSEA found the EMT Hallmark cluster enriched post-NAC
- Suggests surviving tumor cells are becoming more migratory
- Galluzzi et al., 2017 discussed this as a strategy for malignant cells

References

R Packages

- Yu G, et al. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* 16(5):284–287.
- Wu T, et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The Innovation* 2(3):100141.
- Korotkevich G, et al. (2021). Fast gene set enrichment analysis. *bioRxiv*. doi:10.1101/060012
- Robinson MD, et al. (2010). edgeR: a Bioconductor package for differential expression analysis. *Bioinformatics* 26(1):139–140.
- Carlson M. (2024). org.Hs.eg.db: Genome wide annotation for Human. R package version 3.19.1.
- Dolgalev I. (2022). msigdb: MSigDB Gene Sets for Multiple Organisms. R package version 7.5.1.

Gene Sets & Databases

- Ashburner M, et al. (2000). Gene Ontology: tool for the unification of biology. *Nature Genetics* 25(1):25–29.
- Subramanian A, et al. (2005). Gene set enrichment analysis. *PNAS* 102(43):15545–15550.

Visualization

- Shannon P, et al. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Research* 13(11):2498–2504.
- Merico D, et al. (2010). Enrichment Map: a network-based method for gene-set enrichment visualization. *PLoS ONE* 5(11):e13984.
- Kucera M, et al. (2016). AutoAnnotate: A Cytoscape App for Summarizing Networks with Semantic Annotations. *F1000Research* 5:1717.

Literature Support

- Daschner PJ, et al. (1999). Increased AP-1 activity in drug resistant human breast cancer MCF-7 cells. *Breast Cancer Research and Treatment* 53(3):229–240.
- Galluzzi L, et al. (2017). Immunogenic cell death in cancer and infectious disease. *Nature Reviews Immunology* 17(2):97–111.
- Takebe N, et al. (2015). Targeting Notch, Hedgehog, and Wnt pathways in cancer stem cells. *Nature Reviews Clinical Oncology* 12(8):445–464.

Questions?