

Mechanics of the Knee

Ralph Nisell

To cite this article: Ralph Nisell (1985) Mechanics of the Knee, Acta Orthopaedica Scandinavica, 56:sup216, 1-42, DOI: [10.3109/17453678509154159](https://doi.org/10.3109/17453678509154159)

To link to this article: <https://doi.org/10.3109/17453678509154159>



© 1985 Informa UK Ltd All rights reserved:
reproduction in whole or part not permitted



Published online: 08 Jul 2009.



Submit your article to this journal [↗](#)



Article views: 5618



Citing articles: 101 View citing articles [↗](#)

Mechanics of the Knee

A study of joint and muscle
load with clinical applications

Ralph Nisell

Kinesiology Research Group
Department of Anatomy
Karolinska Institute
P.O. Box 60400
S-10401 Stockholm
Sweden

ACTA ORTHOPAEDICA SCANDINAVICA SUPPLEMENTUM NO. 216, VOL. 56, 1985

MUNKSGAARD · COPENHAGEN

ISBN 87-16-06327-9

ISBN 0300-8827

Printed in Sweden by

Delta and Minab/Gotab, Stockholm, 1985

To Susanne
To my parents

ABSTRACT

The load moment of force about the knee joint during machine milking and when lifting a 12.8 kg box was quantified using a computerized static sagittal plane body model. Surface electromyography of quadriceps and hamstrings muscles was normalized and expressed as a percentage of an isometric maximum voluntary test contraction. Working with straight knees and the trunk flexed forwards induced extending knee load moments of maximum 55 Nm. Lifting the box with flexed knees gave flexing moments of 50 Nm at the beginning of the lift, irrespective of whether the burden was between or in front of the feet. During machine milking, a level difference between operator and cow of 0.70 m – 1.0 m significantly lowered the knee extending moments.

To quantify the force magnitudes acting in the tibio-femoral and patello-femoral joints, a local biomechanical model of the knee was developed using a combination of cadaver knee dissections and lateral knee radiographs of healthy subjects. The moment arm of the knee extensor was significantly shorter for women than for men, which resulted in higher knee joint forces in women if the same moment was produced. A diagram for quantifying patellar forces was worked out. The force magnitudes given by the knee joint biomechanical model correlated well with experimentally forces measured by others.

During the parallel squat in powerlifting, the maximum flexing knee load moment was estimated to 335 – 550 Nm when carrying a 382.5 kg burden and the *in vivo* force of a complete quadriceps tendon-muscle rupture to between 10,900 and 18,300 N. During isokinetic knee extension, the tibio-femoral compressive force reached peak magnitudes of 9 times body weight and the anteroposterior shear force was close to 1 body weight at knee angles straighter than 60 degrees, indicating that high forces stress the anterior cruciate ligament. A proximal resistance pad position decreased the shear force considerably, and this position is recommended in early rehabilitation after anterior cruciate ligament repairs or reconstructions.

The methods presented quantify muscle activity, sagittal knee joint moments and forces, enabling assessments to be made of different work postures, training exercises and joint derangements.

This thesis is based on the following papers which will be referred to in the text by their Roman numerals I-VI.

- I. Ekholm J, R Nisell, UP Arborelius, C Hammarberg & G Németh: Load on knee joint structures and muscular activity during lifting. *Scandinavian Journal of Rehabilitation Medicine*, 16:1-9, 1984
- II. Ekholm J, R Nisell, UP Arborelius, O Svensson & G Németh: Load on knee joint and knee muscular activity during machine milking. *Ergonomics*, 28:665-682, 1985
- III. Nisell R, G Németh & H Ohlsén: Joint forces in extension of the knee. Analysis of a mechanical model. *Acta Orthopaedica Scandinavica*, Accepted for publication, 1985
- IV. Nisell R & J Ekholm: Patellar forces during knee extension. *Scandinavian Journal of Rehabilitation Medicine*, 17:63-74, 1985
- V. Nisell R & J Ekholm: Joint load during the parallel squat in powerlifting and force analysis of *in vivo* bilateral quadriceps tendon rupture. Submitted for publication, 1985
- VI. Nisell R, MO Ericson, G Németh & J Ekholm: Tibio-femoral joint forces during isokinetic knee extension. Submitted for publication, 1985

CONTENTS

ABSTRACT	4
LIST OF PAPERS I-VI	5
ABBREVIATIONS AND DEFINITIONS	8
INTRODUCTION	9
AIMS OF THE STUDY	11
LITERATURE REVIEW	11
Knee load in different work postures	11
Tibio-femoral joint	11
Patello-femoral joint	12
Quadriceps rupture	13
Isokinetic exercise	13
MATERIALS	13
Subjects	13
Cadavers	14
METHODS	14
Calculation of load moments	14
Electromyographical recordings	15
Activities investigated	15
<i>Lifting</i>	15
<i>Work postures with hand-held implement</i>	15
<i>Parallel squat</i>	15
<i>Isokinetic knee extension</i>	16
Knee joint model	16
<i>Dissection</i>	16
<i>Radiography</i>	16
<i>Forces</i>	16
<i>Mechanical analysis</i>	18
Statistics	19
SUMMARY OF RESULTS	20
Lifting	20
Work postures with hand-held implement	21
Knee joint model	22
<i>Tibio-femoral contact point</i>	22
<i>Patellar tendon angle</i>	22
<i>Patellar tendon moment arm</i>	24
<i>Joint forces</i>	25
Quadriceps rupture	26
Isokinetic knee extension	28
DISCUSSION	28
Assumptions of the knee model	28
Compressive force per area	29
Knee joint load during various activities	30
Work postures	31
Clinical implications	32
CONCLUSIONS	34
ACKNOWLEDGEMENTS	36
REFERENCES	37

Abbreviations and definitions

BW	= Body weight
C	= Centre of contact pressure point between femoral and tibial condyles
deg	= Degree
de	= Moment arm of external force acting on tibia to bilateral knee axis
dp	= Knee extensor moment arm, perpendicular distance of patellar tendon force to tibio-femoral contact point
EMG	= Electromyography
Fcp	= Compressive force in patello-femoral joint, perpendicular to joint surface
Fcq	= Compressive force between quadriceps tendon and femoral intercondylar groove
Fct	= Compressive force in tibio-femoral joint, perpendicular to joint surface
Fet	= External force acting on tibia tending to flex the knee
FKC	= Lift with flexed knees and box between feet (close to body)
FKFF	= Lift with flexed knees and box in front of feet (far from body)
Fp	= Force in patellar (infrapatellar) tendon
Fq	= Force in quadriceps (suprapatellar) tendon
Fqm	= Force in quadriceps muscle
Fs	= Anteroposterior shear force in tibio-femoral joint along joint surface. Positive value indicates when tibia tends to slide anteriorly in relation to femur, negative when sliding posteriorly
kg	= Kilogram
m	= Meter
N	= Newton
Nm	= Newtonmeter
sec	= Second
SK	= Lift with straight knees
TAMP-R	= Time averaged myoelectrical potential ratio, <i>i.e.</i> muscle activity recorded during the experiment divided by the reference level recorded during an isometric maximum voluntary test contraction (normalized EMG)
WCR	= Working cycle ratio during lifting, runs from 0 to 1
α (alfa)	= Knee flexion angle, between long axes of tibia and femur. Straight knee equals 0 deg
β (beta)	= Angle between Fp and perpendicular line of tibial plateau. Positive when tibia is pulled anteriorly in relation to femur
γ (gamma)	= Angle between Fp and long axis of tibia. Positive when the intersection is below the tibial plateau
δ (delta)	= Angle between external force and tibial plateau
ϵ (epsilon)	= Angle between Fq and Fcp
λ (lambda)	= Angle between Fq and Fqm
ψ (psi)	= Angle between Fp and Fq
ω (omega)	= Angle between tibial plateau and perpendicular line of tibial long axis

Introduction

The concept of biomechanics may be defined in different ways. A definition recently suggested by LeVeau (1984) is: "the study of forces and the effects of those forces on and within the human body". Hence, musculoskeletal load in all its aspects becomes a substantial part of the biomechanical concept. Biomechanics has become an interdisciplinary between existing fields such as mechanical engineering which develop methods, equipment and accuracy, and medicine where for example ergonomics, orthopaedics and rehabilitation abound with practical questions and applications. As a consequence, one of the main goals for the medically and biomechanically trained, might be to understand and use engineering knowledge and methods to solve practical, clinical and preventive problems seen in medical care among patients. In this way the biomechanical field of science may function as a bridge between medical and engineering specializations and, as such, it has become a new and most fascinating field with a variety of practical applications (*e.g.* Frankel & Burstein 1972, Carlsöö 1975, LeVeau, Williams & Lissner 1977, Winter 1979, Frankel & Nordin 1980, Dowson & Wright 1981, Maquet 1984, Chaffin & Andersson 1984).

Quantifying joint moments, forces and muscle activities in various situations or during different activities may be of value from the medical viewpoint, with applications in three different areas:

1) In *ergonomics*, to study the amount of mechanical load and thus to indicate possible load reduction during work; to assess harmful effects of work situations and for designing work equipment; to reduce pain from the locomotor system and load elicited diseases; to improve possibilities of assessing the long-term cause of workload-related diseases and, in a longer perspective, to reduce the number of sick-leave days due to such symptoms.

2) In *orthopaedic surgery*, to assess mechanically the effects of bone and joint derangements (such as those following osteotomy, patellectomy or fractures) and thus to influence the pre- and postoperative investigation of patients; to obtain criteria for material strength in the design of endoprosthesis; to optimize post-surgical rehabilitation after for example anterior cruciate ligament repairs and reconstructions.

3) In *rehabilitation*, to individualize and optimize physical therapy without risks for damaged, injured, diseased or weak structures; to favour exercises that entail a high degree of muscle activity while keeping the load on joint structures low; to estimate the physiological load during normal activities (*e.g.* walking, running, stair climbing, taking a step down, rising from a chair) so as to avoid extreme loads; to improve movement patterns and performance and avoid harmful exercises, which is important also in sports medicine.

"Joint load" is not a very well-defined concept (Németh 1984). The moment of force (= moment, sometimes also called torque) acting about a joint axis is often a useful way to express either load (= load moment) or strength (= muscle moment). However, for different joint angles but the same moments, the joint forces may vary considerably due to changes in muscle moment arm lengths and, consequently, the muscle force magnitudes change. Quantifying the joint force is preferred as it better describes the loading effect on the joint surface. The joint force can be either in the normal (= compressive) or tangential (= shearing) direction of the joint surface or, most commonly, a combination of both. The joint compressive force acts over an area, which indicates that another way to express joint load would be in terms of pressure (N/m^2). This is probably a very good parameter of "joint load". When the menisci are removed from the knee joint, the

compressive force magnitude will remain the same, but the pressure will be larger as the tibio-femoral contact area is considerably smaller (Walker & Erkman 1975, Seedhom 1979, Fukubayashi & Kurosawa 1980). This increased joint load is probably the main reason why degenerative changes are frequent many years after meniscectomy (e.g. Fairbank 1948, Jackson 1968, Johnson et al. 1974). In this thesis the expression "knee joint load" is synonymous with knee load moment and/or joint force.

"Muscle load" can be studied in many ways. Direct EMG was earlier used frequently (e.g. Basmajian 1962, Pocock 1963, Jonsson & Synnerstad 1966, Hallén & Lindahl 1967, Carlsöö 1975). Today normalized and thus quantified EMG has been developed and has definite advantages as it is possible to compare the EMG-level between muscles of different subjects (Ekholm et al. 1978, Philips & Petrofsky 1983, Reynolds et al. 1983, Soderberg & Cook 1983, Andriacchi et al. 1984, Brask et al. 1984, Németh 1984, Yang & Winter 1984). Another approach is to quantify muscle load as a ratio between the muscle moment produced during an activity and the maximum strength measured during an isometric voluntary test contraction (Schüldt et al. 1983, Németh et al. 1984, Svensson et al. 1985). These kinds of "muscle load" quantification were used in Studies I and II where the EMG was normalized and expressed as a percentage of an isometric maximum voluntary test contraction. Muscle load has also been frequently studied by EMG frequency or amplitude distribution, which makes fatigue estimations possible (e.g. Kadefors et al. 1968, Komi & Tesch 1979, Kadefors 1978, Petrofsky 1979, Hagberg 1979, Herberts et al. 1980). Muscle force magnitude as a function of EMG is not always linear, especially at muscle activities close to maximum (e.g. Möller 1966, Zuniga et al. 1970, Bouisset 1973, Vredenburg & Rau 1973, Jonsson 1978, Woods & Bigland-Ritchie 1983). The EMG-force relation seems to change between muscles in the same individual and between the same muscle of different individuals, probably due to differences in muscle fibre composition (Woods & Bigland-Ritchie 1983). So, unfortunately, the muscle force magnitudes exerted cannot easily be read from the EMG.

There are, in principle, two different methodological approaches of quantifying joint load; 1) *experimental measurements* and, 2) *biomechanical models* and of course these two methods may be combined. In experimental measurements, force, pressure, etc., are measured directly by a specific sensor (e.g. strain gauge, force plate, pressure sensitive film). In the biomechanical modelling approach, the unknown parameter is calculated indirectly using reasonable approximations, mechanical laws and if any, only a few pure experimental measurements. The advantage of a model, if good, is that, besides being non-invasive, it can be generally applied to many different situations without performing new, time-consuming experiments. However, it is important that such a model is realistic. Its validity can be studied by comparing experimental measurement results with results from the model. If there is a bad correlation, either the experimental set up or the biomechanical model solution, or both, will be inadequate for describing the reality.

This thesis presents biomechanical methods for quantifying knee joint load during various activities and exercises. Examples of knee load magnitudes are reported and the effects and implications of the induced load are discussed. The present work is one part of a series of investigations from the Kinesiology Research Group (e.g. Ekholm et al. 1978, Arborelius & Ekholm 1978, Ekholm et al. 1982, Schüldt et al. 1983, Harms-Ringdahl et al. 1983, Németh 1984, Svensson et al. 1985, Ericson et al. 1985a and 1985b) concerning levels of muscle activity and the loads on the major joints during various activities and exercises.

Aims of the Study

The general aim of the present investigation was to quantify the load induced to the knee joint in some specific situations or activities. At the beginning of the work on this thesis, only the knee load moment was quantified (Studies I and II), but it was soon obvious that there was a need for a local knee biomechanical model to determine the joint forces induced during the activities. Therefore, a general knee model was worked out (Studies III and IV). The purpose of this model was to be a simplistic and useful, but still realistic, two-dimensional biomechanical model that could be applied to any knee extending activity. In Studies V and VI, examples of how the model can be applied are given.

More specifically, the following topics have been studied:

- (1) The muscle activity and load moment about the bilateral knee axis during different standing work postures with the trunk flexed forwards (machine-milking) and during lifting a box, with the knees either flexed or straight.
- (2) A local knee biomechanical model for quantifying the forces in the tibio-femoral joint and in the patello-femoral joint.
- (3) The *in vivo* force magnitude in the quadriceps tendon at the moment of a traumatic rupture.
- (4) The magnitude of the tibio-femoral joint forces during isokinetic knee extension and the stress on the anterior cruciate ligament.
- (5) The clinical and practical applications of the results in fields such as ergonomics, orthopaedics and rehabilitation

Literature review

Knee load in different work postures

When studying the mechanical load during work with the trunk leaning forwards, the main interest hitherto has been focused on the load induced to the low back (*e.g.* Andersson et al. 1976, Ekholm et al. 1982, Nordin 1982, Chaffin & Andersson 1984). However, dysfunction of the knee joint is also, along with that of the other major joints, a common cause of impaired working capacity among unskilled labourers (Undeutsch et al. 1982). Mechanical knee joint loads in different standing work situations have not earlier been reported in the literature.

Tibio-femoral joint

In a biomechanical model of knee extension, the patellar tendon moment arm length is decisive for the magnitudes of the forces calculated. The moment arm length has been determined mainly in three different ways;

1. Haxton (1945) and Kaufer (1971) calculated the moment arm at different knee flexion angles by using measured data of flexing knee moments which were related to counteracting forces from the knee extensor. The same method was also recently used by Wendt & Johnson (1985).
2. Lindahl & Movin (1967) and Lindahl et al. (1969) using lateral radiographs showed how the tibio-femoral force magnitudes may be calculated when using the distance between the tibio-femoral contact point (C) and the patellar tendon as the knee extensor moment arm. Several authors have subsequently used a similar "tibio-

femoral contact” approach in their biomechanical knee models (e.g. Reilly & Martens 1972, Haffajee et al. 1972, Perry et al. 1975, Seedhom & Terayama 1976, Bishop 1977, Ellis et al. 1979).

3. The moment arm length has been determined on lateral x-rays as the perpendicular distance between the sagittal knee centre axis of rotation and the patellar tendon (Bandi 1972, Smidt 1973). Smidt (1973) presented useful average data of knee joint moment arms from 26 normal men in order to calculate the joint forces during maximum isometric knee extension and flexion.

Bresler & Frankel (1950), using ground reaction forces and mechanical equations, calculated the maximum extending muscular moment about the knee joint during human gait to be between 40 Nm and 88 Nm. Similar methods have subsequently been used for studying joint load during level walking (Radcliffe 1962, Winter 1980, Boccardi et al. 1981) and by Morrison (1968, 1969 and 1970) who related the calculated load to anatomical data of one knee dissection in order to determine knee joint and ligament forces. Morrison (1970) calculated the maximum mean tibio-femoral compressive force (Fct) to be about 3 times BW during level walking. When walking up stairs the maximum Fct was above 4 BW and down stairs just below 4 BW (Morrison 1969). Andriacchi et al. (1980) calculated the knee extensor muscular moment to 54 Nm when ascending stairs and as high as 147 Nm when descending.

Tibio-femoral joint load has also been quantified by means of biomechanical models in other activities such as flexed knee stance and rising from seated and squatting positions (Seedhom & Terayama 1976, Bishop 1977, Ellis et al. 1979 and 1984, Kelley et al. 1978, Dahlqvist et al. 1982, Schüldt et al. 1983), jumping (Alexander & Vernon 1975, Smith 1975, Nisell et al. 1985), kicking (Wahrenberg et al. 1978, Lindbeck 1983) and bicycling (Gregor et al. 1985, Ericson et al. 1985b).

Patello-femoral joint

Several biomechanical models have suggested how to determine the magnitude of the patello-femoral compressive force (Fcp). The oldest and most easily-used model assumes the quadriceps (Fq) and patellar (Fp) tendon forces to have the same directions as the tibial and femoral long axes. Other, more advanced, models may consider the angle of the patellar tendon in relation to the tibial long axis (Reilly & Martens 1972, Bandi 1972, Smidt 1973, Ellis et al. 1979, Hungerford & Lennox 1983) or as Matthews et al. (1977), who suggested a special linear relationship between the knee flexion angle and angle ψ . However, all these models consider Fp to have the same magnitude as Fq throughout the whole range of motion.

As early as 1945, Haxton showed that the patellar and quadriceps tendon forces for the flexed knee joint are not of the same magnitude, the patellar tendon taking up lower forces than the quadriceps tendon. In recent years several authors have reported the same at knee flexion angles above 60 deg, where the Fp/Fq ratio is suggested to range between 0.50 – 0.75 (Bishop 1977, Bishop & Denham 1977, Ellis et al. 1980, Maquet 1984, Huberti et al. 1984).

Reilly & Martens (1972) calculated Fcp to 0.5 times BW during level walking and 3.3 BW during stair-climbing. Smidt (1973) calculated Fcp at isometric maximum knee extension to 2.6 BW. Dahlqvist et al. (1982) estimated maximum Fcp values during squatting to be between 4600 N and 5900 N or about 7 BW. Fcp has also been estimated during rising from a chair with and without the aid of the arms (Seedhom & Terayama 1976, Ellis et al. 1979 and 1984).

Quadriceps rupture

Bilateral simultaneous complete rupture of the quadriceps tendons is a rare injury with 24 cases reported in the literature (Frey 1928, Goldenberg & Paterson 1949, Steiner & Palmer 1949, Wetzler & Merkow 1950, Wilson 1957, Preston & Adicoff 1962, Dalal & Whittam 1966, MacDonald 1966, Levy et al. 1971, Firooznia et al. 1973, Brotherton & Ball 1975, Peiro et al. 1975, Morein et al. 1977, Siwek & Rao 1978, Lotem et al. 1978, Grenier & Guimont 1983, MacEachern & Plewes 1984, Bhole et al. 1985). Most of the patients were elderly and the causes of the tendon ruptures were reported to be degenerative diseases or general metabolic disorders such as gout, obesity, uraemia, hyperparathyroidism or rheumatoid arthritis.

During power lifting, two traumatic cases of knee extensor ruptures have been reported, both at the deepest position of the parallel squat (Grenier & Guimont 1983, Bjerneld et al. 1984) but no biomechanical investigations were made. Zernicke et al. (1977) have performed the only reported biomechanical calculations regarding an *in vivo* tendon force magnitude at the time of rupture. Their subject was a 29-year-old (82 kg) champion weight-lifter who ruptured his right infrapatellar tendon during a "clean and jerk" lift of 175 kg. The load moment was calculated to 550 Nm and the patellar tendon force to 14,500 N or 17.5 times BW at the time of rupture.

Isokinetic exercise

The isokinetic exercise was first reported in 1967 by Hislop & Perrine and Thistle et al. The isokinetic way of exercise is very efficient in muscle strengthening as the muscle may exert maximum effort throughout the range of motion (*e.g.* Thistle et al. 1967, Pipes & Wilmore 1975, Grimby et al. 1980). In addition, isokinetic exercise is very useful in the study and understanding of biomechanics, kinesiology and physiology because of its standardized motion and its possibility to record directly the joint torque produced. Hence, isokinetic exercise has been widely used for many different purposes for example as a method for analysing force-velocity or power-velocity relationships (*e.g.* Thorstensson et al. 1976, Perrine & Edgerton 1978, Ingemann-Hansen & Halkjaer-Kristensen 1979) or as a training/testing device for healthy athletes (Öberg 1985).

In clinical practice, the isokinetic exerciser has become frequently used for example to assess knee rehabilitation after meniscectomy or ligament injury (Campbell & Glenn 1979, Grimby et al. 1980, Arvidsson et al. 1981, Hamberg & Gillquist 1984, Murray et al. 1984), patello-femoral disorders (Nordesjö & Nordgren 1978, Rauschnig et al. 1983, Nordgren et al. 1983, Lysholm et al. 1984), ankle joint function (Nistor et al. 1979, Markhede & Nistor 1979, Fugl-Meyer 1980, Tropp 1985), rehabilitation after femoral shaft fracture (Mira et al. 1980), and spasticity after hemiparesis (Knutsson & Mårtensson 1980). Despite the widespread use of isokinetic exercise, there are no biomechanical reports concerning the forces induced in the knee joint, except for the work by Johnson (1982) who calculated the tibio-femoral shear forces and showed that a dual-pad resistance considerably diminished the stress on the anterior cruciate ligament. However, the compressive force magnitudes were not estimated.

Materials

Subjects

In Study I, fifteen healthy subjects (one woman) volunteered to participate. Mean age was 28 years, weight 76 kg, and height 1.78 kg. During the course of Study I, a new

EMG technique and biomechanical approach were developed. Therefore, only seven subjects were used for moment calculations and five of them for the presentation of EMG recordings.

As subjects in Study II, ten professional milkers (four women) were used with machine milking experience of between 1 and 40 years. The subjects mean age was 31 years, weight 72 kg, and height 1.74 m.

In Studies III and IV, twenty healthy subjects (10 women and 10 men) were radiographed. Their mean age was 23 years (women) and 27 years (men), mean height was 1.67 m (women) and 1.82 m (men), mean weight 59 kg (women) and 75 kg (men). The subjects were without knee disorders or dysfunctions and had not undergone any prior knee surgery.

Four highly-trained power lifters participated in Study V. Their age ranged from 21–34 years, height from 1.64–1.91 m, weight 74–146 kg, and personal record in the parallel squat from 305–430 kg. In a Scandinavian Championship one of the subjects (31 years, 1.91 m, 146 kg) experienced a bilateral simultaneous quadriceps tendon/muscle rupture during a parallel squat while carrying 382.5 kg.

In Study VI, eight healthy men, free from any lower extremity disorder, volunteered as subjects. Their mean age was 27 years, mean weight 72.4 kg, and mean height 1.80 m.

Cadavers

In Studies III and IV, twenty left knees of cadavers (10 women and 10 men) were dissected. Their mean age at death was 76 years (women) and 68 years (men). Their mean height was 1.62 m (women) and 1.74 m (men).

Methods

Calculation of load moments

(Studies I, II, V and VI) In studies I, II and V body position was photographed perpendicularly to the sagittal plane with a motorized camera (Olympus OM-1) at a picture frequency of 4 frames/sec. To make time synchronization possible, a timer with a light-emitting diode display was used. Either photocopies or negative images were projected on a semiautomatic coordinate registration device (Tektronix digitizer, 4953). The digitizer recorded the coordinates of the reference distance points and the positions of the bilateral motion axes of the major joints. The digitizer was connected to a graphics terminal (Tektronix 4012) which was connected to a Nord-100 computer. A sagittal plane mechanical body model based on static mechanics was designed and the load moments about the bilateral knee joint axis and about other major joint axes were calculated (Ekholm et al. 1982). The body segment parameters necessary for the calculations were taken from the anthropometric data given by Dempster (1955), and the moment caused by the burden weight was included in the calculations. The program allowed a change of the burden weight parameter, and this function was used to calculate the contribution of the body segments to the total load moment.

In study VI, a Cybex II isokinetic dynamometer was used connected to an Apple II computer. Controlled acceleration was performed through the computer, and the gravity moments caused by the weights of the leg, foot and resistance arm were corrected for with software (Gransberg & Knutsson 1983).

Electromyographical recordings

(Studies I and II) Muscle activity was recorded by means of EMG from the muscle bellies of the quadriceps and hamstrings muscles. Two flexible disposable Ag-AgCl surface electrodes were attached to the skin over each muscle belly in the direction of the muscle fibers. The distance between the electrode centres was 35–40 mm. The myoelectrical signals were amplified in a Devices AC8 and full wave rectified low pass filtered electromyograms (=linear envelope, Winter 1979) were recorded, using heat-sensitive paper. The bandwidth was 10 Hz to 1.5 kHz and the time integral constant 0.50 sec. For control purposes, amplified unfiltered direct EMG signals were recorded in parallel on a UV-recorder (Honeywell Visicorder, 1508).

For comparison of levels of muscle activity between different muscles and between different individuals, an EMG normalization was performed. The muscle activity was presented as a ratio, TAMP-R, between the EMG activity recorded during the experiments and the activity recorded during an isometric voluntary maximum test contraction. To standardize the isometric test contractions the subject was firmly bound to a specially designed chair and knee angle was held constant at 90 deg for both knee extension and flexion.

Activities investigated

Lifting (Study I)

A box weighing 12.8 kg placed on the floor in front of the subjects was lifted to a bench at the same level as the subject's umbilicus. Three different lifts were studied; 1) Lift with straight knees (SK), 2) Lift with flexed knees and the box in front of the knees (FKFF), 3) Lift with flexed knees and the box between the feet and thus close to the trunk. All the lifts were performed at a moderate speed, *i.e.* approximately two seconds. To make comparisons possible between the different modes of lifting and between the subjects, time was expressed as a WCR, from 0 to 1. A strain gauge was mounted in each box handle to measure the vertical acceleration of the box and thus obtain an idea of the dynamic force components.

Work postures with hand-held implement (Study II)

Twenty different selected postures were studied in our laboratory at the most loading part of the machine milking procedure, *i.e.* when the teat-cups, weighing 3.3 kg (2.8 kg in the right hand and 0.5 kg in the left hand), were attached to the udder. To simulate pit depth, vertical level differences between milker and cow of 0.17, 0.33, 0.50, 0.70, 0.85, and 1.0 m were arranged. The horizontal distance from the centre of the udder to the milker was 0.50 m. At 1.0 m level difference, horizontal distances of 0.6 and 0.7 m were also studied. Standing postures with the knees straight and, at the lower level differences (0.50 m and less) also with flexed knees, were studied. Nine different postures at zero level-difference were also studied, one of them sitting on a one-legged stool.

Parallel squat (Study V)

In power-lifting, the squat is performed by supporting a weight on the shoulders in a standing position, from which the body is lowered by flexing the knees, so the proximal part of the thighs will become lower than the knees. Three of the four subjects performed parallel squats and the deepest position and the ascending phase were

studied. Each subject performed squats in his normal way and speed *i.e.* 1.25 – 1.50 sec for the ascent. Burden weights of between 60 and 260 kg were studied. Using the sagittal plane body model described, the subjects were simulated to the same body size as the subject injured with the bilateral quadriceps rupture.

Isokinetic knee extension (Study VI)

The subjects were seated with back support and the thighs were attached by straps to the table. The arms were held along the trunk and the hands gripped the table sides. The bilateral motion axis about which flexion and extension of the knee occur was aligned with the axis of the resistance arm. The following three knee extensions were conducted in a randomized sequence: 1) velocity of 180 deg/sec, resistance pad distal; 2) velocity of 180 deg/sec, resistance pad proximal; 3) velocity of 30 deg/sec, resistance pad distal.

The distal pad position was just proximal to the ankle joint on the anterior part of the lower at a mean distance of 0.37 m (SD=0.02) from the motion axis. The proximal pad position was 0.20 m from the axis. The knee extension movement started at 90 deg knee angle and finished at straight knee (= 0 deg).

Knee joint model (Studies III and IV)

Dissection

The cadaver's left patella was removed. The thickness of the patellar and quadriceps tendons was measured at their insertions on the patella. The measured distances ap, gt, and bq (Fig. 1) were used to find the tendon midpoints T, P, and Q (Fig. 1) on the radiographs.

Radiography

Lateral x-ray films (30×40 cm) were taken of the loaded knee with the subjects standing at knee flexion angles (α , Fig. 1) ranging from straight (=zero deg) to 120 deg flexed. One anteroposterior and one lateral radiograph picture were taken with the knee straight and the quadriceps activated. The film-focus distance was 1.55 m and the knee centre – film distance was 0.15 m, giving a magnification factor of 1.11 which was corrected for.

The tibio-femoral contact point (C, Fig 1) was defined as the midpoint of the smallest distance between the femoral and tibial condyles. The long axes of the femur and tibia were defined from the radiographs as the lines between the two shaft midpoints located 75 mm and 150 mm from the condyle surfaces.

Forces

Many force vectors act on the bone through one particular tendon. All biomechanical models use approximations, and in this study the forces in the various parts of the tendon were presumed to act as one total single force vector from tendon midpoint at one place to tendon midpoint at the other. The same approach in biomechanical modelling has been used earlier (Jensen & Davy 1975, Dostal & Andrews 1981, Németh & Ohlsén 1985). This assumption is fair for straight tendons, but at knee angles of 90 deg or more where the quadriceps and patellar tendon are no longer straight, two extra points (E and K) were defined (Fig. 1). Point E was located on the radiographs as the midpoint of the quadriceps tendon, 15 mm proximal to the patellar base. Point K was located at a distance corresponding to half patellar tendon thickness in front of the anterior border of the tibial plateau, and the direction of the patellar tendon force (F_p)

was defined as the straight line between points K and P.

The compressive force in the patello-femoral joint (F_{cp}) was projected through the centre of the patello-femoral contact point (M), perpendicularly to the joint surface (Fig. 1). The force in the quadriceps tendon (F_q) was assumed to run parallel to the long axis of femur at knee angles of 30 deg and 60 deg. F_q passes through the midpoint of the quadriceps tendon insertion at the patellar base (point Q). To find point Q on the x-rays, the proximal dorsal bony corner of the patellar base (point B) was identified. The distance between these two points, bq-distance, was estimated in the morphological study and its mean value was used on the x-rays. The corresponding approach was used to locate the tendon midpoints at patellar apex (P) and at tuberositas tibiae (T). At the straight knee the direction of F_q was determined as the line between the F_p – F_{cp} intersection point (X) and the midpoint of the quadriceps tendon (point Q) at the patellar base.

At knee angles 90 deg and 120 deg the quadriceps tendon was curved in the femoral intercondylar groove. The force magnitude in the proximal part of the quadriceps muscle (F_{qm}) was considered to be the same as in the distal part (F_q) inserting on the patellar base. The angle between these two forces was defined the λ angle. It gave rise to a force between the distal part of the quadriceps tendon and femoral intercondylar groove (F_{cq}), (Fig. 1). At knee angles above 60 deg, the direction of F_q was estimated as the line between points E and Q (Fig. 1).




Fig. 1. Forces, angles, distances and bony landmarks at 30 deg knee angle (α). The patellar tendon moment arm (dp) is the perpendicular distance between patellar tendon force (F_p) and centre of the tibio-femoral contact (C). F_p acts through proximal patellar tendon midpoint (P) with a distance "ap" from patellar apex (A) and through distal midpoint (T) with distance "gt" from posterior tendon border (G). Tibial plateau slopes backwards (ω , women 7.2 deg and men 9.2 deg) in relation to tibial long axis. The tibio-femoral forces F_{ct} and F_s act through C perpendicularly to and along the joint surface, respectively. Quadriceps tendon force (F_q) parallels femoral long axis and goes through tendon midpoint (Q) at the insertion on patellar base. Patello-femoral compressive force (F_{cp}) projects through F_p - F_q intersection point (X) and centre of patello-femoral contact point (M), perpendicularly to the joint surface.

Mechanical analysis

The tibio-femoral contact point (C) was chosen as the origin for the calculation of moments. For the tibia in a free body diagram and in a static situation (Fig. 2a), the moment (M) about C can be determined:

$$M = (de \times Fet) + (df \times Fef) + (dm \times mg) \quad (1)$$

where Fet , Fef and mg are the "external" forces acting on the tibia and de , df , and dm their moment arms. The force in the patellar tendon (Fp) is given by:

$$Fp = M / dp \quad (2)$$

where dp is the moment arm of Fp . From Fig. 2a, we get the vector equation:

$$Fp + Fct + Fs + Fet + Fef + mg = 0 \quad (3)$$

Projecting (3) in the normal (or Fct) direction of the tibial plateau;

$$Fct = Fp \times \cos\beta + Fet \times \sin\delta_1 + Fef \times \sin\delta_2 - mg \times \sin\delta_3 \quad (4)$$

in tangential (or Fs) direction;

$$Fs = Fp \times \sin\beta - Fet \times \cos\delta_1 - Fef \times \cos\delta_2 - mg \times \cos\delta_3 \quad (5)$$

where β , δ_1 , δ_2 and δ_3 are angles in relation to tibial plateau. In the example above the external forces act on the anterior border of tibia and on the sole of the foot, but it is obvious that these forces may act anywhere on the "free body".




Fig. 2a. Free body diagram of lower leg. The magnitudes of the patellar tendon force (Fp), tibio-femoral shear (Fs) and compressive force (Fct) can be calculated (see equations above) if the external forces (Fet , Fef and mg), their angles in relation to the tibial plateau (δ_1 , δ_2 and δ_3) and their moment arms (de and df) have been determined.




Fig. 2b. Free body diagram of patella and of the quadriceps tendon. Patello-femoral compressive force (F_{cp}) equals the vector sum of patellar tendon force (F_p) and quadriceps tendon force (F_q). F_{cp} projects through F_p - F_q intersection point (X) and centre of patello-femoral joint contact point (M), perpendicularly to the joint surface. Tendon-bone compression (F_{cq}) equals the vector sum of F_q and F_{qm} . Tendon midpoints E and K are used because the knee extensor tendon is curved at 90 and 120 deg knee flexion angles.

Three forces act on the patella. If we consider the patella in a free body diagram (Fig. 2b) we get the vector equation:

$$\mathbf{F}_p + \mathbf{F}_q + \mathbf{F}_{cp} = \mathbf{0} \quad (6)$$

From basic trigonometry we know that:

$$F_{cp} = F_p (\sin \psi) / (\sin \epsilon) \quad (7)$$

Projecting the vectors F_{cp} and F_p on F_q gives the magnitude of F_q :

$$F_q = F_{cp} (\cos \epsilon) + F_p (\cos \psi) \quad (8)$$

Considering the distal part of the quadriceps tendon in a free body diagram we get the vector equation:

$$\mathbf{F}_q + \mathbf{F}_{qm} + \mathbf{F}_{cq} = \mathbf{0} \quad (9)$$

As forces F_q and F_{qm} are assumed to have the same magnitude we get from equation (9):

$$F_{cq} = 2 F_q (\sin \lambda/2) \quad (10)$$

Statistics

Two- and three-way analysis of variance (ANOVA) were used in Studies I-IV. Multiple comparisons using studentized range test ($p = 0.01$) were used if ANOVA showed significant difference at 0.01 level. In studies III, IV and VI, Student's unpaired or paired t-tests were used. The significance levels chosen were consistently $p = 0.01$.

Means with 95 % confidence intervals were calculated in Studies I-IV and means with standard deviations in Study VI. For the EMG-recordings in Study II, the medians with their 95 % confidence intervals were calculated.

Summary of Results

Lifting (Study I)

Lifting the box with straight knees (SK) gave throughout the lift an extending knee load moment which was highest at the beginning and at the end of the lift (Fig. 3). In the middle of the lift (WCR=0.6) the load was significantly lower. Lifting with flexed knees (FKFF and FKC) induced a flexing knee moment of about 50 Nm at the beginning of the lift, and during the course of the lift the moment changed to an extending direction. There was no significant difference in load moment between the two lifts with flexed knees. During the final phase (WCR 0.8 and 1.0) of all lifts, there was no load difference between the lifts, and the extending knee moment was about 55 Nm. During the SK-lift, the knee extensor muscle showed low activity throughout. The hamstrings, which are knee flexors and also hip extensors, were activated to a medium level during the first third of the lift. During the flexed knee lifts, there was medium to high activity level in the knee extensors. Although to a low level, the hamstrings were activated, indicating that a certain co-contraction was present.




Fig. 3. Diagrams above: load moment (y-axis) about bilateral knee axis during the course (x-axis, WCR) of the three different lifts SK, FKFF and FKC. Positive values show extending load moment, negative flexing load moment. Upper curve: total load moment. Lower curve: moment due to body segments only (burden subtracted). Means and 95 % confidence intervals, $n = 7$. **Diagrams below:** level of EMG-activity in TAMP-R in five knee muscles, $n = 5$.

Work postures with hand-held implement (Study II)

In the straight knee postures, a level difference of 1.0 m between milker and cow showed the lowest extending moment (Fig. 4). At small level differences (0.0 m – 0.50 m) the knee moment was significantly higher, and some subjects utilized a high percentage of the muscle capacity. Increased horizontal distance between milker and cow induced higher knee extending moments. In flexed knee postures the knee load was of flexing direction, and the vastus lateralis muscle was activated between medium and high level and more than the rectus femoris muscle. In both straight- and flexed-knee postures, the biceps femoris muscle was activated between low and medium levels. A knee flexion angle of 65 deg gave a knee moment close to zero, regardless of body weight.




Fig. 4. Knee load moment in different standing working postures (x-axis) illustrated above. Filled circles = straight knees. Unfilled = flexed knees. Positive values indicate extending knee load moment, negative values flexing moment. The figures on the x-axis indicate different level differences between operator and cow. At 1.0 m three different horizontal distances between cow and operator were studied. 95% confidence intervals of the means. $n = 9$ for 0.0 m. $n = 5$ for 1.0 m with horizontal distance 0.6 and 0.7 m. $n = 10$ for the remaining postures. Reprinted with kind permission from *Ergonomics* (Study II); copyright 1985 by Taylor & Francis Ltd.

Knee joint model (Studies III and IV)

Tibio-femoral contact point

The tibio-femoral contact point (C) moved anteriorly on the tibial plateau when the knee was extended (Fig. 5). Over the last 30 deg of knee extension, point C was much displaced. The displacement of C during extension from 120 deg to straight was about 20 mm, or 40 % of the tibial plateau sagittal length.

Comments. The displacement pattern of point C indicates that a considerable rolling motion occurs in the tibio-femoral joint during the last deg of knee extension. The results are mainly in agreement with earlier reports (Zuppinger 1904, Lindahl & Movin 1967, Walker & Hajek 1972). However, this study showed a tibio-femoral rolling motion also at more flexion angles greater than 30 deg. A point C displacement of as much as 20 mm has not earlier been reported.




Fig. 5. Location of tibio-femoral contact point (C, y-axis) at various knee flexion angles (α , x-axis). 95 % confidence intervals of the means are indicated. Y-axis shows the displacement in % of the sagittal tibial plateau length. 100 % equals the anterior border of the plateau.

Patellar tendon angle

The magnitude of the patellar tendon angle γ varies with knee angle as shown in Fig. 6a. The horizontal axis shows the radiographically determined knee angle (α) which for the straight knee was more negative for women (mean -11.7 deg) than for men (-7.8 deg). The angle β is of greater biomechanical interest than γ , as β represents the patellar tendon direction in relation to the tibial plateau. β equals the sum of γ and ω . The magnitude of β increases in the same way as γ during knee extension. These two angles change from a negative magnitude to a positive at a knee angle of 75 and 100 deg, respectively.

Comments. The magnitude of γ at various knee angles has been reported by Smidt (1973). Smidt's pattern of change is similar to that in the present study but consistently somewhat higher. The magnitude of β at various knee angles is of importance when




Fig. 6a. Magnitude of γ (y-axis) at various knee flexion angles (α , x-axis). 95 % confidence intervals of the means are drawn for both γ and α . Zero deg α equals straight knee. The difference in γ is statistically significant between men and women.




Fig. 6b. Magnitude of β (y-axis) at various knee flexion angles (α , x-axis). 95 % confidence intervals of the means are indicated. Note the intersection of the zero-line at about 100 deg knee angle.

constructing a local biomechanical model of the knee but has not earlier been determined. When the knee is straight the patellar tendon pulls tibia forwards in relation to femur due to β being positive and when the knee is flexed to more than 100 deg, the patellar tendon pulls tibia backwards.

Patellar tendon moment arm

The maximum length of the patellar tendon moment arm was found at 30 to 60 deg knee flexion angle, and the shortest at the straight knee (Fig. 7). The same pattern was seen for both sexes but women had significantly shorter moment arms than men throughout the whole range of motion.




Fig. 7. Patellar tendon moment arm (dp, y-axis) for men and women at various knee flexion angles (x-axis). 95 % confidence intervals of the means are indicated.

Joint forces

Fig. 8 shows an isometric knee extension against a resistance, F_{et} , applied at a distance, d_e , perpendicularly to the anterior side of the distal leg, with the subject lying on one side. As women had shorter moment arms than men (Fig. 7), the calculated forces for women were consistently about 20 % higher. The patellar tendon force (F_p) and tibio-femoral compressive force (F_{ct}) were of the same magnitude and did not change dramatically over the various knee angles. When the knee was extended by a moment of 40 Nm, it was found that F_p and F_{ct} were about 1100 N at knee angles of 30–120 deg (Fig. 8 a+b). For the straight knee these forces were higher, about 1230 N and 1400 N respectively. The tibio-femoral shear force (F_s) changed sign between 50 and 90 deg flexion angle, depending on the magnitudes of F_{et} and d_e (Fig. 8c). F_s was around –200 N at 120 deg knee angle and as much as 600 N anteriorly shearing when the knee was straight.




Fig. 8. a) Patellar tendon force (F_p), b) tibio-femoral compressive force (F_{ct}) and, c) tibio-femoral shear force (F_s) for women (unfilled symbols) and men (filled circles) during isometric knee extension at various knee angles (x-axis). The external force (F_{et}) and its varying moment arm (d_e) are indicated. Note in a) and b) that the same magnitudes of F_{et} and d_e give higher forces for women than for men. Note in c) that the intersection of the F_s zero line is between 50 and 90 deg knee angle. Positive values in c) indicate that tibia tends to shear anteriorly in relation to femur, negative values – posteriorly shearing.

The magnitudes of the patello-femoral compressive force (F_{cp}) and tibio-femoral shear force (F_s) were determined during knee extension against gravity with and without a 31 N boot (Fig. 9). F_{cp} reached its maximum at 50 deg knee angle, and at straighter knee angles with higher extending muscle moment this force declined. The anteriorly directed F_s reached its maximum during the last few degrees of knee extension.




Fig. 9. Calculated patello-femoral joint compressive force (F_{cp}) and tibio-femoral shear force (F_s) during knee extension from 90 to zero deg (x-axis). a) Without boot, b) With a 31-N-boot.

The patellar forces may be calculated using equations (2), (7), (8), and (10). In Fig. 10c it is seen that the magnitude of the patello-femoral compressive force (F_{cp}) reached its maximum at 90 deg knee angle and decreased slightly towards 120 deg. The patellar tendon force (F_p) was lower than the quadriceps tendon force (F_q) at knee flexion angles of 60 deg to 120 deg and the minimum F_p/F_q ratio was around 0.70. A ratio of about 0.70 was also seen when comparing the patellar tendon thickness with that of the quadriceps tendon. The magnitude of the compressive force between the quadriceps tendon and the femoral intercondylar groove (F_{cq}) increased considerably at knee flexion angles more than 60 deg.

Comments. Note that the patellar forces can be quantified by using the force prediction diagram (Fig. 10) if three variables are known; 1) sex, 2) knee moment and, 3) knee flexion angle.

Quadriceps rupture (Study V)

The forces in the knee joint at the situation of complete bilateral quadriceps muscle rupture were calculated. The force in the quadriceps tendon (F_q) was greater than the other forces calculated, and its magnitude was between 10,900 N and 18,300 N. The flexing knee load moment ranged between 335 Nm and 550 Nm while the maximum flexing moments about the hip (410–670 Nm) and the lumbo-sacral joint (575–1100 Nm) were even greater.




Fig. 10. Prediction diagram for quantifying forces in a) patellar tendon, b) quadriceps tendon, c) patello-femoral joint and d) between quadriceps tendon and intercondylar groove at various knee flexion angles (x-axis) for women (unfilled circles) and men (filled circles). 95 % confidence intervals of the means are indicated. Let us look at one example of how F_{cp} can be determined. Consider a load moment of 210 Nm and a knee angle of 50 deg (as during jogging, Winter 1983). Enter Fig. 10c at the horizontal axis at 50 deg, continue vertically to the lower curve (male subjects), then horizontally to the vertical axis and read "24 N", which is multiplied by the moment value "210". This gives the force magnitude 5040 N, which is above 7 times BW for a 72 kg man. The other patellar forces can be calculated in a corresponding way. Reprinted with kind permission from the Scandinavian Journal of Rehabilitation Medicine (Study IV).

Isokinetic knee extension (Study VI)

At 180 deg/sec, the maximum moments produced by the quadriceps muscle reached mean peak magnitudes of 181 Nm at 65 deg knee flexion angle with the resistance pad placed distally. With the pad placed proximally, the peak moment occurred at the same knee flexion angle but was significantly lower (163 Nm). At 30 deg/sec, the mean magnitude of the peak moment was significantly higher (284 Nm) than at the faster speed, and the peak occurred somewhat earlier in the knee extension movement, at 70 deg knee flexion angle.

The tibio-femoral shear force (Fs) changed direction from being negative (tibia tends to move posteriorly in relation to femur) at the starting position, to a positive value when the knee was extended. Having the resistance pad proximal gave significantly lower anteriorly-directed shear forces but, on the other hand, higher posteriorly-directed shear forces. Knee extension of 30 deg/sec induced the highest anteriorly-directed shear force, with maximum mean magnitudes close to 1 BW (= 710 N), seen between 60 and 40 deg knee flexion angle.

Comments. Johnson (1982) noted also that the maximum moments induced by the knee extensor muscles become lower when the resistance pad is placed proximally on the leg. The reason for this difference is obscure but may possibly be an unfamiliar or uncomfortable feeling experienced when the pad is positioned proximally. The higher resistance forces may induce pain or other inhibiting influences from the cutaneous, periosteal and/or ligamentous mechanoreceptor afferents, which reflexly inhibit the activation of the motoneurons (Ekholm et al. 1960). Another explanation would be that the higher resistance force influences the tibial position in relation to femur (the proximal end of the tibia is kept more backwards), making the patellar tendon moment arm shorter. If this is true, the same maximum forces exerted by the quadriceps muscle and patellar tendon would produce *lower* knee extending moments due to an impaired mechanical geometry.

Discussion

Assumptions of the knee model

The two-dimensional local biomechanical knee model was made possible by using dissection and radiography in combination, and applying the anatomical findings to biomechanical principles. However, for a complete picture of the forces transmitted in the knee, a three-dimensional force analysis would be needed. Furthermore, the present model did not take into account any antagonistic muscle activity neither from hamstrings nor gastrocnemius muscles. In Studies I and II it was seen that the hamstrings were activated together with quadriceps action. The same has been shown during other knee extending activities when the foot is "fixed", either to the ground as in standing positions or in a closed chain as in bicycling (e.g. Carlsöö & Molbeck 1966, Kelley et al. 1978, Schüldt et al. 1983, Németh et al. 1984, Gregor et al. 1985, Ericson et al. 1985a). However, during knee extension with the foot not fixed – as during isokinetic knee extension – antagonistic muscle activity seems to be low (Osternig et al. 1984).

If the antagonistic muscle forces had been considered, the calculated joint force magnitudes would have been even higher. *However, there are no biomechanical models that consider antagonistic muscles, since the major problem is to quantify their tendon forces.* With electromyography it is possible to study and quantify the activity of such antagonistic muscles but difficult to quantify the tendon force magnitudes because the

relation between muscle activity and tendon force is not always linear (Jonsson 1978, Woods & Bigland-Ritchie 1983).

Using a permutation technique, Nissan (1980) showed the sensitivity of several variables in knee joint biomechanics. He concluded that accurate anteroposterior localisation of the contact point between tibia and femur is most important in the analysis of the knee joint. If the tibio-femoral contact point were moved backward 10 mm in our model, the calculated force in the patellar tendon would decrease by a maximum of approximately 22 %. If the contact point were moved 10 mm anteriorly this force would increase up to 40 %. Since tibio-femoral contact changes with knee angle, especially towards the end of knee extension, *it is important to identify the location of the contact point accurately in order to get proper data of the knee extensor moment arm and, thus, of the forces calculated.*

In the present study, the moment point chosen was the centre point of contact (C) between the condyles. It is a mathematical advantage to choose the contact point as the moment point because the compressive force acts through it and thus gives rise to a zero moment. The shear force is balanced by structures with short or zero moment arms to the contact point (such as cruciates, menisci, collateral ligaments and joint capsule) and therefore it causes a small or zero moment about this point. *In models that use the centre of the axis as the moment point (e.g. Smidt 1973), the moments caused by the joint compressive and shear forces have been omitted, which may give incorrect values of the force magnitudes calculated.*

With regard to the assumptions discussed above, the knee biomechanical model is considered as a useful instrument for quantifying knee joint forces. Comparing results from the biomechanical model with experimentally measured forces in the anterior cruciate ligament (Paulos et al. 1981, Arms et al. 1984), and in the quadriceps tendon (Grood et al. 1984), gives similar patterns and magnitudes of the force curves. In addition, experimentally measured relations between patellar (F_p) and quadriceps tendon forces (F_q) (Haxton 1945, Bishop & Denham 1977, Ellis et al 1980, Huberti et al. 1984) correlate well with the relation found from the present model. Furthermore, experimentally measured patello-femoral compressive forces (Ahmed et al. 1983, Huberti & Hayes 1984) have magnitudes and change patterns similar to the results from the present model. *These facts indicate that the model should be valid.*

Compressive force per area

When considering the magnitude of the tibio-femoral compressive force (F_{ct}), it should not be forgotten that this force acts through an area which varies with knee angle (Kettelkamp & Jacobs 1972, Maquet et al. 1975, Seedhom & Hargreaves 1979, Fukubayashi & Kurosawa 1980, Ahmed & Burke 1983). The tibio-femoral contact area is larger at straighter knee angles, indicating that tibio-femoral pressure is lower for straighter knees for the same F_{ct} magnitude. In Fig. 8 it is seen that F_{ct} reaches its maximum magnitude when the knee is straight but the tibio-femoral pressure is not highest in this position. Using the contact area magnitude found by Ahmed & Burke (1983), the area is about 50 % higher at zero deg than at 90 deg knee flexion angle. Thus, the tibio-femoral pressure calculated with the present biomechanical knee model (Fig. 10) should for the straight knee be about 25 % *lower* than for the 90-deg-flexed, although the compressive force is somewhat higher.

The same discussion applies to the patello-femoral joint since, here also, the contact area varies with knee joint angle. The largest area is reported at 60–90 deg knee flexion

angle and the smallest when the knee is straight (Matthews et al. 1977, Ficat & Hungerford 1977, Huberti & Hayes 1984). From this follows that *the patello-femoral pressure, also when the knee is almost straight, may be considerable although the compressive force seems to be comparatively low.*

Knee joint loads during various activities

In Table I the maximum knee load moments and patello-femoral compressive forces (Fcp) during various activities are presented. The knee moments were taken both from the literature and from Studies I, II, V and VI. Fcp was calculated using the "force prediction diagram" presented in Fig. 10 since the three variables sex, knee moment and knee angle were known. During level walking the flexing knee load moment ranges between 35 and 61 Nm and Fcp is not above 850 N. During jogging, descending stairs, kicking or rising from a chair, the knee moment is considerably higher and the knee flexion angle more pronounced which induce much higher Fcp magnitudes.

TABLE I. Maximum patello-femoral compressive force (Fcp) and knee extensor moments (Mk) produced by quadriceps muscle during various activities. The number of female subjects in the different experiments is given in brackets. Fcp was calculated using the force diagram presented in Fig. 10 and, as there is a difference in this diagram between men and woman, the sexes were considered.

Author(s)	Activity	Number of subjects	Mean weight (kg)	Knee angle (deg)	Mk (Nm)	Fcp (N)
Bresler & Frankel 1950	Level walking	4	71	20	60	840
Radcliffe 1962	Level walking	4	—	20	61	850
Morrison 1968	Level walking	1	—	—	35*	490
Boccardi et al. 1981	Level walking	1	65	—	60*	840
Winter 1980	Level walking	1(1)	—	—	45*	830
Winter 1983	Jogging	11	72	50	210	5000
Andriacchi et al. 1980	Ascending stairs	10	71	65	54	1500
Andriacchi et al. 1980	Descending stairs	10	71	60	147	4000
Lindahl et al. 1969	Isometric max	12	—	60	225	6100
Smidt 1973	Isometric max	26	82	60	120	3400
Wahrenberg et al. 1978	Kicking	6	76	100	180	5800
Kelley et al. 1978	Rising from chair	6(3)	—	90	110	3800
Schüldt et al. 1983	Rising from squat	9(5)	63	105	70	2500
Ericson et al. 1985b	Bicycling	6	71	29	79	880
Study I	Lifting	7	77	90	50	1600
Study II	Machine milking	10(4)	72	108	31	1100
Study II	Isometric max	10(4)	72	90	198	6900
Study V	Parallel squat	3	146**	120	455	14900
Study VI	Isokinetic knee ext	8	72	70	284	8300

* Knee flexion angle of 20 deg was used

** The body weight was simulated




Fig. 11. Knee load moment (y-axis, Nm) for one subject (72 kg) at various knee angles (x-axis, deg) and at various horizontal distances between hand held equipment (3.3 kg) and ankle joint (z-axis, m). The illustrations show the body positions at the four "corners". The zero knee load moment is the intersection of the diagram at the x-z plane.

Work postures

The use of the sagittal plane body model presented makes it possible to quantify knee joint loads in terms of load moments and also to compare different work postures or working tasks with each other. It is important to remember that to find the "best" work posture is to minimize the loading effect in a whole-body perspective and therefore all major joints must be considered. Studies I and II have focused only on the knee joint and the optimum working situation – with respect only to the knee joint – is found when the knee moment and muscle activity are low, which give low compressive and tensile forces in and around the joint. The lowest knee load calculated during machine milking (for the standing low level difference postures) occurred when the knees were flexed to about 65 degrees. In this position the knee load moment was around zero.

The sagittal knee load moment – in static or slow motioned standing activities such as those studied in the present work – is dependent on four parameters; 1) *body weight*, 2) *knee angle*, 3) *horizontal distance* to the working area, 4) *burden weight*. Fig. 11 shows the loading effects of changing two of these parameters, namely knee angle and horizontal distance. The most extending knee load moment occurs when the knees are straight and the horizontal distance to the working area is pronounced (top left in Fig. 11). If the hands with the burden move closer to the body (and the knees are still straight) the extending knee moment decreases (lower left). If the knees start bending

from this position the knee load changes to a flexing moment, which is maximum when the burden is close to the body and the knees most flexed. The flexing knee load moment decreases if the burden moves away from the trunk (to the very right) and changes to an extending moment when knee straightening occurs back to the top left position.

Thus, it is possible to determine the sagittal knee load moment when the parameters described here are known. The same principles can be used for determining knee moments in other similar standing work situations. In most kinds of lifting work and in many kinds of repair work this model can be generally used in order to find the least load on the knee joints.

Clinical implications

The pattern of change in patellar tendon moment arm (Fig. 7) was quite similar to that in earlier reports (Haxton 1945, Lindahl & Movin 1967, Kaufer 1971, Bandi 1972, Smidt 1973). The main difference between the present results and others is seen at straighter knee angles, where the moment arm was suggested to be about 10 mm shorter than what has been reported earlier. This fact, together with the physiological muscular length-tension relationship (Wilkie 1956, Lunnen et al. 1981), may be one explanation why knee extending strength decreases during the last 30–40 deg of knee extension (Williams & Stutzman 1959, Hallén & Lindahl 1967, Lieb & Perry 1968 and 1971, Lindahl & Movin 1968, Lindahl et al. 1969, Moritz et al. 1973, Smidt 1973). *It may also explain the difficulty, often seen in clinical practice, in achieving the last few degrees of knee extension.*

Due to differences between men and women in length of the patellar tendon moment arm (Fig. 7), the female hyaline cartilage will be exposed to higher forces if the extending muscular moment is of the same magnitude. In addition, the contact area over which the compressive force acts, is smaller in women than in men due to smaller knees, indicating that the women knee joint pressure will be even higher. *This may be one explanation, among others, why gonarthrosis is more frequent among women than men (Danielsson & Hernborg 1970, Hernborg & Nilsson 1973).* The sex anatomical differences of the hip extensor moment arms are considerably smaller than that of the knee extensors, and therefore the hip force magnitudes in women and men are of the same magnitude (Németh & Ohlsén 1985). *This might mechanically explain why there is no sex difference in the prevalence of hip joint arthrosis but there is in the case of the knee (Danielsson 1966, Lindberg 1985).*

Using the biomechanical model the situation after patellectomy can be assessed for various knee flexion angles. An example of this is illustrated in Fig. 12 at 30 deg knee angle. The moment arm (dp) decreased 15 % which means that the knee extensor strength will decrease 15 % at this particular knee flexion angle. The patellar tendon angle (β) decreased from 21 to 5 deg, indicating that tibia will shear less in an anterior direction. The biomechanical effects of anteversion of the tuberositas tibiae can be assessed in a similar way.

During knee extension exercises at knee flexion angles straighter than 30–40 deg, there is a considerable anteriorly directed tibio-femoral shear force which gives rise to high forces in the anterior cruciate ligament (Lindahl & Movin 1967, Paulos et al. 1981, Arms et al. 1984, Noyes et al. 1984). Butler et al. (1980) showed that the anterior cruciate takes up 86 % of the anterior shear force. By means of the biomechanical model presented, such forces may be estimated and consequently, different adjust-




Fig. 12. Biomechanical effects of simulated patellectomy. Note the decrease of patellar tendon moment arm length (dp) and patellar tendon angle (β).

ments of knee extension exercises may be assessed with regard to *e.g.* anterior cruciate ligament stress. Such an example is given for isokinetic knee extension (Study VI), where it was concluded that having the resistance pad placed proximally on the anterior part of the lower leg considerably decreased the shear force and consequently the stress on the anterior cruciate ligament. Similar conclusions were found by Johnson (1982) when using a dual pad resistance and in a recent study by Jurist & Otis (1985). *Consequently, patients with repairs to or reconstructions of the anterior cruciate ligament should be advised to start their quadriceps exercise carefully at straighter knee angles and with the resistance force proximally on the lower leg.*

Over the last few degrees during knee extension against gravity (Fig. 9), the anteriorly directed shear force reached its maximum at the same time as the patello-femoral compressive force was low. The peak compressive force occurred at 50 deg knee flexion angle and now the shear force was low. However, during isokinetic knee extension, the shear force was also high at knee angles as flexed as 60 deg (Fig. 12b). Consequently, it is of interest to know and consider the internal force distribution during different quadriceps exercises in patients with different diagnoses such as patello-femoral arthrosis or anterior cruciate ligament deficiency.

The force between the quadriceps tendon and femoral intercondylar groove (F_{cq}) was found to be considerable, especially when the knee was flexed to more than 90 deg (Fig. 10d). *It cannot be excluded that this force might compress the upper part of the suprapatellar bursa, and in some cases even cause pain.* This tendon-bone compression has been discussed earlier (Bishop 1977, Goodfellow et al. 1976, Goyman & Müller 1974, Grood et al. 1984), and Huberti & Hayes (1984) measured its magnitude using pressure sensitive film. However, its quantification in a biomechanical model has not earlier been performed.

A painful patello-femoral joint sensitive to high stresses (*e.g.* oosteoarthritis) is better exercised at straighter knee angles so that high flexing load moments and high compressive forces are avoided. Note that the force magnitudes received by the patello-femoral joint are about the same between 60 deg and 120 deg, so for patients with pain elicited from this joint there should be no reason to avoid 120 deg more than 60 deg. Exercises aiming at an optimal range of motion are of course also necessary, but should be undertaken mainly passively, *i.e.* with no or very little load. Patients with load-elicited pain from the patello-femoral joint might be advised to avoid knee angles above 30 deg under loaded conditions such as in deep squatting, lifting objects from the floor with flexed knees, ascending or descending stairs. The knee extensor apparatus is less activated at low knee angles and the patello-femoral compression is low. *A patello-femoral arthrosis patient might even be advised to lift with straight knees and flexed trunk* – if this does not cause low back pain – or to use technical aids.

Conclusions

The knee load moments and muscle activities during lifting and machine milking were quantified and a local knee biomechanical model presented. It was shown how to use the knee model in order to quantify the tibio-femoral and patello-femoral forces during ascending from the parallel squat in powerlifting and during isokinetic knee extension. The model may also be used to quantify knee joint forces during other knee extending activities and thus, to assess different work postures, training exercises and joint derangements.

The following main conclusions can be stated:

- Lifting a 12.8 kg box with flexed knees, from floor to table level, gave a maximum flexing knee load moment of about 50 Nm at the beginning of the lift, with a knee flexion angle of 90 deg. The quadriceps muscle was highly activated in order to counteract the load moment.
- No significant difference in knee joint load was seen with flexed knees during lifting the box from between or in front of the feet.
- Straight-knee lifts gave an extending knee load moment throughout the lift. During the final phase of all three lifts (SK, FKFF and FKC) there was an extending moment of about 55 Nm.
- During machine milking, the horizontal distance between the operator and the cow was found to be of considerable importance for the magnitude of the extending knee load moment in the straight-knee standing postures. The extending load moment decreased when the milker came closer to the cow and to the working area. Thus the horizontal distance to the cow should be considered when designing milking equipment.
- In flexed-knee postures, normalized EMG-activity in the quadriceps muscle was higher than that in the hamstrings muscles in straight-knee postures.
- The knee load moment in the standing flexed-knee postures was in the direction of flexion and comparatively low, while in the straight-knee postures it was in the direction of extension and comparatively high.

- In a static situation, with a defined hand-held implement and a horizontal distance to the working area, it is possible to determine what knee angle gives the least, or even zero, load moment on the knees. Flexing the knees gave a reduction of the extending load moment in the postures with small level differences between operator and cow. A knee angle of around 65 deg gave a zero load moment for implement weight and posture about the same as in Study II. This knee angle was not influenced by body weight.
- Women were found to have shorter patellar tendon moment arms than men, due to smaller knees, and thus develop about 20 % higher knee joint forces for the same knee extending moment.
- The angle between the tibial plateau and patellar tendon force changed considerably with knee flexion angle. Therefore, the anteroposterior tibio-femoral shear force was found to change direction from negative, with proximal tibia sliding posteriorly in relation to distal femur, to positive during the extension movement, indicating that high forces may arise in the anterior cruciate ligament when the knee is straight.
- The patellar tendon was found to be 25–30 % thinner and narrower than the quadriceps tendon, and the forces in these tendons at knee angles of 60–120 deg had a relation (F_p/F_q) of about 0.70–0.80. At straighter knee angles, the two forces were of the same magnitude.
- A prediction diagram was designed and used for quantifying patellar forces in women and men for given moment magnitude and knee angle.
- For a constant knee extending moment, the patello-femoral joint compressive force reached a maximum at 90 deg knee angle and decreased in magnitude beyond this knee angle.
- Above knee angles of 60 deg, the compressive force between the femoral intercondylar groove and the quadriceps tendon was quantified and found to increase linearly with increasing knee angle.
- The force magnitudes calculated using the knee joint biomechanical model developed, correlated well with forces measured experimentally by others, indicating that the model is valid.
- Performing a parallel squat while carrying a 382.5 kg weight induced a knee flexing load moment of 335–550 Nm. The *in vivo* force of a complete quadriceps tendon-muscle rupture was estimated to between 10,900 and 18,300 N.
- During isokinetic knee extension at slower speed (30 deg/sec), the patellar tendon force and tibio-femoral compressive force magnitudes peaked to almost 9 times BW and during the faster speed (180 deg/sec), to 5 times BW.
- During isokinetic knee extension at 30 deg/sec with the resistance pad distally, the anteriorly directed tibio-femoral shear force was found to be of considerable magnitude between 30 and 60 deg knee flexion angles. A proximal position of the resistance pad decreased the tibio-femoral shear force significantly, indicating that the anterior cruciate ligament receives lower stresses. This position of the pad was recommended for early postsurgical isokinetic rehabilitation of anterior cruciate ligament repairs and reconstructions.

Acknowledgements

This thesis investigation was carried out at the Department of Anatomy at Karolinska Institute in Stockholm. During the work I have had the privilege and pleasure of cooperating with some outstanding people, who have given me not only my proficiency in science but also valuable knowledge of human behaviour and of life in general. I should like to express my sincere appreciation and gratitude to:

Professor GUNNAR GRANT and Professor LARS-GÖSTA ELFVIN for making the research facilities at the Department of Anatomy available to me;

Professor JAN EKHOLM, my tutor, former Senior Lecturer at the Department of Anatomy and now at the Department of Physical Medicine & Rehabilitation, Karolinska Hospital, Head of the Kinesiology Research Group, for introducing me to kinesiology and biomechanics, for great generosity and support and for his open and non-authoritarian attitude which has done much to generate a congenial and inspiring research atmosphere;

ULF ARBORELIUS, MB, for introducing me to kinesiology, biomechanics and the world of computers, for his friendship and for many stimulating discussions;

Dr GUNNAR NÉMETH, for showing me "how to do it", for constructive criticism and permanent practical help throughout the work;

Dr OLA SVENSSON, for fruitful cooperation and friendship, and for many interesting and stimulating discussions;

MATS ERICSON, MB, for his enthusiasm, creativity and close friendship, and for innumerable inspiring discussions;

Physiotherapist KARIN HARMS-RINGDAHL and Dr TINA SCHÜLDT, the other members of the Kinesiology Research Group, for stimulating cooperation and friendship;

Miss CHRISTINE INGEMANSSON, for invaluable help with invoices and applications, and for her excellent coordination of the Kinesiology Research Group;

Dr LARS LINDBECK, National Board of Occupational Safety and Health, for fruitful discussions and expert help regarding the biomechanical analyses;

Dr HANS OHLSÉN, Department of Diagnostic Radiology, Karolinska Hospital, co-writer in Study III, for expert help with the radiographics;

TIM CROSFIELD, MA, for expert revision of the English;

LENNART WAHLSTRÖM, Alfa Laval, for professional help with illustrations in Study II;

INGE JOHANSSON, National coach at the Swedish Weightlifting Federation, for help in collecting subjects for Study V;

MARIANNE GOTTFRIDSSON, MARGIT JEPSEN and MARIANNE RAPP, for secretarial assistance; PIPPI LINDOVIST, for kindly introducing me how to make drawings; BRITT DAHLGREN, for help with illustrations in Studies V and VI; SOLVEIG SIÖCRONA and PETER HOLMBERG for photographic help;

All other friends at the Department of Anatomy, for general help and support;

My friends at the Department of Medical Information Processing, Karolinska Institute, for valuable, time-saving help in how to manage word processing and computer graphics, and especially thanks to ELISABETH BERG for professional statistical advice;

The staff at the Library of the Karolinska Institute, for invaluable help in finding literature and references;

MY PARENTS, for their interest, help and never-failing support.

Last but not least I thank my wife SUSANNE for her honest and sincere love, for her spirit and help, and for her seldom-failing support during these years.

Financial support was kindly given by: The Swedish Work Environment Fund (79/1340, 84/0520), Alfa Laval, the Swedish Medical Research Council (5720), and the Karolinska Institute.

References

1. **Ahmed AM & DL Burke:** In-vitro measurement of static pressure distribution in synovial joints – Part I: Tibial surface of the knee. *J Biomech Engin*, 105:216–25, 1983
2. **Ahmed AM, DL Burke & A Yu:** In-vitro measurement of static pressure distribution in synovial joints – Part II: Retropatellar surface. *J Biomech Engin*, 105:226–36, 1983
3. **Alexander RM & A Vernon:** The dimensions of knee and ankle muscles and the forces they exert. *J Human Movem Stud*, 1:115–23, 1975
4. **Andersson GBJ, R Örtengren & A Nachemson:** Quantitative studies of back loads in lifting. *Spine*, 1:178–85, 1976
5. **Andriacchi TP, GBJ Andersson, RW Fermier, D Stern & JO Galante:** A study of lower-limb mechanics during stair-climbing. *J Bone Joint Surg*, 62-A:749–57, 1980
6. **Andriacchi TP, GBJ Andersson, R Örtengren & RP Mikosz:** A study of factors influencing muscle activity about the knee joint. *J Orthop Res*, 1:266–75, 1984
7. **Arborelius UP & J Ekholm:** Mechanics of shoulder locomotor system during exercises resisted by weight-and-pulley-circuit. *Scand J Rehab Med*, 10:171–7, 1978
8. **Arms SW, MH Pope, RJ Johnson, RA Fischer, I Arvidsson & E Eriksson:** The biomechanics of the anterior cruciate ligament and reconstruction. *Am J Sports Med*, 12:8–18, 1984
9. **Arvidsson I, E Eriksson, T Häggmark & RJ Johnson:** Isokinetic thigh muscle strength after ligament reconstruction in the knee joint. Results from a 5–10 year follow up after reconstruction of the anterior cruciate ligament in the knee joint. *Int J Sports Med*, 2:7–11, 1981
10. **Bandi W:** Chondromalacia Patellae and Femoro-Patellare Arthrose. *Helv Chir Acta*, Suppl. 11, 1972
11. **Basmajian JV:** Muscles Alive. Their Function Revealed by Electromyography. Williams & Wilkins., Baltimore, 1962
12. **Bhole R, J Flynn & TC Marbury:** Quadriceps tendon ruptures in uremia. *Clin Orthop*, 195:200–6, 1985
13. **Bishop RED:** On the mechanics of the human knee. *Engineering in Medicine*, 6:46–52, 1977
14. **Bishop RED & RA Denham:** A note on the ratio between tensions in the quadriceps tendon and infra-patellar ligament. *Engineering in Medicine*, 6:53–4, 1977
15. **Bjernelid H, K Eriksson & G Hallin:** Styrkelyftning, doping och skador – analgetika tystar varningssignaler. *Läkartidningen*, 81:2151, 1984 (In Swedish)
16. **Boccardi S, A Pedotti, R Rodano & GC Santambrogio:** Evaluation of muscular moments at the lower limb joints by an on-line processing of kinematic data and ground reaction. *J Biomech*, 14:35–45, 1981
17. **Bouisset S:** EMG and muscle force in normal motor activities. In: JE Desmedt (ed), *New Developm in EMG and Clin Neurophys*, Karger, Basel, Vol 1:547–83, 1973
18. **Brask B, RH Lueke & GL Soderberg:** Electromyographic analysis of selected muscles during the lateral step-up exercise. *Physical Therapy*, 64:324–8, 1984
19. **Bresler B & JP Frankel:** The Forces and Moments in the Leg During Level Walking. *Transact ASME*, Jan:27–36, 1950
20. **Brotherton BJ & J Ball:** Bilateral simultaneous rupture of the quadriceps tendons. *Br J Surg*, 62:918–20, 1975
21. **Butler DL, FR Noyes & ES Grood:** Ligamentous restraints to anterior-posterior drawer in the human knee. *J Bone Joint Surg*, 62-A:259–70, 1980
22. **Campbell DE & W Glenn:** Foot-pounds of torque of the normal knee and the rehabilitated postmeniscectomy knee. *Phys Ther*, 59:418–21, 1979
23. **Carlsöö S & S Molbeck:** The functions of certain two-joint muscles in a closed muscular chain. *Acta Morph Neerl-Scand*, 6:377–86, 1966
24. **Carlsöö S:** How Man Moves. Heinemann, London, 1975
25. **Chaffin DB & GBJ Andersson:** Occupational Biomechanics. Wiley & Sons, New York, 1984
26. **Dahlqvist NJ, P Mayo & BB Seedhom:** Forces during squatting and rising from a deep squat. *Engineering in Medicine*, 11:69–76, 1982
27. **Dalal VD & DE Whittam:** Bilateral simultaneous rupture of the quadriceps tendons. *Brit Med J*, 2:1370, 1966

28. **Danielsson L & J Hernborg:** Morbidity and mortality of osteoarthritis of the knee (gonarthrosis) in Malmö, Sweden. *Clin Orthop*, 69:224–6, 1970
29. **Danielsson LG:** Incidence of osteoarthritis of the hip. *Clin Orthop*, 45:67–72, 1966
30. **Dempster WT:** Space requirements of the seated operator. WADC Technical report, (Wright Patterson Air Force Base, Ohio: Wright Air Development Center):55–159, 1955
31. **Dostal WF & JG Andrews:** A three-dimensional biomechanical model of hip musculature. *J Biomech*, 14:803–12, 1981
32. **Dowson D & V Wright:** An Introduction to the Biomechanics of Joints and Joint Replacement. Mechanical Engineering Publications LTD, London, 1981
33. **Ekholm J, G Eklund & S Skoglund:** On the reflex effects from the knee joint of the cat. *Acta Physiol Scand*, 50:167–74, 1960
34. **Ekholm J, U Arborelius, L Hillered & Å Örtqvist:** Shoulder muscle EMG and resisting moment during diagonal exercise movements resisted by weight-and-pulley-circuit. *Scand J Rehab Med*, 10:179–85, 1978
35. **Ekholm J, UP Arborelius & G Németh:** The load on the lumbo-sacral joint and trunk muscle activity during lifting. *Ergonomics*, 25:145–61, 1982
36. **Ellis MI, BB Seedhom, AA Amis, D Dowson & V Wright:** Forces in the knee joint whilst rising from normal and motorized chairs. *Engineering in Medicine*, 8:33–40, 1979
37. **Ellis MI, BB Seedhom, V Wright & D Dowson:** An evaluation of the ratio between the tensions along the quadriceps tendon and the patellar ligament. *Engineering in Medicine*, 9:189–94, 1980
38. **Ellis MI, BB Seedhom & V Wright:** Forces in the knee joint whilst rising from a seated position. *J Biomed Eng*, 6:113–20, 1984
39. **Ericson MO, R Nisell, U Arborelius & J Ekholm:** Muscular activity during ergometer cycling. *Scand J Rehab Med*, 17:53–61, 1985a
40. **Ericson MO, R Nisell, G Németh & J Ekholm:** Hip and knee load moments during ergometer cycling. *Scand J Rehab Med*, Accepted for publication, 1985b
41. **Fairbank TJ:** Knee joint changes after meniscectomy. *J Bone Joint Surg*, 30-B:664–70, 1948
42. **Ficat RP & DS Hungerford:** Disorders of the Patello-Femoral Joint. Masson SA, Williams & Wilkins, 1977
43. **Firooznia H, G Selliger, RM Abrams & M Sanz:** Bilateral spontaneous and simultaneous rupture of the quadriceps tendon. *Bull Hosp Joint Dis*, 34:65–9, 1973
44. **Frankel VH & AH Burstein:** Orthopaedic Biomechanics. Lea & Febiger, Philadelphia, 1971
45. **Frankel VH & M Nordin:** Basic Biomechanics of the Skeletal System. Lea & Febiger, Philadelphia, 1980
46. **Frey S:** Doppelseitige subkutane Quadricepssehnrupturn. *Deutsche Zeitschrift für Chirurgie*, 209:284–6, 1928
47. **Fugl-Meyer AR, L Gustafsson & Y Burstedt:** Isokinetic and static plantar flexion characteristics. *Eur J Appl Phys*, 45:221–34, 1980
48. **Fukubayashi T & H Kurosawa:** The contact area and pressure distribution pattern of the knee. A study of normal and osteoarthrotic knee joints. *Acta Orthop Scand*, 51:871–9, 1980
49. **Goldenberg RR & NJ Paterson:** Bilateral rupture of the quadriceps tendon. *J Med Soc New Jersey*, 46:349, 1949
50. **Goodfellow J, DS Hungerford & M Zindel:** Patello-femoral joint mechanics and pathology. 1. Functional anatomy of the patello-femoral Joint. *J Bone Joint Surg*, 58-B:287–90, 1976
51. **Goymann V & HG Müller:** New calculations of the biomechanics of the patellofemoral joint and its clinical significance. In: *The Knee Joint*, (ed Ingwersen et al.):16–21, 1974
52. **Gransberg L & E Knutsson:** Determination of dynamic muscle strength in man with acceleration controlled isokinetic movements. *Acta Phys Scand*, 119:317–20, 1983
53. **Gregor RJ, PR Cavanagh & M LaFortune:** Knee flexor moments during propulsion in cycling – a creative solution to Lombard's paradox. *J Biomech*, 18:307–16, 1985
54. **Grenier R & A Guimont:** Simultaneous bilateral rupture of the quadriceps tendon and leg fractures in a weightlifter. *Am J Sports Med*, 11:451–3, 1983
55. **Grimby G, E Gustafsson, L Peterson & P Renström:** Quadriceps function and training after knee ligament surgery. *Med Sci Sports*, 12:70–5, 1980
56. **Grood ES, WJ Suntay, FR Noyes & D Butler:** Biomechanics of the knee-extension exercise. *J Bone Joint Surg*, 66-A:725–34, 1984
57. **Haffajee D, U Moritz & G Svantesson:** Isometric knee extension strength as a function of joint angle, muscle length and motor unit activity. *Acta Orthop Scand*, 43:138–47, 1972
58. **Hagberg M:** The amplitude distribution of surface EMG in static and intermittent static muscular performance. *Eur J Appl Physiol*, 40:265–72, 1979
59. **Hallén LG & O Lindahl:** Muscle function in knee extension. *Acta Orthop Scand*, 38:434–44, 1967
60. **Hamberg P & J Gillquist:** Knee function after arthroscopic meniscectomy. *Acta Orthop Scand*, 55:172–5, 1984

61. **Harms-Ringdahl K, J Ekholm, U Arborelius, G Németh & K Schüldt:** Load moment, muscle strength and level of muscular activity during internal rotation training of the shoulder. *Scand J Rehab Med*, Suppl 9:125–35, 1983
62. **Haxton H:** The function of the patella and the effects of its excision. *Surg Gynecol Obstet*, 80:389–95, 1945
63. **Herberts P, R Kadefors & H Broman:** Arm positioning in manual tasks. An electromyographic study of localized muscle fatigue. *Ergonomics*, 23:655–65, 1980
64. **Hernborg J & BE Nilsson:** Age and sex incidence of osteophytes in the knee joint. *Acta Orthop Scand*, 44:66–8, 1973
65. **Hislop HJ & JJ Perrine:** The isokinetic concept of exercise. *Phys Ther*, 47:114–7, 1967
66. **Huberti HH & WC Hayes:** Patellofemoral contact pressures. *J Bone Joint Surg*, 66-A:715–24, 1984
67. **Huberti HH, WC Hayes, JL Stone & GT Shybut:** Force ratios in the quadriceps tendon and ligamentum patellae. *J Orthop Res*, 2:49–54, 1984
68. **Hungerford DS & DW Lennox:** Rehabilitation of the knee in disorders of the patellofemoral joint: relevant biomechanics. *Orthop Clin North Am*, 14:397–402, 1983
69. **Ingemann-Hansen T & J Halkjaer-Kristensen:** Force-velocity relationships in the human quadriceps muscles. *Scand J Rehab Med*, 11:85–9, 1979
70. **Jackson JP:** Degenerative changes in the knee after meniscectomy. *Br Med J*, 2:525–7, 1968
71. **Jensen RH & DT Davy:** An investigation of muscle lines of action about the hip: A centroid line approach vs the straight line approach. *J Biomech*, 8:103–10, 1975
72. **Johnson D:** Controlling anterior shear during isokinetic knee extension. *J Orthop & Sports Phys Ther*, 4:23–31, 1982
73. **Johnson RJ, DB Kettelkamp, W Clark & P Leaverton:** Factors affecting late results after meniscectomy. *J Bone Joint Surg*, 56-A:719–29, 1974
74. **Jonsson B & B Synnerstad:** Electromyographic studies of muscle function in standing. A methodological study. *Acta Morph Neerl-Scand*, 6:361–70, 1966
75. **Jonsson B:** Kinesiology. With special reference to electromyographic kinesiology. In: WB Cobb & H VanDuijn (eds), *Contemp Clin Neurophysiol*, Elsevier Scientific Publishing Co, Amsterdam, Suppl 34:417–28, 1978
76. **Jurist KA & JC Otis:** Anteroposterior tibiofemoral displacements during isometric extension efforts. *Am J Sports Med*, 13:254–8, 1985
77. **Kadefors R, E Kaiser & I Petersén:** Dynamic spectrum analysis of myopotentials with special reference to muscle fatigue. *Electromyogr*, 8:39–74, 1968
78. **Kadefors R:** Application of electromyography in ergonomics: new vistas. *Scand J Rehab Med*, 10:127–33, 1978
79. **Kauffer H:** Mechanical function of the patella. *J Bone Joint Surg*, 53-A:1551–60, 1971
80. **Kelley DL, A Dainis & GK Wood:** Mechanics and muscular dynamics of rising from a seated position. In *Biomechanics V*, (ed PV Komi):127–34, 1978
81. **Kettelkamp DB & AW Jacobs:** Tibio-femoral contact area – determination and implications. *J Bone Joint Surg*, 54-A:349–56, 1972
82. **Knutsson E & A Mårtensson:** Dynamic motor capacity in spastic paresis and its relation to prime mover dysfunction, spastic reflexes and antagonistic co-activation. *Scand J Rehab Med*, 12:93–106, 1980
83. **Komi PV & P Tesch:** EMG frequency spectrum, muscle structure, and fatigue during dynamic contractions in man. *Eur J Appl Phys*, 42:41–50, 1979
84. **LeVeau B:** Williams and Lissner: *Biomechanics of Human motion*. WB Saunders Company, 2:nd edition, 1977
85. **LeVeau BF:** Biomechanics. A Summary of Perspectives. *Phys Ther*, 64:1812, 1984
86. **Levy M, M Seelenfreund, P Maor, A Fried & M Lurie:** Bilateral spontaneous and simultaneous rupture of the quadriceps tendons in gout. *J Bone Joint Surg*, 53-B:510–3, 1971
87. **Lieb FJ & J Perry:** Quadriceps function. An anatomical and mechanical study using amputated limbs. *J Bone Joint Surg*, 50-A:1535–48, 1968
88. **Lieb FJ & J Perry:** Quadriceps function. An electromyographic study under isometric conditions. *J Bone Joint Surg*, 53-A:749–58, 1971
89. **Lindahl O & A Movin:** The mechanics of extension of the knee joint. *Acta Orthop Scand*, 38:226–34, 1967
90. **Lindahl O & A Movin:** Active extension capacity of the knee joint in the healthy subject. *Acta Orthop Scand*, 39:203–8, 1968
91. **Lindahl O, A Movin & I Ringqvist:** Knee extension. *Acta Orthop Scand*, 40:79–85, 1969
92. **Lindbeck L:** Impulse and moment of impulse in the leg joints by impact from kicking. *J Biomech Engin*, 105:108–11, 1983
93. **Lindberg H:** Epidemiological Studies on Primary Coxarthrosis. Dept of Orthop Surg, Malmö General Hospital, University of Lund, Thesis, 1985

94. **Lotem M, J Bernheim & B Conforty:** Spontaneous rupture of tendons. A complication of hemodialyzed patients treated for renal failure. *Nephron*, 21:201–8, 1978
95. **Lunnen JD, J Yack & BF LeVeau:** Relationship between muscle length, muscle activity, and torque of the hamstring muscles. *Phys Ther*, 61:190–5, 1981
96. **Lysholm J, M Nordin, J Ekstrand & J Gillquist:** The effect of a patella brace on performance in a knee extension strength test in patients with patellar pain. *Am J Sports Med*, 12:110–2, 1984
97. **MacDonald J:** Bilateral subcutaneous rupture of the quadriceps tendon: Report of a case with delayed repair. *Can J Surg*, 9:74–6, 1966
98. **MacEachern AG & JL Plewis:** Bilateral simultaneous spontaneous rupture of the quadriceps tendons. *J Bone Joint Surg*, 66-B:81–3, 1984
99. **Maquet P, AJ Van de Berg & JC Simonet:** Femorotibial weight-bearing areas. *J Bone Joint Surg*, 57-A:766–71, 1975
100. **Maquet P:** Biomechanics of the Knee. Springer Verlag, Berlin-Heidelberg-New York, 1984
101. **Markhede G & L Nistor:** Strength of plantar flexion and function after resection of various parts of the triceps surae muscle. *Acta Orthop Scand*, 50:693–7, 1979
102. **Matthews LS, DA Sonstegard & JA Henke:** Load bearing characteristics of the patello-femoral joint. *Acta Orthop Scand*, 48:511–6, 1977
103. **Mira AJ, CK Markley & RB Greer:** A critical analysis of quadriceps function after femoral shaft fracture in adults. *J Bone Joint Surg*, 62-A:61–7, 1980
104. **Morein G, Z Goldschmidt, M Pauker, M Seelenfreund, JB Rosenfeld & A Fried:** Spontaneous tendon ruptures in patients treated by chronic hemodialysis. *Clin Orthop*, 124:209–13, 1977
105. **Moritz U, G Svantesson & D Haffajee:** A biomechanical study of muscle torque as affected by motor unit activity, length-tension relationship and muscle force lever. In: JE Desmedt (ed), *New Developm in EMG and Clin Neuropphys*, Karger, Basel, Vol 1:675–82, 1973
106. **Morrison JB:** Bioengineering analysis of force actions transmitted by the knee joint. *Bio-Med Engin*, 3:164–70, 1968
107. **Morrison JB:** Function of the knee joint in various activities. *Bio-Med Engin*, 4:573–80, 1969
108. **Morrison JB:** The mechanics of the knee joint in relation to normal walking. *J Biomech*, 3:51–61, 1970
109. **Murray SM, RF Warren, JC Otis, M Kroll & TL Wickiewicz:** Torque-velocity relationships of the knee extensor and flexor muscles in individuals sustaining injuries of the anterior cruciate ligament. *Am J Sports Med*, 12:436–40, 1984
110. **Möller E:** The chewing apparatus. An electromyographic study of the action of the muscles of mastication and its correlation to facial morphology. *Acta Physiol Scand*, 69, Suppl 280, 1966
111. **Nisell R, J Mizrahi & J Ekholm:** Knee and ankle loads during step down activities. 10:th International Congress of Biomechanics, Umeå, June 15–20, Proceedings:195, 1985
112. **Nissan M:** Review of some basic assumptions in knee biomechanics. *J Biomech*, 13:375–81, 1980
113. **Nistor L, GBJ Andersson & R Örtengren:** Studies of the function of the triceps surae after surgical and non-surgical treatment of achilles tendon rupture. *Orthop Trans*, 3:198–9, 1979
114. **Nordesjö L-O & B Nordgren:** Static and dynamic measuring of muscle function. *Scand J Rehab Med*, Suppl 6:33–42, 1978
115. **Nordgren B, L-O Nordesjö & W Rauschnig:** Isokinetic knee extension strength and pain before and after advancement osteotomy of the tibial tuberosity. *Acta Orthop Trauma Surg*, 102:95–101, 1983
116. **Nordin M:** Methods for studying work load with special reference to the lumbar spine. Dept of Orthop Surg I, Sahlgren Hospital, University of Gothenburg, Thesis, 1982
117. **Noyes FR, CS Keller, ES Grood & DL Butler:** Advances in the understanding of knee ligament injury, repair, and rehabilitation. *Med Sci Sports*, 16:427–43, 1984
118. **Németh G, J Ekholm, UP Arborelius, K Schöldt & K Harms-Ringdahl:** Hip joint load and muscular activation during rising exercises. *Scand J Rehab Med*, 16:93–102, 1984
119. **Németh G:** On Hip and Lumbar Biomechanics: A study of joint load and muscular activity. *Scand J Rehab Med*, Suppl 10, 1984
120. **Németh G & H Ohlsén:** In vivo moment arm lengths for hip extensor muscles at different angles of hip flexion. *J Biomech*, 18:129–40, 1985
121. **Osternig LR, J Hamill, DM Corcos & J Lander:** Electromyographic patterns accompanying isokinetic exercise under varying speed and sequencing conditions. *Am J Phys Med*, 63:289–97, 1984
122. **Paulos L, FR Noyes, E Grood & DL Butler:** Knee rehabilitation after anterior cruciate ligament reconstruction and repair. *Am J Sports Med*, 9:140–9, 1981
123. **Peiro A, R Ferrandis, L Garcia & E Alcazar:** Simultaneous spontaneous bilateral rupture of the patellar tendon in rheumatoid arthritis. *Acta Orthop Scand*, 46:700–3, 1975
124. **Perrine JJ & VR Edgerton:** Muscle force-velocity and power-velocity relationships under isokinetic loading. *Med Sci Sports*, 10:159–66, 1978
125. **Perry J, D Antonelli & W Ford:** Analysis of knee-joint forces during flexed-knee stance. *J Bone Joint Surg*, 57-A:961–7, 1975

126. **Petrofsky JS:** Frequency and amplitude analysis of the EMG during exercise on the bicycle ergometer. *Eur J Appl Physiol*, 41:1-5, 1979
127. **Phillips CA & JS Petrofsky:** Quantitative electromyography: Response of the neck muscles to conventional helmet loading. *Aviation, Space, and Environment Medicine*, 57:452-7, 1983
128. **Pipes TV & JH Wilmore:** Isokinetic vs isotonic strength training in adult men. *Med Sci Sports*, 4:262-74, 1975
129. **Pocock G:** Electromyographic study of the quadriceps during resistive exercise. *J Am Phys Ther Assoc*, 43:427-34, 1963
130. **Preston FS & A Adicoff:** Hyperparathyroidism with avulsion of three major tendons. *New Engl J Med*, 266:968-71, 1962
131. **Radcliffe CW:** The biomechanics of below-knee prostheses in normal, level, bipedal walking. *Artificial Limbs*, 6:16-24, 1962
132. **Rauschnig W, L-O Nordesjö & B Nordgren:** Isokinetic knee extension strength and pain before and after correction of recurrent patellar dislocation. *Acta Orthop Trauma Surg*, 102:102-6, 1983
133. **Reilly DT & M Martens:** Experimental analysis of the quadriceps muscle force and patello-femoral joint reaction force for various activities. *Acta Orthop Scand*, 43:126-37, 1972
134. **Reynolds L, TA Levin, JM Medeiros, NS Adler & A Hallum:** EMG activity of the vastus medialis oblique and the vastus lateralis in their role in patellar alignment. *Am J Phys Med*, 62:61-70, 1983
135. **Schüldt K, J Ekholm, G Németh, UP Arborelius & K Harms-Ringdahl:** Knee load and muscle activity during exercises in rising. *Scand J Rehab Med, Suppl.* 9:174-88, 1983
136. **Seedhom BB & K Terayama:** Knee forces during the activity of getting out of a chair with and without the aid of arms. *Bio-Med Engin*, Aug:278-82, 1976
137. **Seedhom BB:** Transmission of the load in the knee joint with special reference to the role of the menisci. Part I: anatomy, analysis and apparatus. *Engineering in Medicine*, 8:207-19, 1979
138. **Seedhom BB & DJ Hargreaves:** Transmission of the load in the knee joint with special reference to the role of the menisci. Part II: experimental results, discussion and conclusions. *Engineering in Medicine*, 8:220-8, 1979
139. **Siwek KW & JP Rao:** Bilateral simultaneous rupture of the quadriceps tendons. *Clin Orthop*, 131:252-4, 1978
140. **Smidt GL:** Biomechanical analysis of knee flexion and extension. *J Biomech*, 6:79-92, 1973
141. **Smith AJ:** Estimates of muscle and joint forces at the knee and ankle during a jumping activity. *J Human Movem Stud*, 1:78-86, 1975
142. **Soderberg GL & TM Cook:** An electromyographic analysis of quadriceps femoris muscle setting and straight leg raising. *Phys Ther*, 63:1434-8, 1983
143. **Steiner CA & LH Palmer:** Simultaneous bilateral rupture of the quadriceps tendon. *Am J Surg*, 78:752-5, 1949
144. **Svensson O, J Ekholm, U Arborelius, R Nisell & G Németh:** Ankle joint load and leg muscle activity in various working postures during machine milking. *Ergonomics*, 28:683-98, 1985
145. **Thistle HG, HJ Hislop, M Moffroid & EW Lowman:** Isokinetic contraction: A new concept of resistive exercise. *Arch Phys Med Rehab*, 48:279-82, 1967
146. **Thorstensson A, G Grimby & J Karlsson:** Force-velocity relations and fiber composition in human knee extensor muscles. *J Appl Phys*, 40:12-6, 1976
147. **Tropp H:** Functional Instability of the Ankle Joint. *Depts of Orthop Surg and of Clin Neurophysiol*, Linköping University, Thesis, No 202, 1985
148. **Undeutsch K, R Kupper, I Löwenthal, KH Gärtner, T Luopajarvi, K Rauterberg, MJ Karvonen & J Rutenfranz:** Arbeitsmedizinische Untersuchungen über Ladearbeiten auf einem Grossflughafen. *Int Arch Occup Environ Health*, 50:59-75, 1982
149. **Vredenburg J & G Rau:** Surface electromyography in relation to force, muscle length and endurance. In: JE Desmedt (ed), *New Developm in EMG and Clin Neurophys*, Vol. 1, Karger, Basel:607-22, 1973
150. **Wahrenberg H, L Lindbeck & J Ekholm:** Knee muscular moment, tendon tension force and EMG during a vigorous movement in man. *Scand J Rehab Med*, 10:99-106, 1978
151. **Walker PS & JV Hajek:** The load-bearing area in the knee joint. *J Biomech*, 5:581-9, 1972
152. **Walker PS & MJ Erkman:** The role of the menisci in force transmission across the knee. *Clin Orthop*, 109:184-90, 1975
153. **Wendt PP & RP Johnson:** A study of quadriceps excursion, torque, and the effect of patellectomy on cadaver knees. *J Bone Joint Surg*, 67-A:726-32, 1985
154. **Wetzler SH & W Merkow:** Bilateral, simultaneous and spontaneous rupture of the quadriceps tendon. *JAMA*, 144:615-6, 1950
155. **Wilkie DR:** The mechanical properties of muscle. *Br Med Bull*, 12:177-82, 1956
156. **Williams M & L Stutzman:** Strength variation through the range of joint motion. *Phys Ther Rev*, 39:145-52, 1959
157. **Wilson JN:** Bilateral rupture of rectus femoris tendons in chronic nephritis. *Brit Med J*, 5032:1402-3, 1957

158. **Winter D:** Overall principle of lower limb support during stance phase of gait. *J Biomech*, 13:923–7, 1980
159. **Winter DA:** *Biomechanics of Human Movement*. John Wiley & Sons, 1979
160. **Winter DA:** Moments of force and mechanical power in jogging. *J Biomech*, 16:91–7, 1983
161. **Woods JJ & B Bigland-Ritchie:** Linear and non-linear surface emg/force relationships in human muscles. *Am J Phys Med*, 62:287–99, 1983
162. **Yang JF & D Winter:** Electromyographic amplitude normalization methods: improving their sensitivity as diagnostic tools in gait analysis. *Arch Phys Med Rehab*, 65:517–21, 1984
163. **Zernicke RF, J Garhammer & FW Jobe:** Human patellar-tendon rupture. *J Bone Joint Surg*, 59-A:179–83, 1977
164. **Zuniga EN, XT Truong & DG Simons:** Effects of skin electrode position on averaged electromyographic potentials. *Arch Phys Med Rehab*, 51:264–72, 1970
165. **Zuppinger H:** *Die aktive Flexion im unbelasteten Kniegelenk*. Wiesbaden, Verlag von JF Bergmann, 1904
166. **Öberg B:** Lower Extremity Muscle Strength in Soccer Players. Depts of Orthop Surg and of Clin Neurophysiol, Linköping University, Thesis, No 190, 1984