

Fuel Type Mapping with Remote Sensing and Machine Learning



What we do:

- Gather information and analysis using best available science, technology & human expertise
- Assist risk-based decision making for wildfire prevention, preparedness, response & recovery

Who we are: 3 prongs

1. Fire Weather Forecasters,
2. Fire Behavior Specialists, and
3. IT and Data Scientists



Overview



1) BCWS FTL project: move towards NRT / continuous forest monitoring for improved situational awareness

- A. Partners/ Engagement.. Your feedback requested for coordinated approach
- B. Projects + components

2) Satellite fire mapping 2022 Operational Trial

- A. Method
- B. Learnings + next steps

BCWS Predictive Services
BCWS Geospatial Services

3) Satellite fuels mapping (goal: 2022 op. trial)

- A. Demo of proposed iteration towards method
 - A. Check data consistency on 10,000 sq km area! (1% of prov)
- B. Learnings + next steps



Partners and collaborators



BC Wildfire
Service

- Hopefully you?
- Your input appreciated!



RPAS




Sentinel1



Sentinel2




RCM




NiSAR (2023)



JAXA PALSAR-2



WILDFIRESAT



BCWS FTL project components





- Emerging tech collab w:
 - Canadian Space Agency (CSA): radar imaging
 - Italian Space Agency (ASI): hyperspectral imaging
 - Japanese Space Agency (JAXA): radar imaging. Int'l collaborators
- Mobile fuel typing app (ground data collection)
- Telegraph creek ground fuels study (2020, 2021, 2022 UAV study)
- **Active fire detection w Sentinel2**
 - operational trial 2022 season
- **Fuel type monitoring w Sentinel2**
 - Aiming for operational trial 2023 season
- Active fire detection (thermal UAV)
 - Aiming for operational trial 2023 season

Fort McMurray 2016 (Horse River fire) radar



BC Wildfire
Service



JAXA Quad-pol L-band SAR data

National Fire Polygon Database Extent

Quad-pol processing steps

- 1.Radiometric calibration
- 2.Faraday rotation correction (Default option)
- 3.Multilook (2x2)
- 4.T4 matrix generation
- 5.Box filter 7x7
- 6.Range doppler terrain correction (Copernicus 30m dem)
- 7.Convert to PolSARPro format, use NAN in no-data areas, project ground-reference on top, assign no-data areas (where no image overlap)

References

- https://www.eorc.jaxa.jp/ALOS-2/en/img_up/dis_pal2_can-forest_fire_20160509.htm
- <https://ieeexplore.ieee.org/document/8517822>

White Rock Lake (2021, BC) radar image



BC Wildfire
Service

JAXA L-band Dual-pol

Shown Difference:
(2022 - 2021) of


Re(C12)(red)
C22(green)
C11(blue)

** Fire scar is:
the big
pink/purple
thing **

Processing steps (SNAP):

- 1) Radiometric calibration
- 2) Multilook to approx square
- 3) 5x5 box filter
- 4) Range-Doppler TC
(Copernicus 30m dem)
- 5) Convert to PolSARPro/
ENVI format
- 6) Co-locate with GDAL

Scene R,G,B=[2:band 2, 4:band 4, 1:band 1]



Survey 123 Phone app



BC Wildfire
Service

The image displays the Survey 123 mobile application interface. On the left, the main screen shows the Survey 123 logo and "My Surveys" section, which includes a thumbnail of a forest fire scene labeled "BCWS Fuels Field Survey". Two open survey forms are shown on the right:

- Fuel Characterisation and FBP Fuel Typing**: A form for assessing fuel characteristics. It includes fields for "Survey Date and Time" (set to Tuesday, September 10, 2019, at 3:32 PM), "Assessor Name" (with a note about identifying surveyors), and "Location point" (a map showing a pin at 49°27'N 122°54'W).
- Fire Observations**: A form for reporting fire details. It includes fields for "Date and time", "Incident number", and "Location of observed fire behaviour" (with a map pinning feature). Below this is a "Weather Conditions" section with fields for "Temperature (°C)" and "Relative humidity (%)".

BCWS.PredictiveServices@gov.bc.ca


Telegraph creek study




BC Wildfire
Service



Field Observations and
Remote Sensing of
Wildland Fuels: A Case
Study, Bulkley Valley
Research Centre, Oct 8
2021 2021



Key finding:
Winter dates Help
Discern Fuel types!




Field Observations and
Remote Sensing of
Wildland Fuels: A Case
Study, Bulkley Valley
Research Centre, Oct 8
2021 2021

Key finding:
Multitemporal could
Help get 3d parameters

Telegraph creek: add Drone imagery



BC Wildfire
Service



Operational trial 2022 (BCWS)



1. Goal: satellite fire perimeter mapping

2. Approach

- Web app w PSU Agile team <https://github.com/bcgov/wps-fire-perimeter>
- ESA Copernicus download (better latency)
- Simple "A.I." Method(s)

3. What we learned

- Wins

4. Next steps

BCWS Predictive Services
BCWS Geospatial Services

Satellite fire mapping




- Opportunity to monitor progression of fires not actively suppressed
 - Add situational awareness
 - Reduce high-risk flight reqt's
- Complement existing methods
- Additional frequency & fidelity for perimeter updates
 - Better intel for growth projections & other predictive services products
- ***Stepping-stone, towards continuous fuels mapping***
- Start w Sentinel-2 & add more (tried Landsat & Sentinel-3)

Why the method is unique



- Somewhat unusual
 - Not a retrospective pre/post veg comparison
 - Single-date imagery used (*)
 - NBR / dNBR not used
 - Thermal IR band not used
 - Fixed threshold method (v fast)
 - Not a hotspot detection, can catch transient fires
 - Accessible:
 - no math/stats req'd
 - Free/open-source software & open data used!
 - Can try a simple "band math" expression in your preferred Geomatics tool (Arc, ENVI, SNAP, PCI, QGIS, ..etc)
- (*) Can bring multi-date back to help with tricky cases

Approach: direct "low latency" data access



https://github.com/bcgov/wps-research/blob/master/py/binary_polygonize.py

1. Band selection: False color coding (right)




BC Wildfire
Service

- Color encoding to generate map at right:


- Red: "B12" 2190 nm = 2.2 μm
- Green: "B11" 1610 nm = 1.6 μm
- Blue: "B9" 945 nm = 0.95 μm

i.e. The B12, B11 and B9 are respectively plotted as Red, Green and Blue on the screen

- Vegetation is blue
- Hotspots are red
- Burned areas are orange



"Somewhere between
visible and thermal"





R90436 2022


1) Sentinel2 data: Why use longest-waves?

RGB = RGB (visible) Battleship mountain (G72150) 20220910

RGB= (B12, B11, B9) shortwave IR. 3 longest-wave bands!



Battleship mountain (G72150) 20220910






2) Threshold

https://github.com/bcgov/bcws-psu-research/blob/master/cpp/sentinel2_active.cpp
https://github.com/bcgov/wps-research/blob/master/cpp/raster_dominant.cpp



BC Wildfire
Service



(B12 > B11 && B12 > B9)





- Find image areas that are "more red"
- False positives incl. Reflection off water
- In-house private cloud app uses GEE land-cover to exclude water

Battleship mountain (G72150) 20220910

3) Scrub



BC Wildfire
Service




- Remove water areas etc.
- GIMP used for manual scrubbing
- Weather, illumination or other image quality issues could necessitate more scrubbing
 - Or more AI

Battleship mountain (G72150) 20220910

4) Convert to polygon and compare



BC Wildfire
Service




Battleship mountain (G72150) 20220910

- KML outline (RED) viewed in Google Earth
- SWIR band preview saved to TIFF
 - Scaled to 8-bit (each band)
- 5. SME post processing not shown

4. Polygon: compare w existing data!



BC Wildfire
Service



Battleship mountain (G72150) 20220910 Poly
data 2022091021 (9-10pm)


Note: 12 Sep 2022 poly to boots on ground in 4h24m from image capture (approx noon)

5. Post processing



BC Wildfire
Service

- Post processing completed in conjunction with current perimeter, Plans Chief and GIS Specialist
- Left: Sentinel2 derived detection. Middle: Generalized shape Right: comparison




Comparing Sentinel2 result (right) with Firms





BC Wildfire
Service

NASA Firms (MODIS
and VIIRS) detection Aug 17



Purple – Sentinel-2 detection Aug. 17th
Orange – public perimeter as of 20220819



SWIR false color





Fire: N20881 Date: 20220817

Finding unknown fires





BC Wildfire
Service

- Found on July 28 from Jul 27 imagery
- Aug 20 we confirmed it was assigned a fire number (status out)
 - R90522 / Tahoots Lake




Small fires detected

- Sub-hectare fires observed
- Fire: G82427
- Date: 20220913
- Size: 0.20 ha (Google Earth)



Wins

- More than 300 "low latency" fire mapping updates generated & vetted
 - Sentinel-2 (ESA) and Landsat (NASA)
 - Low latency: less than 12h possible
- Mapped small or unknown fires
- Mapped fires under smoke cover
- Value recognized by front-line staff in 2022
 - Increasing number of requests from incidents for operational use





Learnings




BC Wildfire
Service

- Largest limitations: Frequency & Latency
 - 1-5 day repeat (Sentinel2), <= 9 day repeat (Landsat)
 - NRT access for Sentinel2 is possible (< 3 hours)
- Challenging cases:
 - Atmosphere, illumination, altitude, low-intensity fire, sub-canopy fire, data variation!
 - Multiple dates, more sophisticated algorithms needed to improve results
- Artificial Intelligence:
 - "*Computer-based Decision Support Systems*" ([NRCAN PFC AFT group Definition](#))



Next: *find the sweet spot between automatic vs explainable/accessible!*
Want to be a little more "AI"-like than (*)





Next Steps

- Also want physical models: Keen to explore **energy modelling w WildfireSat team!**
 - And sensor fusion, too!
- **NRT/URT data access?**
- Automate & extend
 - Train "more-automatic" methods --> ***Reduce human intervention!***
 - Add terrain, geometry, climate variables?
 - Machine / API access for Landsat data
 - IT resources needed to scale up
- Continue work w CSA, BC Forest Inventory, NRCAN, JAXA, ESA, NASA, ASI & more partners
 - Cloud penetrating imaging to support Fire & Fuels mapping
 - Moisture, wind speed or other applications

Satellite fuels mapping overview



- Concept + Data preparation
- Algorithm
 - Abcd
 - Kabcd
- Kamloops example
- Next steps

Province-wide fuel type mapping concept



BC Wildfire
Service

- Extend data eng. as Used in fire mapping
- Use as industry standard statistical inference
 - E.g. KNN (NFI & others)
 - More-automatic methods if necessary (ANN)
- Use existing fuels maps as ground ref
 - Use algorithms to find areas where ground-ref and satellite images are consistent! (*)
 - Incorporate local fuels studies & Annotation by local SME
 - Note: existing BCWS FTL is derived from Vegetation Resource Inventory (VRI) polygons
 - Multiple time stamps, rumored as old as 1950
- Use a "trajectory" approach (**)

(*) agreement between fixed-point of classifier w ground-ref
(*) Cross-reference RS observations with VRI changes

(**) Start with dense time series (year on year)
(**) Improvement: segment signatures w.r.t. time (change detection w.r.t a seasonal baseline)



A screenshot of a 'Data Catalogue' page from the BC Wildfire Service website. The page features a header with the BC logo and the title 'Data Catalogue'. Below the header is a search bar and a table of dataset information. The table includes columns for 'Dataset Name', 'Field Name', 'Type', 'Length', and 'Description'. The first few rows of the table are:

Dataset	Field	Type	Length	Description
BCLCS_LEVEL_3	BCLCS_LV_3	VARCHAR	10	drainage, and is described as either alpine, wetland, or upland. In rare cases, the polygon may be alpine wetland.
BCLCS_LEVEL_4	BCLCS_LV_4	VARCHAR	10	Classifies the vegetation types and Non-Vegetated cover types (as described by the presence of distinct types upon the land base within the polygon).
BCLCS_LEVEL_5	BCLCS_LV_5	VARCHAR	10	Classifies the vegetation density classes and Non-Vegetated categories.
INTERPRETATION_DATE	INTRP_DATE	DATE	7	The date on which the data was photo interpreted.
PROJECT	PROJECT_ID	VARCHAR	100	The business assigned name of the project. The name typically reflects a Timber Supply Area, an initiating Agency, or a land area.
REFERENCE_YEAR	REF_YR_ID	NUMBER	4	The year of the source data on which the interpretation is based. Known as the Reference Year in the VIF file.

Data preparation (proposed) Sentinel-2 & ref



- L1C data download from Google Cloud Platform (fast)
 - Select data with $\leq 7\%$ total cloud cover only
- Convert L1C to L2A (surface reflectance)
- Warp all bands to 20m, order by wavelength (increasing)
- Set no-data areas and $\geq 7\%$ cloud probability, to NAN (etc.)
 - Set cloud shadow to NAN
- Option: convert to ieee-standard float16 (save 50% space)
- BC Fuel type layer 2020 dataset (based on 2019 VRI)
 - rasterize (1-hot encoding)

Abcd algo

- NN inference (multithread C++ version)
 - Special case of [KNN](#) ($K=1$)
 - Fast results. But grainy(*)
 - ***Unique: Conceptually unifies regression & classification***
- Answers the question: A is to B as C is to ____? For square images
 - Answer: D
- No K. Data skip factor:
 - increase it, get faster results
 - ***Inspired by Emergency geomatics! Optional "shift" factor that does something cool (see *)***
- ***Three functions (also test cases for implementing classifiers)***
 - A != B != C: regression or classification (supervised)
 - Set A = B = C: image compression!
 - Set A = C: image translation
 - Simulate one kind of data from another
 - Abcd method is a crude "babel fish" for image data

(*) A. Beaudoin et al (2022)
Improved k-NN Mapping of Forest Attributes in Northern Canada Using Spaceborne L-band SAR, Multispectral and LiDAR Data

Note: uses texture!

(**) "tiny-AI challenge" to me (write pattern recognition algo in ≤ 100 lines of code)

- ***Use classification maps as "reference data"***
 - ***A = Satellite data***
 - ***B = class map***
 - ***C = new Satellite data***
 - ***D = new class map!***

Abcd method: Sicamous (BC 2021)




Image translation: translate "SWIR" imagery from "Visible" imagery. n=1476225, skip = 1111, 1329 = 0.09% retained
Abcd.exe swir.bin visible.bin swir_large.bin 1111 &


A) (B12,B11, B9) "SWIR"



C) input: "SWIR" (larger chip)



D) prediction: "Visible" (larger chip)



B) (B4, B3, B2) "Visible"




- KNN classification (multithread C++ code)
 - To include classification and regression option
- Data skip factor: increase it, get a result quicker
 - supports "Back of the envelope" tests for relationships in data
- Fuel type application: use class maps as "reference data"
 - A = Satellite data
 - B = existing class map (class labels stored as band names)
 - C = ***new Satellite data***
 - D = ***new class map!*** (class labels in band names)

Approach proposed w. Joanna Wand in 2020 used patches :

 - Such texture info likely needed
 - More computationally intensive
- Both presentations could use non-image ref. Data (just pack into image shape)

K-abcd algo test! 2022 trial output for "training" (K=7)



A) 12-band S2 image
(RGB=B12, B11, B9)
= B) (predict on same image)

B) Binary classification result
from operational trial
Rule-based classifier &
human cleaning


D) result (slightly different)
Skip = 1111
 $N = 511225 / 1111 = 460$
0.09% N retained

D) result (slightly different)
Skip = 111
 $N = 511225 / 111 = 4605$
0.9% N retained
(sampling grid touches the water)

FTL 2020 map (3 of 14 classes shown)



BC Wildfire
Service




Kamloops demo first run (check consistency)



- Use classification maps as "reference data"
 - A = Satellite data
 - B = Class map
 - C = Same satellite data
 - D = output: Compressed class map (Q: *can an image "code" the fuel types*)
- Iterative mindset:
 - Start w "back of envelope" assessment:
 - *do the images and fuel type map match?*

Data consistency check:
abcd.exe T10UFB_20180908ftl_2020.bin T10UFB_2018090811111



Date	Difference (skip = 11111)
20180809 (shown)	8.67e+06 (71 %)
20190806	8.95e+06
20200806	9.20e+06
20210803	9.09e+06

- According to this method the 2018 date matches the Fuel Type map better
- Can "replicate" a fuels map from single date imagery (using .09% of the original fuels map) "approximately"
- There are continuous areas of specific fuels that can be found and used as reference data

Next Steps for Fuels mapping



- Want non-categorical & frequently updated (e.g. monthly) forest parameters
 - Multiresolution updates (lower-res first)?
 - Standardized or data-driven mapping units? Cross-ref VRI changes w Satellite
- How?
- Good models: Computationally expensive
 - Iterate and improve data until consistent & predictive (keep checking)
- **Trajectory methods:** extend fuel types to succession types? Q: "What next"?
 - *Start with linear (temporal) interpolation on "dense" time series (~ 20-40 frames / year)*
 - *Adapt kabcd (KNN) impl. for missing values (cloud, cloud shadow) (*)*
- ***Main Goal: Operational trial 2023 for Fuels mapping***
 - Start at incident level and scale up with humans in loop?
 - Start at high level and drill down where people want more?
(*) decrease denominator in metric, or interpolate to fill in



BC Wildfire
Service

Thanks!

Questions?

Suggestions?

Ashlin.Richardson@gov.bc.ca