

Medication comparison

2 medications · 5 topics · September 2, 2026

Read across a row to compare the same topic.

Use this report to prepare questions for your pharmacist or prescriber. It cannot tell you which medicine is right for you. Do not start, stop, or switch a medicine based on this report.

☐ Highlighted rows contain different source details. They do not rank benefits, risks, or suitability.

What to compare	01	02
	Metformin Hydrochloride Public drug information	Empagliflozin Also listed as JARDIANCE Public drug information
FDA LABEL		
Uses		
Uses described in each medication's FDA label.		
Different source details		
	Public data FDA label excerpt 1 INDICATIONS AND USAGE Metformin hydrochloride tablets are indicated as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients 10 years of age and older with type 2 diabetes mellitus. Metformin hydrochloride tablets is a biguanide indicated as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients 10 years of age and older with type 2 diabetes mellitus. (1) Sources and freshness FDA drug label https://api.fda.gov/drug/label.json?search=set_id%3A%22073cf6d9-7142-4d8b-898e-174c62a1f76f%22 Retrieved Sep 3, 2026. Effective Aug 21, 2026. FDA label source text, not individualized medical advice or a complete pairwise interaction check. Source notices openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability. openfda-label: Selected the latest matching label among the first 12 source SPL sets. Other product labels may differ. nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted. nadac: Benchmarks cover up to 100 of 263 package NDCs, not every product variant.	Public data FDA label excerpt 1 INDICATIONS AND USAGE JARDIANCE is indicated: to reduce the risk of cardiovascular death and hospitalization for heart failure in adults with heart failure. to reduce the risk of sustained decline in eGFR, end-stage kidney disease, cardiovascular death, and hospitalization in adults with chronic kidney disease at risk of progression. to reduce the risk of cardiovascular death in adults with type 2 diabetes mellitus and established cardiovascular disease. as an adjunct to Excerpt only. Additional text or records are omitted from this report. Read the source for the full information. Sources and freshness FDA drug label https://api.fda.gov/drug/label.json?search=set_id%3A%2259ed43d9-b4e0-72a2-e063-6394a90a327d%22 Retrieved Sep 3, 2026. Effective Aug 26, 2026. FDA label source text, not individualized medical advice or a complete pairwise interaction check. Source notices openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability. nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted.

What to compare

Selected report topics

01

Metformin Hydrochloride

Public drug information

02

Empagliflozin

Also listed as JARDIANCE

Public drug information

FDA ADVERSE-REACTION LABEL
SECTION

Side effects

Each FDA label's adverse-reaction section. Different wording does not tell you which medication is safer.

Different source details

Public data

FDA label excerpt

6 ADVERSE REACTIONS The following adverse reactions are also discussed elsewhere in the labeling: Lactic Acidosis [see Boxed Warning and Warnings and Precautions (5.1)] Vitamin B 12 Deficiency [see Warnings and Precautions (5.2)] Hypoglycemia [see Warnings and Precautions (5.3)] For metformin hydrochloride tablets, the most common adverse reactions (>5.0%) are diarrhea, nausea/vomiting, flatulence, asthenia, indigestion, abdominal discomfort, and headache. (6.1) To report

Excerpt only. Additional text or records are omitted from this report. Read the source for the full information.

Sources and freshness

- [FDA drug label](https://api.fda.gov/drug/label.json?search=set_id%3A%22073cf6d9-7142-4d8b-898e-174c62a1f76f%22)
https://api.fda.gov/drug/label.json?search=set_id%3A%22073cf6d9-7142-4d8b-898e-174c62a1f76f%22

Retrieved Sep 3, 2026. Effective Aug 21, 2026.

FDA label source text, not individualized medical advice or a complete pairwise interaction check.

Source notices

- openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability.
- openfda-label: Selected the latest matching label among the first 12 source SPL sets. Other product labels may differ.
- nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted.
- nadac: Benchmarks cover up to 100 of 263 package NDCs, not every product variant.

Public data

FDA label excerpt

6 ADVERSE REACTIONS The following important adverse reactions are described below and elsewhere in the labeling: Diabetic Ketoacidosis in Patients with Type 1 Diabetes Mellitus and Other Ketoacidosis [see Warnings and Precautions (5.1)] Volume Depletion [see Warnings and Precautions (5.2)] Genitourinary Infections, including Urosepsis, Pyelonephritis, Necrotizing Fasciitis of the Perineum (Fournier's Gangrene), and Genital Mycotic Infections [see Warnings and Precautions

Excerpt only. Additional text or records are omitted from this report. Read the source for the full information.

Sources and freshness

- [FDA drug label](https://api.fda.gov/drug/label.json?search=set_id%3A%2259ed43d9-b4e0-72a2-e063-6394a90a327d%22)
https://api.fda.gov/drug/label.json?search=set_id%3A%2259ed43d9-b4e0-72a2-e063-6394a90a327d%22

Retrieved Sep 3, 2026. Effective Aug 26, 2026.

FDA label source text, not individualized medical advice or a complete pairwise interaction check.

Source notices

- openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability.
- nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted.

What to compare Selected report topics	01 Metformin Hydrochloride Public drug information	02 Empagliflozin Also listed as JARDIANCE Public drug information
FDA LABEL Warnings Label warnings and conditions when the medication should not be used. Different source details Missing warnings do not mean a medication is safe for a particular person.	Public data FDA label excerpt 5 WARNINGS AND PRECAUTIONS Lactic Acidosis: See boxed warning. (5.1) Vitamin B 12 Deficiency: Metformin may lower vitamin B 12 levels. Measure hematological parameters annually and vitamin B 12 at 2 to 3 year intervals and manage any abnormalities. (5.2) Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues: Increased risk of hypoglycemia when used in combination with insulin and/or an insulin secretagogue. Lower dose of insulin or insulin secretagogue may Contraindication 4 CONTRAINDICATIONS Metformin hydrochloride tablets are contraindicated in patients with: Severe renal impairment (eGFR below 30 mL/min/1.73 m 2) [see Warnings and Precautions (5.1)]. Hypersensitivity to metformin. Acute or chronic <i>Excerpt only. Additional text or records are omitted from this report. Read the source for the full information.</i> Sources and freshness FDA drug label https://api.fda.gov/drug/label.json?search=set_id%3A%22073cf6d9-7142-4d8b-898e-174c62a1f76f%22 Retrieved Sep 3, 2026. Effective Aug 21, 2026. FDA label source text, not individualized medical advice or a complete pairwise interaction check. Source notices - openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability. - openfda-label: Selected the latest matching label among the first 12 source SPL sets. Other product labels may differ. - nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted. - nadac: Benchmarks cover up to 100 of 263 package NDCs, not every product variant.	Public data FDA label excerpt 5 WARNINGS AND PRECAUTIONS Diabetic Ketoacidosis in Patients with Type 1 Diabetes Mellitus and Other Ketoacidosis: Consider ketone monitoring in patients with type 1 diabetes mellitus and consider ketone monitoring in others at risk for ketoacidosis, as indicated. Assess for ketoacidosis regardless of presenting blood glucose levels and discontinue JARDIANCE if ketoacidosis is suspected. Monitor patients for resolution of ketoacidosis before restarting. (5.1) Volume Contraindication 4 CONTRAINDICATIONS JARDIANCE is contraindicated in patients: with a hypersensitivity to empagliflozin or any of the excipients in JARDIANCE, reactions such as angioedema have occurred [see Warnings and Precautions (5.6)] . <i>Excerpt only. Additional text or records are omitted from this report. Read the source for the full information.</i> Sources and freshness FDA drug label https://api.fda.gov/drug/label.json?search=set_id%3A%2259ed43d9-b4e0-72a2-e063-6394a90a327d%22 Retrieved Sep 3, 2026. Effective Aug 26, 2026. FDA label source text, not individualized medical advice or a complete pairwise interaction check. Source notices - openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability. - nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted.

What to compare

Selected report topics

01

Metformin Hydrochloride

Public drug information

02

Empagliflozin

Also listed as JARDIANCE

Public drug information

FDA LABEL

FDA-labeled drug interactions

Each medication's label section, not a check of whether these medications interact with each other.

Different source details

These are individual medication label sections, not a complete pairwise interaction check. Selecting two drugs does not establish that they interact.

Public data

FDA label excerpt

7 DRUG INTERACTIONS Table 3 presents clinically significant drug interactions with metformin hydrochloride. Table 3: Clinically Significant Drug Interactions with Metformin Hydrochloride Carbonic Anhydrase Inhibitors Clinical Impact: Carbonic anhydrase inhibitors frequently cause a decrease in serum bicarbonate and induce non-anion gap, hyperchloremic metabolic acidosis. Concomitant use of these drugs with metformin hydrochloride may increase the risk for lactic acidosis.

Excerpt only. Additional text or records are omitted from this report. Read the source for the full information.

Sources and freshness

- [FDA drug label](https://api.fda.gov/drug/label.json?search=set_id%3A%22073cf6d9-7142-4d8b-898e-174c62a1f76f%22)
https://api.fda.gov/drug/label.json?search=set_id%3A%22073cf6d9-7142-4d8b-898e-174c62a1f76f%22

Retrieved Sep 3, 2026. Effective Aug 21, 2026.

FDA label source text, not individualized medical advice or a complete pairwise interaction check.

Source notices

- openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability.
- openfda-label: Selected the latest matching label among the first 12 source SPL sets. Other product labels may differ.
- nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted.
- nadac: Benchmarks cover up to 100 of 263 package NDCs, not every product variant.

Public data

FDA label excerpt

7 DRUG INTERACTIONS See Table 4 for clinically relevant interactions with JARDIANCE. Table 4 Clinically Relevant Interactions with JARDIANCE Diuretics Clinical Impact Coadministration of empagliflozin with diuretics resulted in increased urine volume and frequency of voids, which might enhance the potential for volume depletion. Intervention Before initiating JARDIANCE, assess volume status and renal function. In patients with volume depletion, correct this condition before

Excerpt only. Additional text or records are omitted from this report. Read the source for the full information.

Sources and freshness

- [FDA drug label](https://api.fda.gov/drug/label.json?search=set_id%3A%2259ed43d9-b4e0-72a2-e063-6394a90a327d%22)
https://api.fda.gov/drug/label.json?search=set_id%3A%2259ed43d9-b4e0-72a2-e063-6394a90a327d%22

Retrieved Sep 3, 2026. Effective Aug 26, 2026.

FDA label source text, not individualized medical advice or a complete pairwise interaction check.

Source notices

- openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability.
- nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted.

What to compare

Selected report topics

01

Metformin Hydrochloride

Public drug information

02

Empagliflozin

Also listed as JARDIANCE

Public drug information

EACH PRICE KEEPS ITS OWN SOURCE

Pricing and benchmarks

Check the product, quantity, date, and price type. NADAC is a pharmacy acquisition benchmark, not a patient price.

Different source details

Compare only like-for-like products and quantities. Public benchmarks never become purchasable cart offers.

Public data

Public acquisition benchmark

NADAC estimates pharmacy acquisition cost. This is not a retail cash price, a patient copay, or evidence of pharmacy availability.

\$0.69

30 EA

NADAC benchmark for 30 EA

Package NDC 42385094911

Effective or as-of date Aug 19, 2026.

CMS Medicaid NADAC. Retrieved Sep 3, 2026.

First of 16 source records, not a lowest-price selection.

Excerpt only. Additional text or records are omitted from this report. Read the source for the full information.

Sources and freshness

- CMS Medicaid NADAC
<https://data.medicaid.gov/dataset/fbb83258-11c7-47f5-8b18-5f8e79f7e704>
Retrieved Sep 3, 2026. Dataset 2026.

NADAC is a pharmacy acquisition-cost benchmark, not a retail cash price, insured copay, or negotiated patient price.

Source notices

- openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability.
- openfda-label: Selected the latest matching label among the first 12 source SPL sets. Other product labels may differ.
- nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted.
- nadac: Benchmarks cover up to 100 of 263 package NDCs, not every product variant.

Public data

Public acquisition benchmark

NADAC estimates pharmacy acquisition cost. This is not a retail cash price, a patient copay, or evidence of pharmacy availability.

\$335.57

30 EA

NADAC benchmark for 30 EA

Package NDC 00597015337

Effective or as-of date Aug 19, 2026.

CMS Medicaid NADAC. Retrieved Sep 3, 2026.

First of 6 source records, not a lowest-price selection.

Excerpt only. Additional text or records are omitted from this report. Read the source for the full information.

Sources and freshness

- CMS Medicaid NADAC
<https://data.medicaid.gov/dataset/fbb83258-11c7-47f5-8b18-5f8e79f7e704>
Retrieved Sep 3, 2026. Dataset 2026.

NADAC is a pharmacy acquisition-cost benchmark, not a retail cash price, insured copay, or negotiated patient price.

Source notices

- openfda-ndc: Showing one matching finished-product ingredient group. Other formulations may exist; this does not imply interchangeability.
- nadac: NADAC returned the latest 100 rows for a package batch. Older package records may be omitted.

TAKE IT TO YOUR NEXT CONVERSATION

Questions for your pharmacist or prescriber

- Which benefits and risks matter for my health history?
- Could this medicine affect anything else I take, including supplements?
- What would I actually pay for my prescribed product and quantity?

FDA excerpts are not a complete medication guide. Missing information does not mean no risk. Public acquisition benchmarks are not patient prices. Source dates appear with each topic.