

Supplementary Material: “Climate change decouples drought from early winegrape harvests in France”

Benjamin I Cook^{1,2} & Elizabeth M Wolkovich^{3,4}

June 10, 2015

Regional GHD Analyses

To analyze each site individually, we regressed the GHD for each site in GHD-Core against CRU climate data averaged within one degree of the site location. As with GHD-Core, temperature is the dominant control on GHD in all eight regional series (Supplementary Figures 4–11), and all the temperature regressions are highly significant ($p \leq 0.01$). For the period 1901–1980, the temperature sensitivities (regression slopes) across sites range from a minimum of $-3.70 \text{ days } ^\circ\text{C}^{-1}$ at SRv to $-7.39 \text{ days } ^\circ\text{C}^{-1}$ at Cha1. Significant ($p \leq 0.05$) moisture sensitivities (precipitation and PDSI) are also found at seven of the eight sites: Als, Bor, Bur, Cha1, Lan, LLV, and Swi. At the eighth site, SRv, the precipitation regression is marginally significant ($p = 0.055$) and is insignificant for PDSI.

At all sites, GHD temperature sensitivities remain significant for the 1981–2007 interval, indicating that temperature remains an important control on winegrape phenology in more recent decades. By contrast, most sites show a significant weakening of the two moisture sensitivities. Both the precipitation and PDSI regressions with GHD become insignificant ($p > 0.05$) during 1981–2007 for Bor, Bur, Cha1, Lan, LLV, and Swi. At SRv, only precipitation was marginally significant for 1901–1980, and this regression weakens considerably. At Als, the PDSI regression becomes insignificant, while precipitation does not change appreciably. Analyses of the individual regional GHD series therefore support the general conclusions drawn from analyses of the GHD-Core composite index, demonstrating an overall large-scale weakening of moisture controls on GHD in Western Europe.

Temperature versus Moisture Comparisons

We hypothesize that, in Western Europe, moisture impacts on winegrape phenology occur primarily through the indirect effect of moisture on growing season temperatures. Further, we argue that because of anthropogenic warming trends in recent decades, winegrape phenology has become decoupled from moisture variability. Here we, present evidence in support of these hypotheses.

A strong connection between soil moisture and warm season temperatures over the Mediterranean and Western Europe has been well established (1, 2), occurring primarily through the control of soil moisture on surface energy partitioning. In regions where evapotranspiration is limited by the supply of moisture at the surface, wetter soils will lead to increases in evapotranspiration and latent heat flux at the expense of sensible heating, keeping surface soil and air temperatures relatively cool. Conversely, if soils are dry, sensible heat fluxes will dominate, and soil and air temperatures will be higher. If these are the primary physics operating in our GHD-Core region, we would therefore expect negative relationships between temperature and precipitation or PDSI during the spring and summer.

Indeed, for both MJJ (Supplementary Figure 12) and JJA (Supplementary Figure 14) we can see significant negative relationships between both temperature and precipitation and PDSI during the 1901–1980 interval. The relationships are generally stronger during JJA, when solar energy inputs are larger and land-atmosphere interactions are expected to be stronger. During the more recent interval (1981–2007), the temperature-moisture relationships generally weaken, especially during JJA. Additionally, temperatures during 1981–2007 are substantially warmer compared to 1901–1980, reflecting recent warming trends driven primarily by anthropogenic greenhouse gas forcing. This additional warmth (unrelated to moisture variability), combined with the apparent weakening of the temperature-moisture coupling strength itself, supports our conclusions regarding the mechanisms by which moisture influences GHD variability and the recent weakening of this relationship.

Monte-Carlo Analyses

Interpretations of the composite climate anomaly analyses (Figure 4, main manuscript) and comparisons of climate distributions (Supplemental Figure 15, top panel) for extreme early harvest years may be affected by sampling uncertainties or biases, especially during the post 1981 period when the number of early harvest years to

sample from is small. For example, from 1600–1981, 72 years qualify as early harvests with GHD-Core anomalies of -7.67 days or earlier. Because of the last years represented in the climate reconstructions, however, from 1981 to 2007 only 18 years qualify for PDSI, 13 years for temperature, and 11 years for precipitation. To estimate the impact on our analyses of these potential sampling uncertainties, we conducted a Monte-Carlo analysis where, in each of 10,000 iterations, we resampled climate anomalies from the early harvest years with replacement for the 1600–1980 period. In each iteration, we resampled n times, where n is equal to the number of early harvest years from 1981–2007 (so 11 for precipitation, 13 for temperature, and 18 for PDSI).

Kernal density functions for the 10,000 resampled mean precipitation and PDSI anomalies are shown in the bottom panels of Supplemental Figure 15. As in the main analysis, PDSI shows the strongest shift with early harvest dates before and after 1980. Repeating the One Sided Student's t-test during each of these iterations found significant differences ($p \leq 0.05$) in 99% of cases, indicating our finding (that PDSI no longer indicates drought during early harvests post-1980) is quite robust. The Monte-Carlo results were not as robust for precipitation: only 29% of the iterations found marginally significantly ($p \leq 0.10$) drier conditions during the pre-1980 period compared to post-1980. In the precipitation reconstruction, however, we are severely limited in our sample size post-1980 (only 11 years, compared to 18 for PDSI). Our conclusions, however, are still strongly supported by both the observational analysis (using 20th century climate data) and the PDSI reconstruction.

References

1. Fischer, E. M., Seneviratne, S. I., Vidale, P. L., Lüthi, D. & Schär, C. Soil Moisture–Atmosphere Interactions during the 2003 European Summer Heat Wave. *Journal of Climate* **20**, 5081–5099 (2007). URL <http://dx.doi.org/10.1175/JCLI4288.1>.
2. Miralles, D. G., Teuling, A. J., van Heerwaarden, C. C. & Vila-Guerau de Arellano, J. Mega-heatwave temperatures due to combined soil desiccation and atmospheric heat accumulation. *Nature Geoscience* **7**, 345–349 (2014). URL <http://dx.doi.org/10.1038/ngeo2141>.

Table 1: Regional GHD series from DAUX used in construction of the GHD-Core composite series. Included are the three letter codes for each site, their geographic locations (units of decimal degrees), and their mean harvest dates (day of year) for various intervals.

GHD Series	Site Code	Lat	Lon	1600-1900	1600-1980	1901-1950	1951-1980	1981-2007
Alsace	Als	48.17	7.28	282.81	281.87	272.12	287.16	277.70
Bordeaux	Bor	45.18	-0.75	269.01	268.22	263.85	270.41	259.75
Burgundy	Bur	47.32	5.04	269.92	270.07	269.44	272.58	262.15
Champagne 1	Chal	47.98	4.28	266.88	267.52	267.01	270.02	264.92
Languedoc	Lan	43.60	3.87	272.43	270.28	260.97	266.88	263.40
Low Loire Valley	LLV	47.15	0.22	286.12	284.61	282.69	282.56	275.33
Southern Rhone Valley	SRv	43.98	5.05	269.20	269.10	268.84	268.46	257.87
Switzerland (Lake Geneva)	Swi	46.57	6.52	286.39	283.87	273.86	275.22	263.00

Table 2: For each regional GHD series used in GHD-Core, the fraction of years with observations for various time intervals.

	1354-2007	1600-1900	1600-2007	1800-2007	1900-2007
Als	0.401	0.578	0.642	0.837	0.824
Bor	0.500	0.648	0.738	0.995	0.991
Bur	0.925	0.997	0.990	0.986	0.972
Cha1	0.280	0.259	0.449	0.880	0.981
Lan	0.442	0.635	0.689	0.438	0.833
LLV	0.310	0.332	0.498	0.976	0.963
SRv	0.690	0.970	0.951	0.942	0.898
Swi	0.749	1.00	1.00	1.00	1.00

Table 3: Spearman's rank correlations (1354-2007) between the various regional GHD series used to construct GHD-Core.

	Als	Bor	Bur	Cha1	Lan	LLV	SRv	Swi
Als	1.00							
Bor	0.601	1.00						
Bur	0.574	0.615	1.00					
Cha1	0.632	0.675	0.799	1.00				
Lan	0.550	0.467	0.450	0.655	1.00			
LLV	0.503	0.712	0.776	0.709	0.359	1.00		
SRv	0.414	0.249	0.456	0.346	0.765	0.456	1.00	
Swi	0.604	0.554	0.562	0.569	0.705	0.697	0.501	1.00

Table 4: Day of year anomalies in the regional GHD series used to construct GHD-Core. Anomalies are calculated relative to the baseline mean dates from 1600-1900. Also included are results for the GHD-Core index and GHD-All, compositized from all 27 regional GHD series in DAUX. For GHD-Core and GHD-All, the harvest date anomalies were calculated for the regional series individually before averaging together. Because the regional GHD series are not all equally represented in all years, anomalies in GHD-Core and GHD-All are slightly non-zero for the baseline averaging period (1600–1900).

	1600-1900	1600-1980	1901-1950	1951-1980	1981-2007
Als	0	-0.94	-10.69	4.35	-5.11
Bor	0	-0.79	-5.16	1.40	-9.26
Bur	0	0.15	-0.48	2.66	-7.78
Chal	0	0.64	0.13	3.14	-1.96
Lan	0	-2.14	-11.46	-5.55	-9.03
LLV	0	-1.51	-3.42	-3.56	-10.79
SRv	0	-0.10	-0.36	-0.74	-11.33
Swi	0	-2.52	-12.53	-11.17	-23.39
GHD-Core	-0.32	-1.02	-5.16	-1.11	-10.24
GHD-All	-0.25	-0.85	-5.13	0.36	-8.91

Table 5: As Table 4, but for inter-annual standard deviation.

	1600-1900	1600-1980	1901-1950	1951-1980	1981-2007
Als	8.74	9.56	9.12	7.09	8.74
Bor	9.51	9.00	6.27	7.00	9.56
Bur	9.61	9.19	6.55	8.11	7.97
Chal	8.81	8.88	7.85	10.12	8.64
Lan	8.63	9.03	6.51	5.69	5.39
LLV	10.29	9.21	6.56	8.01	7.41
SRv	8.68	8.46	8.62	5.55	5.93
Swi	10.09	10.71	7.19	6.68	8.11
GHD-Core	7.67	7.49	5.39	6.24	7.10
GHD-All	7.03	7.02	5.55	6.55	6.75

Table 6: Coefficients and p-values from ordered logit models of wine quality data (on a scale of 0 to 5) and grape harvest dates (GHD) and CRU May-July seasonal temperatures for the periods 1900-1980 and 1981-2001. For more details on data and analyses see the Methods section in the main text.


	GHD: 1900-1980	GHD: 1981-2001	Temp: 1900-1980	Temp: 1981-2001
Red Bordeaux	-0.1117 (<0.01)	-0.133 (0.03)	1.244 (<0.01)	1.308 (0.01)
White Bordeaux	-0.084 (<0.01)	-0.079 (0.14)	0.951 (<0.01)	1.109 (0.02)
Red Burgundy	-0.102 (<0.01)	-0.101 (0.18)	0.403 (0.06)	0.612 (0.23)
White Burgundy	-0.093 (0.02)	-0.144 (0.07)	0.564 (0.04)	1.262 (0.03)

Table 7: Coefficients and p-values from ordered logit models of wine quality data (on a scale of 0 to 5) and CRU May-July seasonal precipitation and Palmer Drought Severity Index (PDSI) for the periods 1900-1980 and 1981-2001. For more details on data and analyses see the Methods section in the main text.


	Prec: 1900-1980	Prec: 1981-2001	PDSI: 1900-1980	PDSI: 1981-2001
Red Bordeaux	-0.013 (<0.01)	-0.011 (0.12)	-0.457 (<0.01)	-0.119 (0.61)
White Bordeaux	-0.014 (<0.01)	-0.014 (0.06)	-0.291 (0.02)	-0.234 (0.32)
Red Burgundy	-0.018 (<0.01)	0 (0.96)	-0.273 (0.03)	0.12 (0.64)
White Burgundy	-0.011 (0.05)	-0.013 (0.08)	-0.101 (0.52)	-0.199 (0.41)

Table 8: Comparison of regression statistics between GHD-Core and the CRU climate data (May-June-July, MJJ) for 1901-1980. Shown are results for cases where all data was linearly detrended over time prior to the regression calculation (detrended) versus the case where no detrending occurred (normal).

	Slope (normal)	Slope (detrended)	R2 (normal)	R2 (detrended)	p-val (normal)	p-val (detrended)
1901-1981						
Temperature	-6.044534	-6.218814	0.704	0.780	< 0.00000001	< 0.00000001
Precipitation	0.068827	0.068518	0.241	0.252	0.000004	0.000002
PDSI	1.682021	1.667401	0.133	0.137	0.000890	0.000709
1981-2007						
Temperature	-6.354566	-5.038086	0.642	0.473	0.000001	0.000073
Precipitation	0.033541	0.019836	0.051	0.030	0.258433	0.384498
PDSI	1.067015	0.429598	0.043	0.012	0.301859	0.592694




Supplementary Figure 1: Geographic locations for all 27 regional GHD time series in DAUX. Highlighted in red are the sites that comprise the GHD-Core index: Alsace (Als), Bordeaux (Bor), Burgundy (Bur), Champagne 1 (Cha1), Langue-doc (Lan), the Lower Loire Valley (LLV), the Southern Rhone Valley (SRv), and Switzerland at Lake Geneva (Swi). The dashed black box indicates the GHD-Core region (2°W – 8°E , 43°N – 51°N) over which climate anomalies from the CRU instrumental climate datasets and the three climate reconstructions were averaged for the various regression analyses.




Supplementary Figure 2: Number of observations (i.e., regional GHD series) represented in each year of the GHD-Core index.


GHD-Core vs Regional Series




Supplementary Figure 3: Time series (1600–2007) of GHD-Core (blue solid line) and each individual regional GHD series (green dashed lines) used in the construction of GHD-Core.




Supplementary Figure 4: Comparisons between the GHD series from ALS and May-June-July temperature, precipitation, and PDSI from the CRU 3.21 climate grids. Top panels: point-by-point Spearman's rank correlations for 1901–1980 and 1981–2007 (location of ALS is shown by the black dot). Bottom panels: linear regression plots for the same intervals against CRU climate data averaged within one degree of the site location.




Supplementary Figure 5: Same as Figure 4, but for Bor.




Supplementary Figure 6: Same as Figure 4, but for Bur.




Supplementary Figure 7: Same as Figure 4, but for Cha1.




Supplementary Figure 8: Same as Figure 4, but for Lan.




Supplementary Figure 9: Same as Figure 4, but for LLV.




Supplementary Figure 10: Same as Figure 4, but for SRv.




Supplementary Figure 11: Same as Figure 4, but for SWi.




Supplementary Figure 12: Temperature versus moisture (precipitation and PDSI) during May-June-July for the GHD-Core region from the CRU 3.21 climate grids for two periods: 1901–1980 and 1981–2007. Temperature during MJJ has a significant negative relationship with both precipitation and PDSI, indicating the tendency for warmer conditions during drier years. Over the more recent period, the relationship between temperature and PDSI breaks down, but the relationship with precipitation remains largely consistent.



Supplementary Figure 13: Same as Figure 3 from the main manuscript, but for June-July-August (JJA) climate.



Supplementary Figure 14: Same as Supplemental Figure 12, but for June-July-August (JJA) climate. During JJA, the temperature moisture relationship is stronger over this region and, in both precipitation and PDSI, these relationships breakdown in the recent decades (1981–2007).



Supplementary Figure 15: Top row: box plots of pooled precipitation and PDSI anomalies from the climate reconstructions for years with early harvest date anomalies in GHD-Core. Bottom row: kernel density functions of recalculated mean precipitation and PDSI anomalies associated with early harvest dates, based on 10,000 Monte-Carlo resamplings with replacement. Dashed vertical lines represent the mean anomalies calculated from the actual early harvest years from 1600–1980 (blue) and 1981–2007 (green).