

University of Wisconsin-Madison  
Engineering Physics Department  
Spring 2011 Qualifying Exams

# Modern Physics

You must solve 4 out of the 6 problems.  
Start each problem on a new page.

**SHOW ALL YOUR WORK.**  
**WRITE ONLY ON THE FRONT PAGES OF THE**  
**WORKSHEETS, NOT ON THE EXAM PAGES**

Grading is based on both the final answer and work done in reaching your answer. All problems receive an equal number of points.

Clearly indicate which problems you want graded. If you do not indicate which problems are to be graded, the first four solutions you provide will be graded.

1. \_\_\_\_\_
2. \_\_\_\_\_
3. \_\_\_\_\_
4. \_\_\_\_\_
5. \_\_\_\_\_
6. \_\_\_\_\_

Student No. \_\_\_\_\_

Engineering Physics Department  
Spring 2011 Qualifying Exams

**Problem 1.** A sample containing atoms of nuclide  $N_1$  ( ${}^A_Z X_N$ ) is placed in a neutron beam with a flux  $\phi$ . The radionuclide  $N_2$  ( ${}^{A'}_{Z'} X_{N'}$ ) is produced from the (n,p) neutron reaction with a cross section  $\sigma_{p1}$ . Radionuclide  $N_2$  is a beta emitter ( $\beta^-$ ) while in the neutron beam, nuclide  $N_2$  can be transmuted through the (n,alpha) reaction with a cross section  $\sigma_{\alpha 2}$ . The initial concentration of  $N_1$  at  $t = 0$  is  $N_0$  and that of  $N_2$  at  $t = 0$  is zero.

- (a) (1.5pts) What are atomic number  $A'$ , neutron number  $N'$  and proton number  $Z'$  for radionuclide  $N_2$  in terms of the  $A$ ,  $Z$  and  $N$  of the original nuclide?
- (b) (1.5pts) The decay of radionuclide  $N_2$  leads to nuclide  $N_3$  ( ${}^{A''}_{Z''} X_{N''}$ ). What are  $A''$ ,  $N''$ , and  $Z''$  for this nuclide in terms of the  $A$ ,  $Z$  and  $N$  of the original nuclide?
- (c) (3pts) Write out the rate equations for radionuclides  $N_1$  and  $N_2$ .
- (d) (4pts) Assuming that the (n,alpha) reaction rate for  $N_2$  is quite small and can be neglected as a first approximation, develop a general expression for the time-dependent concentrations of the radionuclides  $N_1$  and  $N_2$ .

Engineering Physics Department  
Spring 2011 Qualifying Exams

**Problem 2.**  $K^+$  mesons (positive kaons) decay into other particles, particularly pions. Their proper mean lifetime (i.e. the average time they exist before decaying) is  $\tau_0 = 1.24 \times 10^{-8}$  sec and their rest energy is 494 MeV. Suppose a beam of kaons is created in an accelerator laboratory with velocity  $v = 0.9c$

- a) (2pts) What is the energy of the kaons in the laboratory?
- b) (2pts) What will the kaon's mean lifetime be in the laboratory? Is it longer or shorter than  $\tau_0$ ?
- c) (2pts) How far will the kaon travel (on the average) in the laboratory's frame of reference?
- d) (2pts) Suppose the kaon beam is directed at another kaon beam with the same speed but moving in the opposite direction, so that a head on collision occurs as determined in the laboratory. What is the speed of the second kaon beam as determined in a frame of reference moving along with the first kaon beam?



- e) (2pts) What is the total energy of a colliding pair of kaons, one from each beam, as measured in the laboratory?


Student No. \_\_\_\_\_

Engineering Physics Department  
Spring 2011 Qualifying Exams

**Problem 3.** Write down the Semi-empirical Mass Formula for nuclei, and qualitatively describe the origin of each term in the formula for the binding energy per nucleon. Give a simple physical argument for the functional dependence of each of the terms on  $A$  (mass number) and  $Z$  (charge).

Engineering Physics Department  
Spring 2011 Qualifying Exams

**Problem 4.** A particle of mass  $m$  is subject to the potential distribution shown in the figure below. Consider the case when the energy of the particle is less than  $V_0$ .



- (a) (2pts) Draw qualitatively the wavefunctions of the first two energy eigenvalues for the potential in the figure.
- (b) (1pt) In the limit when  $V_0 \rightarrow \infty$ , is the ground state eigenvalue larger or smaller than in part (a)?
- (c) (2pts) What are the general solutions to the wavefunctions in the various regions?
- (d) (5pts) Determine the relations (expressions) required to find the lowest-energy and the next lowest-energy solution.

Student No. \_\_\_\_\_

Engineering Physics Department  
Spring 2011 Qualifying Exams

**Problem 5.** (a) (8pts) Determine the neutron energy needed to produce neutron-induced fission of the following nuclei (neglect recoil): (i)  ${}^{241}_{95}\text{Am}$  and (ii)  ${}^{242}_{95}\text{Am}$ .

Mass [amu] where 1 amu = 931.502 MeV/c <sup>2</sup>	E <sub>a</sub> [MeV] (for the A+1 nucleus) where E <sub>a</sub> = activation energy
$M({}^{241}_{95}\text{Am}) = 241.056824$	6.5
$M({}^{242}_{95}\text{Am}) = 242.059542$	6.2
$M({}^1_0\text{n}) = 1.008665$	---
$M({}^{243}_{95}\text{Am}) = 243.061375$	---

(b) (2pts) Briefly explain the physical origin of the difference in fissile potential of  ${}^{241}_{95}\text{Am}$  and  ${}^{242}_{95}\text{Am}$ .

Student No. \_\_\_\_\_

Engineering Physics Department  
Spring 2011 Qualifying Exams

**Problem 6.** A proton is incident on  ${}^7_3\text{Li}$  which undergoes a (p,n) reaction.

(a) (2pts) Compute the  $Q$ -value of the reaction.

(b) (8pts) Develop an expression for the threshold energy for the reaction using the basic momentum and energy conservation laws (recall, at threshold, the product particles have equal speed and move in the same direction as the incoming particle).

<u>Mass</u> [amu] (where 1 amu = 931.502 MeV/c <sup>2</sup> )	<u>Mass</u> [amu]
$M({}^1_1\text{H}) = 1.007825$	$M({}^1_0\text{n}) = 1.008665$
$M({}^7_3\text{Li}) = 7.016003$	$M({}^7_2\text{He}) = 7.02803$
$M({}^7_4\text{Be}) = 7.016928$	$M({}^8_4\text{Be}) = 8.005305$