

SMC-Guided Diffusion for General Optimisation

Brendan Dowling, Dr O. Deniz Akyildiz

September 6, 2024

Imperial College London

Table of contents

1. Introduction
2. Background
3. SMCDiffOpt
4. Experimental Results
5. Conclusion

Introduction

Hello, world! Akyildiz et al. (2020), Anderson (1982), Del Moral (2011), Dou and Song (2023), and Ho et al. (2020)

Background

Text

Text

Text

Text

SMCDiffOpt

Text

Text

Text

Text

Experimental Results





GMM Experiment — Setup

Consider a Gaussian mixture model (GMM), p_{data} , with 25 equally weighted ($\omega_{i,j} \propto 1$) d_x -dimensional Gaussian random variables with means $\mu_{i,j} := (8i, 8j, \dots, 8i, 8j) \in \mathbb{R}^{d_x}$ for $(i, j) \in \{-2, \dots, 2\}^2$ and unit variance. We generate some d_y -dimensional measurement, \mathbf{y} , according to the following process:

- Draw $\tilde{A} \sim \mathcal{N}(0, 1)^{d_y \times d_x} \in \mathbb{R}^{d_y \times d_x}$ and compute its SVD decomposition, USV^\top .
- Sample $s_{i,j} \sim \mathcal{U}[0, 1]$ for $(i, j) \in \{-2, \dots, 2\}^2$.
- Set $A := U \text{diag}(\{s_{i,j}\}) V^\top$.
- Draw $\mathbf{x}_* \sim p_{\text{data}}$ and set $\mathbf{y} := A\mathbf{x}_* + \sigma_y \epsilon$, $\epsilon \sim \mathcal{N}(\mathbf{0}_{d_y}, \mathbf{I}_{d_y})$.

The goal of the inverse problem is to take \mathbf{y} (with A and σ_y known), and use it to infer \mathbf{x}_* . We consider $d_y < d_x$, making the problem ill-posed.

GMM Experiment — Setup Visualised



GMM Experiment — Prior Model and TMPD Guidance

GMM Experiment — Performance Comparison

d_x	d_y	SMCDiffOpt	DPS	Π IGD	TMPD
8	1	1.35 ± 1.0	8.18 ± 7.5	3.16 ± 2.72	3.31 ± 2.86
	2	0.55 ± 0.43	2.72 ± 2.62	0.94 ± 0.89	1.34 ± 1.25
	4	0.19 ± 0.09	0.95 ± 0.9	0.09 ± 0.03	0.37 ± 0.33
80	1	1.65 ± 1.41	5.03 ± 4.63	3.08 ± 2.71	2.42 ± 1.71
	2	1.19 ± 1.07	3.14 ± 3.02	1.68 ± 1.62	1.35 ± 1.22
	4	1.11 ± 0.93	1.32 ± 1.21	0.84 ± 0.79	1.14 ± 0.9

Table 1: Sliced-Wasserstein distances of samples from particle samples.


Branin Experiment — Setup

Consider optimising the Branin function as an objective over some constrained region (the data manifold) as detailed in Kong et al. (2024). The Branin function is defined by

$$f(x_1, x_2) = a(x_2 - bx_1^2 + cx_1 - r)^2 + s(1 - t)\cos(x_1) + s \quad (1)$$

where $a = 1$, $b = \frac{5.1}{4\pi^2}$, $c = \frac{5}{\pi}$, $r = 6$, $s = 10$, $t = \frac{1}{8\pi}$. It has three global minima located at $(-\pi, 12.275)$, $(\pi, 2.275)$ and $(9.42478, 2.475)$, with a value of 0.397887. We consider some uniform prior sampling distribution, p_{data} , over an ellipse region centred at $(-0.2, 7.5)$ with semi-axis lengths $(3.6, 8)$, tilted 25° . This region covers two of the global minima.


Branin Experiment — Pitfalls of Gradient Optimisation




Text

Text


Black-Box Experiment — Real Sample



Black-Box Experiment — Generated Sample



Black-Box Experiment — Optimised Sample



Black-Box Experiment — Performance Comparison

CbAS	CMA-ES	Gradient Ascent	MINs	REINFORCE	DDOM	DiffOpt	SMCDiffOpt
0.433 ± 0.027	0.474 ± 0.021	0.510 ± 0.009	0.473 ± 0.003	0.483 ± 0.015	0.560 ± 0.044	0.614 ± 0.029	0.644 ± 0.024

Table 2: Normalised scores for SuperConductor experiment.

CbAS	CMA-ES	Gradient Ascent	MINs	REINFORCE	DDOM	DiffOpt	SMCDiffOpt
80.082 ± 4.933	87.774 ± 3.965	94.414 ± 1.719	87.483 ± 0.481	89.351 ± 2.708	103.600 ± 8.139	113.545 ± 5.322	119.059 ± 4.497

Table 3: Raw scores for SuperConductor experiment.

Conclusion

Text

Text

Thank you!





Akyildiz, Ö. D., Crisan, D., & Míguez, J. (2020). Parallel sequential Monte Carlo for stochastic gradient-free nonconvex optimization. *Statistics and Computing*, 30(6), 1645–1663. <https://doi.org/10.1007/s11222-020-09964-4>



Anderson, B. D. (1982). Reverse-time diffusion equation models. *Stochastic Processes and their Applications*, 12(3), 313–326. [https://doi.org/10.1016/0304-4149\(82\)90051-5](https://doi.org/10.1016/0304-4149(82)90051-5)



Del Moral, P. (2011). Central Limit Theorems. In *Feynman-Kac formulae: Genealogical and interacting particle systems with applications* (pp. 291–330). Springer
OCLC: 1063493341.

-  Dou, Z., & Song, Y. (2023). Diffusion Posterior Sampling for Linear Inverse Problem Solving: A Filtering Perspective. Retrieved June 8, 2024, from <https://openreview.net/forum?id=tpIXNcHZs1>
-  Ho, J., Jain, A., & Abbeel, P. (2020). Denoising Diffusion Probabilistic Models. <https://doi.org/10.48550/ARXIV.2006.11239>



Kong, L., Du, Y., Mu, W., Neklyudov, K., De Bortoli, V., Wang, H., Wu, D., Ferber, A., Ma, Y.-A., Gomes, C. P., & Zhang, C. (2024, April 29). *Diffusion Models as Constrained Samplers for Optimization with Unknown Constraints*. arXiv: 2402.18012 [cs]. Retrieved July 26, 2024, from <http://arxiv.org/abs/2402.18012>