

# Theoretical properties of nearest-neighbor distance distributions and novel metrics for high dimensional bioinformatics data

Bryan A. Dawkins<sup>1</sup>, Trang T. Le<sup>2</sup> and Brett A. McKinney<sup>1,3,\*</sup>

<sup>1</sup>Department of Mathematics, University of Tulsa, Tulsa, OK 74104, USA

<sup>2</sup>Department of Biostatistics, Epidemiology and Informatics, University of Pennsylvania, Philadelphia, PA 19104

<sup>3</sup>Tandy School of Computer Science, University of Tulsa, Tulsa, OK 74104, USA.

## Abstract

The performance of nearest-neighbor feature selection and prediction methods depends on the metric for computing neighborhoods and the distribution type of the underlying data. The effects of the distribution and metric, as well as the presence of correlation and interactions, are reflected in the expected moments of the distribution of pairwise distances. We derive general analytical expressions for the mean and variance of pairwise distances for  $L_q$  metrics for Gaussian and uniform data with  $p$  attributes and  $m$  instances. We use extreme value theory to derive results for metrics normalized by the max and min of attributes. These expressions are applicable to the analysis of continuous data such as gene expression. We derive similar analytical expressions for a new metric for genetic variants in GWAS data (categorical predictors) that accounts for minor allele frequency and transition/transversion ratio. We introduce a new metric for resting-state fMRI data that is applicable to correlation-based predictors from time series data. Derivations assume independent data, but empirically we also consider the effect of correlation. This study provides detailed derivations and expressions parameterized by  $q$ ,  $p$ ,  $m$  and other properties for a broad collection of bioinformatics data types.

## Author summary

## Introduction

Statistical models can deviate from expected behavior depending on whether certain properties of the underlying data are satisfied, such as having a Gaussian distribution. Nearest neighbor methods are further influenced by the choice of metric, such as Euclidean or Manhattan. For random normal data, for example, the variance of the pairwise distances of a Manhattan metric is proportional to the number of attributes ( $p$ ) whereas the variance is constant for a Euclidean metric. Relief methods [1–3] and nearest-neighbor projected distance regression (NDPR) use nearest neighbors to compute attribute importance scores and often use adaptive neighborhoods that rely on the mean and variance of the distance distribution. Thus, knowledge of the expected values for a given metric and data distribution may improve the performance of these feature selection methods.

For continuous data, the metrics most commonly used in nearest neighbor methods are  $L_q$  with  $q = 1$  (Manhattan) or  $q = 2$  (Euclidean). For data from standard normal ( $\mathcal{N}(0, 1)$ ) or standard uniform ( $\mathcal{U}(0, 1)$ ) distributions, the asymptotic behavior of the  $L_q$  metrics is known. However, detailed derivations of these distance distribution asymptotics are not readily available in the literature. We provide detailed derivations

of generalized expressions parameterized by  $q$ , attributes  $p$ , and samples  $m$ . We build on this mathematical formalism to derive the asymptotic properties of a new metric for categorical data in genome-wide association studies (GWAS) data [4]. We also derive asymptotic properties of a new metric introduced in the current study for resting-state fMRI (rs-fMRI) correlation data.

We derive asymptotic formulas for the mean and variance for three recently introduced GWAS metrics [4]. These metrics were developed for Relief-based feature selection to account for binary variant differences (two levels), allelic differences (three levels), and transition/transversion differences (five levels). The mean and variance expressions we derive for these multi-level categorical data types are parameterized by the minor allele frequency and the transition/transversion ratio.

The analysis of rs-fMRI data is a growing area for machine learning and feature selection [5–8]. For a given subject, a time series correlation or similar matrix is computed between regions of interest (ROIs), consisting of subsets of voxels from the neuroimage data. The time series represent functional activity of the ROI while the subject is at rest and the ROI typically corresponds to a region with known function for emotion or cognition. Thus, the dataset consists of pairwise ROI correlations for each of the  $m$  subjects. Nearest-neighbor based feature selection was applied to rs-fMRI in the private evaporative cooling method [cite pec], where the predictors were pairwise correlations between ROIs. The use of pairwise correlation predictors is a common practice because of convenience and differential connectivity between brain regions may be of biological importance [9]. However, one may be interested in the importance of features at the ROI level. Thus, in the current study we introduce a new metric to be used in NPDR with resting state correlation matrices that provides feature importance for ROIs. This metric is applicable to general time series-correlation (ts-corr) based data, and we derive asymptotic estimates for the mean and variance of distance distributions induced by our new ts-corr based metric.

The ability of nearest neighbor feature selection to identify association effects, like main effects or interaction effects, depends on neighborhood parameters, such as neighborhood radius or number of neighbors  $k$ . As  $k$  increases, nearest neighbor distance based algorithms are more sensitive to detecting main effects [10]. On the other hand, their ability to detect interaction effects decreases with increasing  $k$  [10, 11]. Correlation and interaction effects impact distance distributions by introducing positive skewness and increased variance, which can lead to changes in neighborhood inclusion. In order to understand how these statistical effects impact distance distributions in continuous and discrete data types, we first derive distance asymptotics for independently and identically distributed data. Using these derivations as a baseline, we can then determine how statistical effects and correlation change distance distributional properties from the null case.

In Section 1, we introduce preliminary notation and apply the Central Limit Theorem (CLT) and the Delta Method to derive asymptotics for pairwise distances. In Section 2, we present general derivations for continuously distributed data sets with  $m$  instances and  $p$  features. We focus on the cases of standard normal and standard uniform data distributions, but we derive analytical expressions parameterized by  $p$  and  $q$ . In Section 2.4 we use Extreme Value Theory (EVT) to address max-min normalized versions of  $L_q$  metrics, which are often used in Relief-based algorithms. In Section 3, we extend the derivations to categorical data with a binomial distribution for GWAS data. In Section 4, we present the final set of asymptotic results for our newly introduced time series correlation-based distance metric, with a particular emphasis on rs-fMRI data. Lastly, in Section 5, we demonstrate the effect of correlation in the attribute space on distance distributional properties.

# 1 Limit distribution for $L_q$ on null data

In the application of nearest-neighbor distance-based methods to continuous data, the distance between instances  $(i, j \in \mathcal{I}, |\mathcal{I}| = m)$  in the data set  $X^{m \times p}$  of  $m$  instances and  $p$  attributes (or features) is calculated in the space of all attributes ( $a \in \mathcal{A}, |\mathcal{A}| = p$ ) using a metric such as

$$D_{ij}^{(q)} = \left( \sum_{a \in \mathcal{A}} |d_{ij}(a)|^q \right)^{1/q}, \quad (1)$$

which is typically Manhattan ( $q = 1$ ) but may also be Euclidean ( $q = 2$ ). We use the terms “feature” and “attribute” interchangeably for the remainder of this work. The quantity  $d_{ij}(a)$ , known as a “diff” in Relief literature, is the projection of the distance between instances  $i$  and  $j$  onto the attribute  $a$  dimension. The function  $d_{ij}(a)$  supports any type of attributes (e.g., numeric and categorical). For example, the projected difference between two instances  $i$  and  $j$  for a continuous numeric ( $d^{\text{num}}$ ) attribute  $a$  may be

$$\begin{aligned} d_{ij}^{\text{num}}(a) &= \text{diff}(a, (i, j)) \\ &= |\hat{X}_{ia} - \hat{X}_{ja}|, \end{aligned} \quad (2)$$

where  $\hat{X}$  represents the standardized data matrix  $X$ . We use a simplified  $d_{ij}(a)$  notation in place of the  $\text{diff}(a, (i, j))$  notation that is customary in Relief-based methods. In NPDR, we omit the division by  $\max(a) - \min(a)$  used by Relief to constrain scores to the interval from  $-1$  to  $1$ , where  $\max(a) = \max_{k \in \mathcal{I}} \{X_{ka}\}$  and  $\min(a) = \min_{k \in \mathcal{I}} \{X_{ka}\}$ . The numeric  $d_{ij}^{\text{num}}(a)$  projection is simply the absolute difference between row elements  $i$  and  $j$  of the data matrix  $X^{m \times p}$  for the attribute column  $a$ .

All derivations in the following sections are applicable to nearest-neighbor distance-based methods in general, which includes not only NPDR, but also Relief-based algorithms. Each of these methods uses a distance metric, such as, Eq. 1 to compute neighbors for each instance  $i \in \mathcal{I}$ . Therefore, our derivations of asymptotic distance distributions are applicable to all methods that compute neighbors in order to weight features. The predictors used by NPDR, however, are the one-dimensional projected distances between two instances  $i, j \in \mathcal{I}$  given by Eq. 2. Hence, all asymptotic estimates we derive for Eq. 2 are particularly relevant to NPDR. Since the distance metric given by Eq. 1 is a function of the one-dimensional projection given by Eq. 2, asymptotic estimates derived for Eq. 2 are implicitly relevant to older nearest-neighbor distance-based methods like Relief-based algorithms. We proceed in the following section by applying the Classical Central Limit Theorem and the Delta Method to derive the limit distribution of pairwise distances on any data distribution that is induced by the metric given in Eq. 1.

## 1.1 Asymptotic normality of pairwise distances

Suppose that  $X_{ia}, X_{ja} \stackrel{iid}{\sim} \mathcal{F}_X(\mu_X, \sigma_X^2)$  for two fixed and distinct instances  $i, j \in \mathcal{I}$  and a fixed attribute  $a \in \mathcal{A}$ .  $\mathcal{F}_X$  represents any data distribution with mean  $\mu_X$  and variance  $\sigma_X^2$ .

It is clear that  $|X_{ia} - X_{ja}|^q = |d_{ij}(a)|^q$  is another random variable. Let  $Z_a^q \sim \mathcal{F}_{Z^q}(\mu_{z^q}, \sigma_{z^q}^2)$  be the random variable such that

$$Z_a^q = |d_{ij}(a)|^q = |X_{ia} - X_{ja}|^q, \quad a \in \mathcal{A}. \quad (3)$$

Furthermore, the collection  $\{Z_a^q | a \in \mathcal{A}\}$  is a random sample of size  $p$  of mutually independent random variables. Hence, the sum of  $Z_a^q$  over all  $a \in \mathcal{A}$  is asymptotically

normal by the Classical Central Limit Theorem (CCLT). More explicitly, this implies  
that

$$\left(D_{ij}^{(q)}\right)^q = \sum_{a \in \mathcal{A}} |d_{ij}(a)|^q = \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q = \sum_{a \in \mathcal{A}} Z_a^q \sim \mathcal{N}(\mu_{z^q} p, \sigma_{z^q}^2 p). \quad (4)$$

Consider the smooth function  $g(z) = z^{1/q}$  that is continuously differentiable for  $z > 0$ .  
Assuming that  $\mu_{z^q} > 0$ , the Delta Method [12] can be applied to show that


$$\begin{aligned} g\left(\left(D_{ij}^{(q)}\right)^q\right) &= g\left(\sum_{a \in \mathcal{A}} Z_a^q\right) \\ &= \left(\sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q\right)^{1/q} \\ &= D_{ij}^{(q)} \sim \mathcal{N}\left(g(\mu_{z^q} p), [g'(\mu_{z^q} p)]^2 \sigma_{z^q}^2 p\right) \\ \Rightarrow D_{ij}^{(q)} &\sim \mathcal{N}\left((\mu_{z^q} p)^{1/q}, \frac{\sigma_{z^q}^2 p}{q^2 (\mu_{z^q} p)^{2(1-\frac{1}{q})}}\right). \end{aligned} \quad (5)$$

Therefore, the distance between two fixed, distinct instances  $i$  and  $j$  given by Eq. 1 is  
asymptotically normal. Specifically, when  $q = 2$ , the distribution of  $D_{ij}^{(2)}$  asymptotically  
approaches  $\mathcal{N}\left(\sqrt{\mu_{z^2} p}, \frac{\sigma_{z^2}^2}{4\mu_{z^2}}\right)$ . When  $p$  is small, however, we observe empirically that a  
closer estimate of the sample mean is

$$\begin{aligned} \mathbb{E}\left(D_{ij}^{(2)}\right) &= \sqrt{\mathbb{E}\left[\left(D_{ij}^{(2)}\right)^2\right] - \text{Var}\left(D_{ij}^{(2)}\right)} \\ &= \sqrt{\mu_{z^2} p - \frac{\sigma_{z^2}^2}{4\mu_{z^2}}}. \end{aligned} \quad (6)$$

We estimate rate of convergence to normality for Euclidean ( $q = 2$ ) and Manhattan  
( $q = 1$ ) metrics by comparing the distribution of pairwise distances in simulated data  
to a Gaussian (Fig. 1). We compute the distance between all pairs of instances in  
simulated datasets of uniformly distributed random data. We simulate data with fixed  
 $m = 100$  instances, and, by varying the number of attributes ( $p = 10, 100, 10000$ ), we  
observe rapid convergence to Gaussian. For  $p$  as low as 10 attributes, Gaussian is a  
good approximation. The number of attributes in bioinformatics data is typically quite  
large, at least on the order of  $10^3$ . The Euclidean metric has stronger convergence to  
a Gaussian than Manhattan. This may be due to Euclidean's use of the square root,  
which is a common transformation of data in statistics. Normality was assessed using  
the Shapiro-Wilk test.

To show asymptotic normality of distances, we did not specify whether the data  
distribution  $\mathcal{F}_X$  was discrete or continuous. This is because asymptotic normality is a  
general phenomenon in high attribute dimension  $p$  for any data distribution  $\mathcal{F}_X$  satisfying  
the assumptions we have made. Therefore, Fig. 1 has an analogous representation for  
discrete data, as well as all other continuous data distributions.



**Fig 1.** Convergence to Gaussian for Manhattan and Euclidean distances for simulated standard uniform data with  $m = 100$  instances and  $p = 10, 100$ , and  $10000$  attributes. Convergence to Gaussian occurs rapidly with increasing  $p$ , and Gaussian is a good approximation for  $p$  as low as 10 attributes. The number of attributes in bioinformatics data is typically much larger, at least on the order of  $10^3$ . The Euclidean metric has stronger convergence to normal than Manhattan. P values from Shapiro-Wilk test, where the null hypothesis is a Gaussian distribution.

For distance based learning methods, all pairwise distances are used to determine relative importances for attributes. The collection of all distances above the diagonal in an  $m \times m$  distance matrix does not satisfy the independence assumption used in the previous derivations. This is because of the redundancy that is inherent to the distance matrix calculation. However, this collection is still asymptotically normal with mean and variance approximately equal to those given in Eq. 5. In the next section, we assume actual data distributions in order to define more specific general formulas for standard  $L_q$  and max-min normalized  $L_q$  metrics. We also derive asymptotic moments for a new discrete metric in GWAS data and a new metric for time series correlation-based data, such as, resting-state fMRI.

## 2 $L_q$ metric moments for continuous data distributions

In this section, we begin by deriving general formulas for asymptotic means and variances of the  $L_q$  distance given by Eq. 1 for standard normal and standard uniform data. With our general formulas for continuous data, we compute moments associated with Manhattan ( $L_1$ ) and Euclidean ( $L_2$ ). We then consider the max-min normalized version of the  $L_q$  distance, where the magnitude difference given by Eq. 2 is divided by the range of each feature  $a$ . Using Extreme Value Theory (EVT), we derive formulas for the moments of feature range in standard normal and standard uniform data. Transitioning into discrete data distributions relevant to GWAS, we derive asymptotic moments for two well known metrics and one new metric. In addition, we derive distance asymptotics for time series correlation-based data, such as, resting-state fMRI.

### 2.1 Distribution of $|\mathbf{d}_{ij}(a)|^q = |X_{ia} - X_{ja}|^q$

Suppose that  $X_{ia}, X_{ja} \stackrel{iid}{\sim} \mathcal{F}_X(\mu_x, \sigma_x^2)$  and define  $Z_a^q = |\mathbf{d}_{ij}(a)|^q = |X_{ia} - X_{ja}|^q$ , where  $a \in \mathcal{A}$  and  $|\mathcal{A}| = p$ . In order to find the distribution of  $Z_a^q$ , we will use the following theorem given in [13].

**Theorem 2.1** *Let  $f(x)$  be the value of the probability density of the continuous random variable  $X$  at  $x$ . If the function given by  $y = u(x)$  is differentiable and either increasing or decreasing for all values within the range of  $X$  for which  $f(x) \neq 0$ , then, for these values of  $x$ , the equation  $y = u(x)$  can be uniquely solved for  $x$  to give  $x = w(y)$ , and for the corresponding values of  $y$  the probability density of  $Y = u(X)$  is given by*

$$g(y) = f[w(y)] \cdot |w'(y)| \quad \text{provided } u'(x) \neq 0$$

Elsewhere,  $g(y) = 0$ .

We have the following cases that result from solving for  $X_{ja}$  in the equation given by  $Z_a^q = |X_{ia} - X_{ja}|^q$ :

- (i) Suppose that  $X_{ja} = X_{ia} - (Z_a^q)^{1/q}$ . Based on the iid assumption for  $X_{ia}$  and  $X_{ja}$ , it follows from Thm. 2.1 that the joint density function  $g^{(1)}$  of  $X_{ia}$  and  $Z_a^q$  is given by

$$\begin{aligned} g^{(1)}(x_{ia}, z_a) &= f_X(x_{ia}, x_{ja}) \left| \frac{\partial x_{ja}}{\partial z_a} \right| \\ &= f_X(x_{ia}) f_X(x_{ja}) \left| \frac{-1}{q} (z_a^q)^{\frac{1}{q}-1} \right| \\ &= \frac{1}{q (z_a^q)^{1-\frac{1}{q}}} f_X(x_{ia}) f_X \left( x_{ia} - (z_a^q)^{1/q} \right), \quad z_a > 0 \end{aligned} \tag{7}$$

The density function  $f_{Z_a^q}^{(1)}$  of  $Z_a^q$  is then defined as

$$\begin{aligned} f_{Z_a^q}^{(1)}(z_a^q) &= \int_{-\infty}^{\infty} g^{(1)}(x_{ia}, z_a^q) dx_{ia} \\ &= \frac{1}{q (z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{\infty} f_X(x_{ia}) f_X \left( x_{ia} - (z_a^q)^{1/q} \right) dx_{ia}, \quad z_a > 0. \end{aligned} \tag{8}$$

(ii) Suppose that  $X_{ja} = X_{ia} + (Z_a^q)^{1/q}$ . Based on the iid assumption for  $X_{ia}$  and  $X_{ja}$ ,  
it follows from Thm. 2.1 that the joint density function  $g^{(2)}$  of  $X_{ia}$  and  $Z_a$  is given  
by

$$\begin{aligned} g^{(2)}(x_{ia}, z_a) &= f_X(x_{ia}, x_{ja}) \left| \frac{\partial x_{ja}}{\partial z_a} \right| \\ &= f_X(x_{ia}) f_X(x_{ja}) \left| \frac{1}{q} (z_a^q)^{\frac{1}{q}-1} \right| \\ &= \frac{1}{q (z_a^q)^{1-\frac{1}{q}}} f_X(x_{ia}) f_X \left( x_{ia} - (z_a^q)^{1/q} \right), \quad z_a > 0. \end{aligned} \quad (9)$$

The density function  $f_{Z_a^q}^{(2)}$  of  $Z_a^q$  is then defined as

$$\begin{aligned} f_{Z_a^q}^{(2)}(z_a^q) &= \int_{-\infty}^{\infty} g^{(2)}(x_{ia}, z_a^q) dx_{ia} \\ &= \frac{1}{q (z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{\infty} f_X(x_{ia}) f_X \left( x_{ia} + (z_a^q)^{1/q} \right) dx_{ia}, \quad z_a > 0. \end{aligned} \quad (10)$$

Let  $F_{Z_a^q}$  denote the distribution function of the random variable  $Z_a^q$ . Furthermore,  
we define the events  $E^{(1)}$  and  $E^{(2)}$  as

$$E^{(1)} = \{|X_{ia} - X_{ja}|^q \leq z_a^q | X_{ja} = X_{ia} - (Z_a^q)^{1/q}\} \quad (11)$$

and

$$E^{(2)} = \{|X_{ia} - X_{ja}|^q \leq z_a^q | X_{ja} = X_{ia} + (Z_a^q)^{1/q}\}. \quad (12)$$

Then it follows from fundamental rules of probability that

$$\begin{aligned} F_{Z_a^q}(z_a^q) &= \mathbb{P}[Z_a^q \leq z_a^q] \\ &= \mathbb{P}[|X_{ia} - X_{ja}|^q \leq z_a^q] \\ &= \mathbb{P}[E^{(1)} \cup E^{(2)}] \\ &= \mathbb{P}[E^{(1)}] + \mathbb{P}[E^{(2)}] - \mathbb{P}[E^{(1)} \cap E^{(2)}] \\ &= \mathbb{P}[E^{(1)}] + \mathbb{P}[E^{(2)}] \\ &= \int_{-\infty}^{z_a^q} f_{Z_a^q}^{(1)}(t) dt + \int_{-\infty}^{z_a^q} f_{Z_a^q}^{(2)}(t) dt \\ &= \int_{-\infty}^{z_a^q} \left( f_{Z_a^q}^{(1)}(t) + f_{Z_a^q}^{(2)}(t) \right) dt \\ &= \frac{1}{q (z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{z_a^q} \left( \int_{-\infty}^{\infty} f_X(x_{ia}) [f_X(x_{ia} - t) + f_X(x_{ia} + t)] dx_{ia} \right) dt, \quad z_a > 0. \end{aligned} \quad (13)$$

It follows directly from the result in Eq. 13 that the density function of the random  
variable  $Z_a^q$  is given by

$$\begin{aligned} f_{Z_a^q}(z_a^q) &= \frac{\partial}{\partial z_a^q} F_{Z_a^q}(z_a^q) \\ &= \frac{1}{q (z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{\infty} f_X(x_{ia}) \left[ f_X \left( x_{ia} - (z_a^q)^{1/q} \right) + f_X \left( x_{ia} + (z_a^q)^{1/q} \right) \right] dx_{ia}, \end{aligned} \quad (14)$$

where  $z_a > 0$ .

Using Eq. 14, we can compute the mean and variance of the random variable  $Z_a^q$  as 182

$$\mu_{z^q} = \int_{-\infty}^{\infty} z_a^q f_{Z^q}(z_a^q) dz_a^q \quad (15)$$

and 183

$$\sigma_{z^q}^2 = \int_{-\infty}^{\infty} (z_a^q)^2 f_{Z^q}(z_a^q) dz_a^q - \mu_{z^q}^2. \quad (16)$$

It follows immediately from Eqs. 15 and 16 and the Classical Central Limit Theorem (CCLT) that 184  
185

$$\left(D_{ij}^{(q)}\right)^q = \sum_{a \in \mathcal{A}} Z_a^q = \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q \sim \mathcal{N}(\mu_{z^q} p, \sigma_{z^q}^2 p). \quad (17)$$

Applying the result given in Eq. 5, the distribution of  $D_{ij}^{(q)}$  is given by 186

$$D_{ij}^{(q)} \sim \mathcal{N}\left((\mu_{z^q} p)^{1/q}, \frac{\sigma_{z^q}^2 p}{q^2 (\mu_{z^q} p)^{2(1-\frac{1}{q})}}\right), \quad \mu_{z^q} > 0 \quad (18)$$

with improved estimate of the mean for  $q = 2$  given by Eq. 6. 187

### 2.1.1 Standard normal data 188

If  $X_{ia}, X_{ja} \stackrel{iid}{\sim} \mathcal{N}(0, 1)$ , then the marginal density functions with respect to  $X$  for  $X_{ia}$ ,  $X_{ia} - (Z_a^q)^{1/q}$ , and  $X_{ia} + (Z_a^q)^{1/q}$  are defined as 189  
190

$$f_X(x_{ia}) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_{ia}^2}, \quad (19)$$

$$f_X\left(x_{ia} - (z_a^q)^{1/q}\right) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(x_{ia} - (z_a^q)^{1/q})^2}, \quad z_a > 0, \text{ and} \quad (20)$$

$$f_X\left(x_{ia} + (z_a^q)^{1/q}\right) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(x_{ia} + (z_a^q)^{1/q})^2}, \quad z_a > 0. \quad (21)$$

Substituting the results given by Eqs. 19-21 into Eq. 14 and completing the square on  $x_{ia}$  in the exponents, we have

$$f_{Z^q}(z_a^q) = \frac{1}{2q\pi (z_a^q)^{1-\frac{1}{q}}} e^{-\frac{1}{4}(z_a^q)^{2/q}} \int_{-\infty}^{\infty} \left( e^{-\frac{1}{2}[\sqrt{2}x_{ia} - \frac{\sqrt{2}}{2}(z_a^q)^{1/q}]^2} + e^{-\frac{1}{2}[\sqrt{2}x_{ia} + \frac{\sqrt{2}}{2}(z_a^q)^{1/q}]^2} \right) dx_{ia} \quad (22)$$

$$= \frac{1}{2q\sqrt{\pi} (z_a^q)^{1-\frac{1}{q}}} e^{-\frac{1}{4}(z_a^q)^{2/q}} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \left( e^{-\frac{1}{2}u^2} + e^{-\frac{1}{2}u^2} \right) du \quad (23)$$

$$= \frac{1}{2q\sqrt{\pi} (z_a^q)^{1-\frac{1}{q}}} e^{-\frac{1}{4}(z_a^q)^{2/q}} (1 + 1) \quad (24)$$

$$= \frac{1}{q\sqrt{\pi}} (z_a^q)^{\frac{1}{q}-1} e^{-\frac{1}{4}(z_a^q)^{2/q}} \quad (25)$$

$$= \frac{\frac{2}{q}}{(2q)^{1/q} \Gamma\left(\frac{1}{q}\right)} (z_a^q)^{\frac{1}{q}-1} e^{-\left(\frac{z_a^q}{2q}\right)^{2/q}}. \quad (26)$$



The density function given by Eq. 22 is a Generalized Gamma density with parameters  $b = \frac{2}{q}$ ,  $c = 2^q$ , and  $d = \frac{1}{q}$ . This distribution has mean and variance given by

$$\begin{aligned}\mu_{z^q} &= \frac{c\Gamma\left(\frac{d+1}{b}\right)}{\Gamma\left(\frac{d}{b}\right)} \\ &= \frac{2^q\Gamma\left(\frac{q+1}{2}\right)}{\sqrt{\pi}}\end{aligned}\quad (27)$$

and

$$\begin{aligned}\sigma_{z^q}^2 &= c^2 \left[ \frac{\Gamma\left(\frac{d+2}{b}\right)}{\Gamma\left(\frac{d}{b}\right)} - \left( \frac{\Gamma\left(\frac{d+1}{b}\right)}{\Gamma\left(\frac{d}{b}\right)} \right)^2 \right] \\ &= 4^q \left[ \frac{\Gamma\left(q + \frac{1}{2}\right)}{\sqrt{\pi}} - \frac{\Gamma^2\left(\frac{1}{2}q + \frac{1}{2}\right)}{\pi} \right].\end{aligned}\quad (28)$$

By linearity of the expected value and variance operators under the iid assumption, Eqs. 27 and 28 allow the  $p$ -dimensional mean and variance of the  $D_{ij}^{(q)}$  distribution to be computed directly as

$$\mu_{(D_{ij}^{(q)})^q} = \mathbb{E} \left[ (D_{ij}^{(q)})^q \right] = \mathbb{E} \left( \sum_{a \in \mathcal{A}} Z_a^q \right) = \sum_{a \in \mathcal{A}} \mathbb{E}(Z_a^q) = \sum_{a \in \mathcal{A}} \frac{2^q\Gamma\left(\frac{q+1}{2}\right)}{\sqrt{\pi}} = \frac{2^q\Gamma\left(\frac{q+1}{2}\right)}{\sqrt{\pi}} p \quad (29)$$

and

$$\begin{aligned}\sigma_{(D_{ij}^{(q)})^q}^2 &= \text{Var} \left[ (D_{ij}^{(q)})^q \right] = \text{Var} \left( \sum_{a \in \mathcal{A}} Z_a^q \right) \\ &= \sum_{a \in \mathcal{A}} \text{Var}(Z_a^q) \\ &= \sum_{a \in \mathcal{A}} 4^q \left[ \frac{\Gamma\left(q + \frac{1}{2}\right)}{\sqrt{\pi}} - \frac{\Gamma^2\left(\frac{1}{2}q + \frac{1}{2}\right)}{\pi} \right] \\ &= 4^q \left[ \frac{\Gamma\left(q + \frac{1}{2}\right)}{\sqrt{\pi}} - \frac{\Gamma^2\left(\frac{1}{2}q + \frac{1}{2}\right)}{\pi} \right] p.\end{aligned}\quad (30)$$

Therefore, the asymptotic distribution of  $D_{ij}^{(q)}$  for standard normal data is

$$\mathcal{N} \left( \left( 2^q \frac{\Gamma\left(\frac{q+1}{2}\right)}{\sqrt{\pi}} p \right)^{1/q}, \frac{4^q p}{q^2 \left( \frac{2^q \Gamma\left(\frac{1}{2}q + \frac{1}{2}\right)}{\sqrt{\pi}} p \right)^{2(1-\frac{1}{q})}} \left[ \frac{\Gamma\left(q + \frac{1}{2}\right)}{\sqrt{\pi}} - \frac{\Gamma^2\left(\frac{1}{2}q + \frac{1}{2}\right)}{\pi} \right] \right). \quad (31)$$

### 2.1.2 Standard uniform data

If  $X_{ia}, X_{ja} \stackrel{iid}{\sim} \mathcal{U}(0, 1)$ , then the marginal density functions with respect to  $X$  for  $X_{ia}$ ,  $X_{ia} - (Z_a^q)^{1/q}$ , and  $X_{ia} + (Z_a^q)^{1/q}$  are defined as

$$f_X(x_{ia}) = 1, \quad 0 \leq x_{ia} \leq 1 \quad (32)$$

$$f_X\left(x_{ia} - (z_a^q)^{1/q}\right) = 1, \quad 0 \leq x_{ia} - (z_a^q)^{1/q} \leq 1, \text{ and} \quad (33)$$

$$f_X \left( x_{ia} + (z_a^q)^{1/q} \right) = 1, \quad 0 \leq x_{ia} + (z_a^q)^{1/q} \leq 1. \quad (34)$$

Substituting the results given by Eqs. 32-34 into Eq. 14, we have

$$\begin{aligned} f_{Z^q}(z_a^q) &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{\infty} f_X(x_{ia}) \left[ f_X \left( x_{ia} - (z_a^q)^{1/q} \right) + f_X \left( x_{ia} + (z_a^q)^{1/q} \right) \right] dx_{ia}, \\ & \quad 0 < z_a \leq 1 \\ &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_0^1 [f_X(x_{ia} - (z_a^q)^{1/q}) + f_X(x_{ia} + (z_a^q)^{1/q})] dx_{ia}, \quad 0 < z_a \leq 1 \\ &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_{(z_a^q)}^1 1 dx_{ia} + \int_0^{1-(z_a^q)} 1 dx_{ia}, \quad 0 < z_a \leq 1 \\ &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} [(1 - (z_a^q)) + (1 - (z_a^q))], \quad 0 < z_a \leq 1 \\ &= \frac{1}{q} \cdot 2 (z_a^q)^{\frac{1}{q}-1} \left[ 1 - (z_a^q)^{1/q} \right]^{2-1}, \quad 0 < z_a \leq 1. \end{aligned} \quad (35)$$

The density given by Eq. 35 is a Kumaraswamy density with parameters  $b = \frac{1}{q}$  and  $c = 2$  with moment generating function (MGF) given by

$$\begin{aligned} M_n &= \frac{c\Gamma(1 + \frac{n}{b})\Gamma(c)}{\Gamma(1 + c + \frac{n}{b})} \\ &= \frac{2}{(nq + 2)(nq + 1)}. \end{aligned} \quad (36)$$

Using the MGF given by Eq. 36, the mean and variance of  $Z_a^q$  are computed as

$$\mu_{z^q} = \frac{2}{(q + 2)(q + 1)} \quad (37)$$

and

$$\sigma_{z^q}^2 = \frac{1}{(q + 1)(2q + 1)} - \left( \frac{2}{(q + 2)(q + 1)} \right)^2. \quad (38)$$

By linearity of the expected value and variance operators under the iid assumption, Eqs. 39 and 40 allow the  $p$ -dimensional mean and variance of the  $(D_{ij}^{(q)})^q$  distribution to be computed directly as

$$\begin{aligned} \mu_{(D_{ij}^{(q)})^q} &= E \left[ (D_{ij}^{(q)})^q \right] = E \left( \sum_{a \in \mathcal{A}} Z_a^q \right) \\ &= \sum_{a \in \mathcal{A}} E(Z_a^q) \\ &= \sum_{a \in \mathcal{A}} \frac{2}{(q + 2)(q + 1)} \\ &= \frac{2p}{(q + 2)(q + 1)} \end{aligned} \quad (39)$$

and

214

$$\begin{aligned}
\sigma^2_{(D_{ij}^{(q)})^q} &= \text{Var} \left[ \left( D_{ij}^{(q)} \right)^q \right] = \text{Var} \left( \sum_{a \in \mathcal{A}} Z_a^q \right) \\
&= \sum_{a \in \mathcal{A}} \text{Var} (Z_a^q) \\
&= \sum_{a \in \mathcal{A}} \left[ \frac{1}{(q+1)(2q+1)} - \left( \frac{2}{(q+2)(q+1)} \right)^2 \right] \\
&= \left[ \frac{1}{(q+1)(2q+1)} - \left( \frac{2}{(q+2)(q+1)} \right)^2 \right] p.
\end{aligned} \tag{40}$$

Therefore, the asymptotic distribution of  $D_{ij}^{(q)}$  for standard uniform data is

215

$$\begin{aligned}
&\mathcal{N} \left( \left( \frac{2p}{(q+2)(q+1)} \right)^{1/q}, \right. \\
&\quad \left. \frac{p}{q^2 \left( \frac{2p}{(q+2)(q+1)} \right)^{2(1-\frac{1}{q})}} \left[ \frac{1}{(q+1)(2q+1)} - \left( \frac{2}{(q+2)(q+1)} \right)^2 \right] \right).
\end{aligned} \tag{41}$$

## 2.2 Manhattan ( $L_1$ )

216

With our general formulas for the asymptotic mean and variance given by Eqs. 31 and 41 for any value of  $q \in \mathbb{N}$ , we can simply substitute a particular value of  $q$  in order to determine the asymptotic distribution of the corresponding distance metric  $D_{ij}^{(q)}$ . We demonstrate this with the example of the Manhattan metric for standard normal and standard uniform data, which is given by Eq. 1 by setting  $q = 1$  ( $L_1$ ).

217

218

219

220

221

### 2.2.1 Standard normal data

222

Using the mean given by Eq. 31 and substituting  $q = 1$ , we have the following for expected  $L_1$  distance between two independently sample instances  $i$  and  $j$  in standard normal data

223

224

225

$$\begin{aligned}
\mathbb{E} \left( D_{ij}^{(1)} \right) &= \left( 2 \frac{\Gamma \left( \frac{1+1}{2} \right)}{\sqrt{\pi}} p \right)^{1/1} \\
&= \frac{2p}{\sqrt{\pi}} \Gamma(1) \\
&= \frac{2p}{\sqrt{\pi}}.
\end{aligned} \tag{42}$$

We see in Eq. 42 that  $D_{ij}^{(1)} \sim p$  in the limit, which implies that this distance is unbounded as feature dimension  $p$  increases.

226

227

Substituting  $q = 1$  into the formula for the asymptotic variance of  $D_{ij}^{(1)}$  given in

228

Eq. 31 leads to the following

229

$$\begin{aligned}
\text{Var} \left( D_{ij}^{(1)} \right) &= \frac{4^1 p}{1^2 \left( \frac{2^1 \Gamma(\frac{1}{2}(1) + \frac{1}{2})}{\sqrt{\pi}} p \right)^{2(1-\frac{1}{1})}} \left[ \frac{\Gamma(1 + \frac{1}{2})}{\sqrt{\pi}} - \frac{\Gamma^2(\frac{1}{2}(1) + \frac{1}{2})}{\pi} \right] \\
&= \frac{4p}{1} \left[ \frac{\frac{1}{2} \Gamma(\frac{1}{2})}{\sqrt{\pi}} - \frac{\Gamma^2(1)}{\pi} \right] \\
&= 4p \left[ \frac{1}{2} - \frac{1}{\pi} \right] \\
&= \frac{2(\pi - 2)p}{\pi}.
\end{aligned} \tag{43}$$

Similar to the mean given by Eq. 42, the limiting variance of  $D_{ij}^{(1)}$  given by Eq. 43 grows on the order of feature dimension  $p$ , which implies that points become more dispersed as the dimension increases.

230  
231  
232

## 2.2.2 Standard uniform data

233

Using the mean given by Eq. 41 and substituting  $q = 1$ , we have the following for the expected  $L_1$  distance between two independently sampled instances  $i$  and  $j$  in standard uniform data

234  
235  
236

$$\begin{aligned}
\mathbb{E} \left( D_{ij}^{(1)} \right) &= \left( \frac{2p}{(1+2)(1+1)} \right)^{1/1} \\
&= \frac{2p}{6} \\
&= \frac{p}{3}.
\end{aligned} \tag{44}$$

Once again, we see that the mean of  $D_{ij}^{(1)}$  given by Eq. 44 grows on the order of  $p$  just as in the case of standard normal data.

237  
238

Substituting  $q = 1$  into the formula given by Eq. 41 of the asymptotic variance of  $D_{ij}^{(1)}$  leads to the following

239  
240

$$\begin{aligned}
\text{Var} \left( D_{ij}^{(1)} \right) &= \frac{p}{1^2 \left( \frac{2p}{(1+2)(1+1)} \right)^{2(1-\frac{1}{1})}} \left[ \frac{1}{(1+1)(2(1)+1)} - \left( \frac{2}{(1+2)(1+1)} \right)^2 \right] \\
&= p \left[ \frac{1}{6} - \frac{1}{9} \right] \\
&= \frac{p}{18}.
\end{aligned} \tag{45}$$

As in the case of the  $L_1$  metric on standard normal data, we have a variance given by Eq. 45 that grows on the order of  $p$ . The distances between points in high-dimensional uniform data become more widely dispersed with this metric.

241  
242  
243

## 2.3 Euclidean ( $L_2$ )

244

Analogous to the previous section, we demonstrate the usage of Eqs. 31 and 41 for the Euclidean metric for standard normal and standard uniform data, which is given by Eq. 1 by setting  $q = 2$  ( $L_2$ ).

245  
246  
247

### 2.3.1 Standard normal data

Using the mean given by Eq. 31 and substituting  $q = 2$ , we have the following for expected  $L_2$  distance between two independently sampled instances  $i$  and  $j$  in standard normal data

$$\begin{aligned} \mathbb{E} \left( D_{ij}^{(2)} \right) &= \left( 2 \frac{\Gamma \left( \frac{2+1}{2} \right)}{\sqrt{\pi}} p \right)^{1/2} \\ &= \left( \frac{2p}{\sqrt{\pi}} \Gamma \left( \frac{3}{2} \right) \right)^{1/2} \\ &= \sqrt{2p}. \end{aligned} \quad (46)$$

In the case of  $L_2$  on standard normal data, we see that the mean of  $D_{ij}^{(2)}$  given by Eq. 46 grows on the order of  $\sqrt{p}$ . Hence, the Euclidean distance does not increase as quickly as the Manhattan distance on standard normal data.

Substituting  $q = 2$  into the formula for the asymptotic variance of  $D_{ij}^{(2)}$  given in Eq. 31 leads to the following

$$\begin{aligned} \text{Var} \left( D_{ij}^{(2)} \right) &= \frac{4^2 p}{2^2 \left( \frac{2^2 \Gamma \left( \frac{1}{2}(2) + \frac{1}{2} \right)}{\sqrt{\pi}} p \right)^{2(1-\frac{1}{2})}} \left[ \frac{\Gamma \left( 2 + \frac{1}{2} \right)}{\sqrt{\pi}} - \frac{\Gamma^2 \left( \frac{1}{2}(2) + \frac{1}{2} \right)}{\pi} \right] \\ &= \frac{16p}{4 \left( \frac{4\Gamma \left( \frac{3}{2} \right)}{\sqrt{\pi}} p \right)} \left[ \frac{\Gamma \left( \frac{5}{2} \right)}{\sqrt{\pi}} - \frac{\Gamma^2 \left( \frac{3}{2} \right)}{\pi} \right] \\ &= 2 \left[ \frac{3}{4} - \frac{1}{4} \right] \\ &= 1. \end{aligned} \quad (47)$$

Surprisingly, the asymptotic variance given by Eq. 47 is just 1. Regardless of data dimensions  $m$  and  $p$ , the variance of Euclidean distances on standard normal data tends to 1. Therefore, most instances are contained within a ball of radius 1 about the mean in high feature dimension  $p$ . This means that the Euclidean distance distribution on standard normal data is simply a horizontal shift to the right of the standard normal distribution.

For the case in which the number of attributes  $p$  is small, an improved estimate of the mean is given by Eq. 6. The lower dimensional estimate of the mean is as follows

$$\begin{aligned} \mathbb{E} \left( D_{ij}^{(2)} \right) &= \left( 2 \frac{\Gamma \left( \frac{2+1}{2} \right)}{\sqrt{\pi}} p - 1 \right)^{1/2} \\ &= \left( \frac{2p}{\sqrt{\pi}} \Gamma \left( \frac{3}{2} \right) - 1 \right)^{1/2} \\ &= \sqrt{2p - 1}. \end{aligned} \quad (48)$$

For high dimensional data sets like gene expression [14, 15], which typically contain thousands of genes (or features), it is clear that the magnitude of  $p$  will be sufficient to use Eq. 46 since  $\sqrt{2p} \approx \sqrt{2p - 1}$  in that case.

### 2.3.2 Standard uniform data

Using the mean given by Eq. 41 and substituting  $q = 2$ , we have the following for expected  $L_2$  distance between two independently sampled instances  $i$  and  $j$  in standard

$$\begin{aligned}
E(D_{ij}^{(2)}) &= \left( \frac{2p}{(2+2)(2+1)} \right)^{1/2} \\
&= \left( \frac{2p}{12} \right)^{1/2} \\
&= \sqrt{\frac{p}{6}}.
\end{aligned} \tag{49}$$

As in the case of standard normal data, the expected value of  $D_{ij}^{(2)}$  given by Eq. 49 grows on the order of  $\sqrt{p}$ .

Substituting  $q = 2$  into the formula for the asymptotic variance of  $D_{ij}^{(2)}$  given in Eq. 41 leads to the following

$$\begin{aligned}
\text{Var}(D_{ij}^{(2)}) &= \frac{p}{2^2 \left( \frac{2p}{(2+2)(2+1)} \right)^{2(1-\frac{1}{2})}} \left[ \frac{1}{(2+1)(2(2)+1)} - \left( \frac{2}{(2+2)(2+1)} \right)^2 \right] \\
&= \frac{3}{2} \left[ \frac{1}{15} - \frac{1}{36} \right] \\
&= \frac{7}{120}.
\end{aligned} \tag{50}$$

Once again, the variance of Euclidean distance surprisingly approaches a constant.

For the case in which the number of attributes  $p$  is small, an improved estimate of the mean is given by Eq. 6. The lower dimensional estimate of the mean is as follows

$$\begin{aligned}
E(D_{ij}^{(2)}) &= \left( \frac{2p}{(2+2)(2+1)} - \frac{7}{120} \right)^{1/2} \\
&= \left( \frac{2p}{12} - \frac{7}{120} \right)^{1/2} \\
&= \sqrt{\frac{p}{6} - \frac{7}{120}}.
\end{aligned} \tag{51}$$

This concludes our analysis with continuous data distributions and the standard  $L_q$  metric. In the next section, we will use extreme value theory to derive the distribution of the sample maximum and minimum for standard normal and standard uniform data. This will lead us to asymptotics for the max-min normalized  $L_q$  metric used frequently in Relief-based algorithms [?, 1, 3] for scoring features.

## 2.4 Distribution of max-min normalized $L_q$ metric

For Relief-based methods [?, 1, 3], the standard numeric diff metric is given by

$$d_{ij}^{\text{num}}(a) = \text{diff}(a, (i, j)) = \frac{|X_{ia} - X_{ja}|}{\max(a) - \min(a)}, \tag{52}$$

where  $\max(a) = \max_{k \in \mathcal{I}} \{X_{ka}\}$ ,  $\min(a) = \min_{k \in \mathcal{I}} \{X_{ka}\}$ , and  $\mathcal{I} = \{1, 2, \dots, m\}$ .

In order to determine moments of asymptotic distance distributions induced by Eq. 52, we must first derive the asymptotic extreme value distributions of the attribute maximum and minimum. Although the exact distribution of the maximum or minimum requires an assumption about the data distribution, the Fisher-Tippett-Gnedenko Theorem allows us to generally categorize the extreme value distribution for a collection of independent and identically distributed random variables into one of three distributional families. Before stating the theorem, we first need the following definition

**Definition 2.1** A distribution  $\mathcal{F}_X$  is said to be **degenerate** if its density function  $f_X$  is the Dirac delta  $\delta(x - c_0)$  centered at a constant  $c_0 \in \mathbb{R}$ , with corresponding distribution function  $F_X$  defined as

$$F_X(x) = \begin{cases} 1, & x \geq c_0, \\ 0, & x < c_0. \end{cases}$$

**Theorem 2.2 (Fisher-Tippett-Gnedenko)** Let  $X_{1a}, X_{2a}, \dots, X_{ma} \stackrel{iid}{\sim} \mathcal{F}_X(\mu_x, \sigma_x^2)$  and let  $X_a^{max} = \max_{k \in \mathcal{I}} \{X_{ka}\}$ . If there exists two non-random sequences  $b_m > 0$  and  $c_m$  such that

$$\lim_{m \rightarrow \infty} P\left(\frac{X_a^{max} - c_m}{b_m} \leq x\right) = G_X(x),$$

where  $G_X$  is a non-degenerate distribution function, then the limiting distribution  $\mathcal{G}_X$  is in the Gumbel, Fréchet, or Weibull family.

The three distribution families given in Thm. 2.2 are actually special cases of the Generalized Extreme Value Distribution. In the context of extreme values, Thm. 2.2 is analogous to the Central Limit Theorem for the distribution of sample mean. Although we will not explicitly invoke this theorem, it does tell us something very important about the asymptotic behavior of sample extremes under certain necessary conditions. For illustration of this general phenomenon of sample extremes, we derive the distribution of the maximum for standard normal data to show that the limiting distribution is in the Gumbel family, which is a well known result. In the case of standard uniform data, we will derive the distribution of the maximum and minimum directly. Regardless of data type, the distribution of the sample maximum is derived as follows

$$\begin{aligned} P[X_a^{max} \leq x] &= P\left[\max_{k \in \mathcal{I}} \{X_{ka}\} \leq x\right] \\ &= P[X_{1a} \leq x, X_{2a} \leq x, \dots, X_{ma} \leq x] \\ &= \prod_{k=1}^m P[X_{ka} \leq x] \\ &= \prod_{k=1}^m F_X(x) \\ &= [F_X(x)]^m. \end{aligned} \tag{53}$$

Therefore, we have the following expression for the distribution function of the maximum

$$F_{\max}(x) = [F_X(x)]^m. \tag{54}$$

Differentiating the distribution function given by Eq. 54 gives us the following density function for the distribution of the maximum

$$\begin{aligned} f_{\max}(x) &= \frac{d}{dx} F_{\max}(x) \\ &= \frac{d}{dx} [F_X(x)]^m \\ &= m[F_X(x)]^{m-1} f_X(x). \end{aligned} \tag{55}$$

The distribution of the sample minimum,  $X_a^{\min}$ , is derived as follows

316

$$\begin{aligned}
P[X_a^{\min} \leq x] &= 1 - P[X_a^{\min} \geq x] \\
&= 1 - P\left[\min_{k \in \mathcal{I}} \{X_{ka}\} \geq x\right] \\
&= 1 - P[X_{1a} \geq x, X_{2a} \geq x, \dots, X_{ma} \geq x] \\
&= 1 - \prod_{k=1}^m P[X_{ka} \geq x] \\
&= 1 - [P[X_{1a} \geq x]]^m \\
&= 1 - [1 - P[X_{1a} \leq x]]^m \\
&= 1 - [1 - F_X(x)]^m.
\end{aligned} \tag{56}$$

Therefore, we have the following expression for the distribution function of the minimum

317

$$F_{\min}(x) = 1 - [1 - F_X(x)]^m. \tag{57}$$

318

Differentiating the distribution function given by Eq. 57 gives us the following density function for the distribution of the minimum

319

320

$$\begin{aligned}
f_{\min}(x) &= \frac{d}{dx} F_{\min}(x) \\
&= \frac{d}{dx} (1 - [1 - F_X(x)]^m) \\
&= m [1 - F_X(x)]^{m-1} f_X(x).
\end{aligned} \tag{58}$$

Given the densities of the distribution of sample maximum and minimum, we can easily compute moments and the variance. The first and second moment about the origin and the variance of the distribution of the maximum are given by the following

321

322

323

$$\begin{aligned}
\mu_{\max}^{(1)}(m) &= E(X_a^{\max}) = \int_{-\infty}^{\infty} x f_{\max}(x) dx \\
&= \int_{-\infty}^{\infty} x (m[F_X(x)]^{m-1} f_X(x)) dx \\
&= m \int_{-\infty}^{\infty} x f_X(x) [F_X(x)]^{m-1} dx.
\end{aligned} \tag{59}$$

324

$$\begin{aligned}
\mu_{\max}^{(2)}(m) &= E[(X_a^{\max})^2] = \int_{-\infty}^{\infty} x^2 f_{\max}(x) dx \\
&= \int_{-\infty}^{\infty} x^2 (m[F_X(x)]^{m-1} f_X(x)) dx \\
&= m \int_{-\infty}^{\infty} x^2 f_X(x) [F_X(x)]^{m-1} dx
\end{aligned} \tag{60}$$

325

$$\sigma_{\max}^2(m) = \mu_{\max}^{(2)}(m) - [\mu_{\max}^{(1)}(m)]^2 \tag{61}$$

Similarly, we have the first and second moment about the origin and variance of the distribution of sample minimum given by the following

326

327

$$\begin{aligned}
\mu_{\min}^{(1)}(m) &= E(X_a^{\min}) = \int_{-\infty}^{\infty} x f_{\min}(x) dx \\
&= \int_{-\infty}^{\infty} x (m[F_X(x)]^{m-1} f_X(x)) dx \\
&= m \int_{-\infty}^{\infty} x f_X(x) [F_X(x)]^{m-1} dx,
\end{aligned} \tag{62}$$



$$\begin{aligned}
\mu_{\min}^{(2)}(m) &= E[(X_a^{\min})^2] = \int_{-\infty}^{\infty} x^2 f_{\min}(x) dx \\
&= \int_{-\infty}^{\infty} x^2 (m[F_X(x)]^{m-1} f_X(x)) dx \\
&= m \int_{-\infty}^{\infty} x^2 f_X(x) [F_X(x)]^{m-1} dx,
\end{aligned} \tag{63}$$

and

$$\sigma_{\min}^2(m) = \mu_{\min}^{(2)}(m) - \left[ \mu_{\min}^{(1)}(m) \right]^2. \tag{64}$$

With the densities of attribute maximum and minimum for sample size  $m$ , the expected range is given by the following

$$\begin{aligned}
E(X_a^{\max} - X_a^{\min}) &= E(X_a^{\max}) - E(X_a^{\min}) \\
&= \mu_{\max}^{(1)}(m) - \mu_{\min}^{(1)}(m).
\end{aligned} \tag{65}$$

For a data distribution that has zero skewness and has support that is symmetric about 0, the result given by Eq. 65 can be simplified to the following expression

$$E(X_a^{\max} - X_a^{\min}) = 2\mu_{\max}^{(1)}(m). \tag{66}$$

For large samples ( $m \gg 1$ ), the covariance between the sample maximum and minimum is approximately zero [16]. Therefore, the variance of the attribute range of a sample of size  $m$  is given by the following

$$\begin{aligned}
\text{Var}(X_a^{\max} - X_a^{\min}) &\approx \text{Var}(X_a^{\max}) + \text{Var}(X_a^{\min}) \\
&= \sigma_{\max}^2(m) + \sigma_{\min}^2(m).
\end{aligned} \tag{67}$$

Under the assumption of zero skewness and support that is symmetric about 0, the result given by Eq. 67 becomes the following

$$\begin{aligned}
\text{Var}(X_a^{\max} - X_a^{\min}) &= 2\text{Var}(X_a^{\max}) \\
&= 2\sigma_{\max}^2.
\end{aligned} \tag{68}$$

Let  $\mu_{D_{ij}^{(q)}}$  and  $\sigma_{D_{ij}^{(q)}}^2$  denote the mean and variance given by Eq. 18. Furthermore, let  $D_{ij}^{(q*)}$  denote the max-min normalized distance between instances  $i$  and  $j$  that is induced by the metric given by Eq. 52. Then the mean of the max-min normalized distance distribution is given by the following

$$\begin{aligned}
\mu_{D_{ij}^{(q*)}} &= E \left[ \left( \sum_{a \in \mathcal{A}} \left( \frac{|X_{ia} - X_{ja}|}{X_a^{\max} - X_a^{\min}} \right)^q \right)^{1/q} \right] \\
&\approx \frac{1}{E(X_a^{\max} - X_a^{\min})} E \left[ \left( \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q \right)^{1/q} \right] \\
&= \frac{\mu_{D_{ij}^{(q)}}}{E(X_a^{\max}) - E(X_a^{\min})} \\
&= \frac{\mu_{D_{ij}^{(q)}}}{\mu_{\max}^{(1)} - \mu_{\min}^{(1)}}.
\end{aligned} \tag{69}$$

The variance of the max-min normalized distance distribution is given by the following 342

$$\begin{aligned}
\sigma_{D_{ij}^{(q*)}}^2 &= \text{Var} \left[ \left( \sum_{a \in \mathcal{A}} \left( \frac{|X_{ia} - X_{ja}|}{X_a^{\max} - X_a^{\min}} \right)^q \right)^{1/q} \right] \\
&= \text{E} \left[ \left( \sum_{a \in \mathcal{A}} \left( \frac{|X_{ia} - X_{ja}|}{X_a^{\max} - X_a^{\min}} \right)^q \right)^{2/q} \right] - \left( \text{E} \left[ \left( \sum_{a \in \mathcal{A}} \left( \frac{|X_{ia} - X_{ja}|}{X_a^{\max} - X_a^{\min}} \right)^q \right)^{1/q} \right] \right)^2 \\
&\approx \frac{\text{E} \left[ \left( \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q \right)^{2/q} \right]}{\text{E}[(X_a^{\max} - X_a^{\min})^2]} - \frac{\left( \text{E} \left[ \left( \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q \right)^{1/q} \right] \right)^2}{\text{E}[(X_a^{\max} - X_a^{\min})^2]} \\
&= \frac{\sigma_{D_{ij}^{(q)}}^2 + \mu_{D_{ij}^{(q)}}^2}{\text{E}[(X_a^{\max} - X_a^{\min})^2]} - \frac{\mu_{D_{ij}^{(q)}}^2}{\text{E}[(X_a^{\max} - X_a^{\min})^2]} \\
&= \frac{\sigma_{D_{ij}^{(q)}}^2}{\text{E}[(X_a^{\max} - X_a^{\min})^2]} \\
&= \frac{\sigma_{D_{ij}^{(q)}}^2}{\text{E}[(X_a^{\max})^2] - 2\text{E}(X_a^{\max})\text{E}(X_a^{\min}) + \text{E}(X_a^{\min})^2}} \\
&= \frac{\sigma_{D_{ij}^{(q)}}^2}{\mu_{\max}^{(2)}(m) - 2\mu_{\max}^{(1)}(m)\mu_{\min}^{(1)}(m) + \mu_{\min}^{(2)}(m)}. \tag{70}
\end{aligned}$$

With the results given by Eqs. 69 and 70, we have the following generalized estimate for the asymptotic distribution of the max-min normalized distance distribution 343  
344

$$D_{ij}^{(q*)} \sim \mathcal{N} \left( \frac{\mu_{D_{ij}^{(q)}}}{\mu_{\max}^{(1)}(m) - \mu_{\min}^{(1)}(m)}, \frac{\sigma_{D_{ij}^{(q)}}^2}{\mu_{\max}^{(2)}(m) - 2\mu_{\max}^{(1)}(m)\mu_{\min}^{(1)}(m) + \mu_{\min}^{(2)}(m)} \right). \tag{71}$$

For data with zero skewness and support that is symmetric about 0, the expected sample maximum is the additive inverse of the expected sample minimum. This allows us to express the formula given by Eq. 69 exclusively in terms of the expected maximum. This result is given by the following 345  
346  
347  
348

$$\mu_{D_{ij}^{(q*)}} \approx \frac{\mu_{D_{ij}^{(q)}}}{2\mu_{\max}^{(1)}(m)}. \tag{72}$$

A similar substitution gives us the following expression for the variance of the max-min normalized distance distribution 349  
350

$$\begin{aligned}
\sigma_{D_{ij}^{(q*)}}^2 &\approx \frac{\sigma_{D_{ij}^{(q)}}^2}{2\mu_{\max}^{(2)}(m) + 2 \left[ \mu_{\max}^{(1)}(m) \right]^2} \\
&= \frac{\sigma_{D_{ij}^{(q)}}^2}{2 \left( \sigma_{\max}^2(m) + \left[ \mu_{\max}^{(1)}(m) \right]^2 \right)}. \tag{73}
\end{aligned}$$

Therefore, the asymptotic distribution of the max-min normalized distance distribu- 351

tion is given by the following

352

$$D_{ij}^{(q*)} \sim \mathcal{N} \left( \frac{\mu_{D_{ij}^{(q)}}}{2\mu_{\max}^{(1)}(m)}, \frac{\sigma_{D_{ij}^{(q)}}^2}{2 \left( \sigma_{\max}^2(m) + [\mu_{\max}^{(1)}(m)]^2 \right)} \right). \quad (74)$$

#### 2.4.1 Standard normal data

353

Standard normal data has zero skewness and has support that is symmetric about 0. This implies that the mean and variance of the distribution of sample range can be expressed exclusively in terms of the sample maximum. Given the nature of the density function of the sample maximum for sample size  $m$ , the integration required to determine the moments given by Eqs. 59 and 60 is not possible. These moments can either be approximated numerically or we can use extreme value theory to determine the form of the asymptotic distribution of the sample maximum. Using the latter method, we will show that the asymptotic distribution of the sample maximum for standard normal data is in the Gumbel family. Let  $c_m = -\Phi^{-1} \left( \frac{1}{m} \right)$  and  $b_m = \frac{1}{c_m}$ . Using Taylor's Theorem, we have the following expansion

354

355

356

357

358

359

360

361

362

363

$$\begin{aligned} \log \Phi(-c_m - b_m x) &= \log \Phi(-c_m) - b_m x \frac{\phi(-c_m)}{\Phi(-c_m)} + \mathcal{O}(b_m^2 x^2) \\ &= \log \left( \frac{1}{m} \right) - x \frac{\phi(-c_m)}{c_m \Phi(-c_m)} + \mathcal{O}(b_m^2 x^2). \end{aligned} \quad (75)$$

In order to simplify the right-hand side of Eq. 75, we will use the well known Mills Ratio Bounds [17] given by the following

364

365

$$1 \leq \frac{\phi(x)}{x\Phi(-x)} \leq 1 + \frac{1}{x^2}, \quad x > 0. \quad (76)$$

The inequalities given by Eq. 76 show that  $\frac{\phi(x)}{x\Phi(-x)} \rightarrow 1$  as  $x \rightarrow \infty$ . This implies that  $\frac{\phi(c_m)}{c_m \Phi(-c_m)} \rightarrow 1$  as  $m \rightarrow \infty$  since  $c_m = -\Phi^{-1} \left( \frac{1}{m} \right) \rightarrow \infty$  as  $m \rightarrow \infty$ . This gives us the following approximation of the right-hand side of Eq. 75

366

367

368

$$\begin{aligned} \log \Phi(-c_m - b_m x) &\approx \log \left( \frac{1}{m} \right) - x + \mathcal{O}(b_m^2 x^2) \\ \Rightarrow \Phi(-c_m - b_m x) &\approx \frac{1}{m} e^{-x + \mathcal{O}(b_m^2 x^2)} \\ \Rightarrow \Phi(c_m + b_m x) &\approx 1 - \frac{1}{m} e^{-x + \mathcal{O}(b_m^2 x^2)}. \end{aligned} \quad (77)$$

Using the result given by Eq. 77, we have the following

369

$$\begin{aligned}
P\left(\frac{X_a^{\max} - c_m}{b_m} \leq x\right) &= P(X_a^{\max} \leq c_m + b_m x) \\
&= \Phi^m(c_m + b_m x) \\
&\approx \left(1 - \frac{1}{m} e^{-x + \mathcal{O}(b_m^2 x^2)}\right)^m \\
&= \left(1 - \frac{1}{m} e^{-x + \mathcal{O}\left(\frac{1}{c_m^2} x^2\right)}\right)^m \\
&\approx \left(1 - \frac{1}{m} e^{-x}\right)^m \\
\Rightarrow \lim_{m \rightarrow \infty} P\left(\frac{X_a^{\max} - c_m}{b_m} \leq x\right) &= \lim_{m \rightarrow \infty} \left(1 - \frac{1}{m} e^{-x}\right)^m \\
&= e^{-e^{-x}}.
\end{aligned} \tag{78}$$

The right-hand side of Eq. 78 is the cumulative distribution function of the standard Gumbel distribution. The mean of the asymptotic distribution is given by the following

370

371

$$E(X_a^{\max}) = \mu_{\max}^{(1)} = -\Phi^{-1}\left(\frac{1}{m}\right) - \frac{\gamma}{\Phi^{-1}\left(\frac{1}{m}\right)}. \tag{79}$$

where  $\gamma$  is the Euler-Mascheroni constant. The median of this distribution is given by the following

372

373

$$\tilde{\mu}_{\max} = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right). \tag{80}$$

Finally, the variance of the asymptotic distribution of the sample maximum is given by the following

374

375

$$\text{Var}(X_a^{\max}) = \frac{\pi^2}{6} \left( \frac{1}{-\Phi^{-1}\left(\frac{1}{m}\right)} \right)^2. \tag{81}$$

For typical sample sizes  $m$  in high-dimensional spaces, the variance estimate given by Eq. 81 exceeds the variance of the sample maximum significantly. Using the fact that  $-\Phi^{-1}\left(\frac{1}{m}\right) \sim \sqrt{2\log(m)}$  [18] and  $\frac{1}{2\log(m)} \leq \left(\frac{1}{-\Phi^{-1}\left(\frac{1}{m}\right)}\right)^2$  for  $m \geq 2$ , we can get a more accurate approximation of the variance with the following

376

377

378

379

$$\begin{aligned}
\sigma_{\max}^2(m) = \text{Var}(X_a^{\max}) &\approx \frac{\pi^2}{6} \left( \frac{1}{\sqrt{2\log(m)}} \right)^2 \\
&= \frac{\pi^2}{12\log(m)}.
\end{aligned} \tag{82}$$

Then the mean of the range of  $m$  iid standard normal random variables are given by the following

380

381

$$E(X_a^{\max} - X_a^{\min}) = 2\mu_{\max}^{(1)}(m) = 2 \left[ -\Phi^{-1}\left(\frac{1}{m}\right) - \frac{\gamma}{\Phi^{-1}\left(\frac{1}{m}\right)} \right]. \tag{83}$$

It is well known that the sample extremes from the standard normal distribution are approximately uncorrelated for large sample size  $m$  [16]. This implies that we can

382

383

approximate the variance of the range of  $m$  iid standard normal random variables with  
the following result

$$\begin{aligned}
\text{Var}(X_a^{\max} - X_a^{\min}) &\approx \text{Var}(X_a^{\max}) + \text{Var}(X_a^{\min}) \\
&= \sigma_{\max}^2(m) + \sigma_{\min}^2(m) \\
&= 2\sigma_{\max}^2(m) \\
&\approx 2 \left( \frac{\pi^2}{2\log(m)} \right) \\
&= \frac{\pi^2}{6\log(m)}.
\end{aligned} \tag{84}$$

For the purpose of approximating the mean and variance of the max-min normalized distance distribution, the formula for the median of the distribution of the attribute maximum yields more accurate results. That is, the approximation of the expected maximum given by Eq. 79 overestimates the sample maximum. The formula for the median of the sample maximum, given by Eq. 80, provides a more accurate estimate of this sample extreme. Therefore, the following estimate for the mean of the attribute range will be used instead

$$\text{E}(X_a^{\max} - X_a^{\min}) = 2\mu_{\max}^{(1)}(m) \approx 2 \left[ \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right) \right]. \tag{85}$$

We have already determined that  $\mu_{D_{ij}^{(q)}}$  and  $\sigma_{D_{ij}^{(q)}}^2$  are given by Eq. 31. Using the results given by Eqs. 85 and 84 and the general formulas for the mean and variance of the max-min normalized distance distribution given in Eq. 74, this leads us to the following asymptotic estimate for the distribution of the max-min normalized distances for standard normal data

$$D_{ij}^{(q*)} \sim \mathcal{N} \left( \frac{\mu_{D_{ij}^{(q)}}}{2\mu_{\max}^{(1)}(m)}, \frac{6\log(m)\sigma_{D_{ij}^{(q)}}^2}{\pi^2 + 24 \left[ \mu_{\max}^{(1)}(m) \right]^2 \log(m)} \right). \tag{86}$$

#### 2.4.2 Standard uniform data

Standard uniform data does not have support that is symmetric about 0. Due to the simplicity of the density function, however, we can derive the distribution of the maximum and minimum of a sample of size  $m$  explicitly. Using the general forms of the distribution functions of the maximum and minimum given by Eqs. 54 and 57, we have the following distribution functions for standard uniform data

$$F_{\max}(x) = x^m \tag{87}$$

and

$$F_{\min}(x) = 1 - (1 - x)^m. \tag{88}$$

Using the general forms of the density functions of the maximum and minimum given by Eqs. 55 and 58, we have the following density functions for standard uniform data

$$f_{\max}(x) = mx^{m-1} \tag{89}$$

and

$$f_{\min}(x) = m(1 - x)^{m-1} \tag{90}$$

Then the expected maximum and minimum are computed through straightforward integration as follows 408  
409

$$\begin{aligned} E(X_a^{\max}) &= \mu_{\max}^{(1)}(m) = \int_0^1 x f_{\max}(x) dx \\ &= \int_0^1 x [mx^{m-1}] dx \\ &= \frac{m}{m+1} \end{aligned} \quad (91)$$

and 410

$$\begin{aligned} E(X_a^{\min}) &= \mu_{\min}^{(1)}(m) = \int_0^1 x f_{\min}(x) dx \\ &= \int_0^1 x [m(1-x)^{m-1}] dx \\ &= \frac{1}{m+1}. \end{aligned} \quad (92)$$

We can compute the second moment about the origin of the sample range as follows 411

$$\begin{aligned} E[(X_a^{\max} - X_a^{\min})^2] &= E[(X_a^{\max})^2 - 2X_a^{\max}X_a^{\min} + (X_a^{\min})^2] \\ &= E[(X_a^{\max})^2] - 2E(X_a^{\max})E(X_a^{\min}) + E[(X_a^{\min})^2] \\ &= \mu_{\max}^{(2)}(m) - 2\mu_{\max}^{(1)}(m)\mu_{\min}^{(1)}(m) + \mu_{\min}^{(2)}(m) \\ &= \int_0^1 x^2 [mx^{m-1}] dx - 2 \left( \frac{m}{m+1} \right) \left( \frac{1}{m+1} \right) \\ &\quad + \int_0^1 x^2 [m(1-x)^{m-1}] dx \\ &= \frac{m}{m+2} - \frac{2m}{(m+1)^2} + \frac{2}{(m+1)(m+2)} \\ &= \frac{m^3 - m + 2}{(m+2)(m+1)^2}. \end{aligned} \quad (93)$$

Using the general formulas given in Eq. 71 and the mean ( $\mu_{D_{ij}^{(q)}}$ ) and variance ( $\sigma_{D_{ij}^{(q)}}^2$ ) 412  
given by Eq. 41, we have the following asymptotic estimate for the max-min normalized 413  
distance distribution for standard uniform data 414

$$D_{ij}^{(q*)} \sim \mathcal{N} \left( \frac{(m+1)\mu_{D_{ij}^{(q)}}}{m-1}, \frac{(m+2)(m+1)^2\sigma_{D_{ij}^{(q)}}^2}{m^3 - m + 2} \right). \quad (94)$$

## 2.5 Normalized Manhattan ( $q = 1$ ) 415

Using the general asymptotic results for mean and variance given by Eqs. 86 and 94 416  
for any value of  $q \in \mathbb{N}$ , we can substitute a particular value of  $q$  in order to determine 417  
a more specified asymptotic distance distribution for  $D_{ij}^{(q*)}$ . The following results are 418  
for the max-min normalized Manhattan ( $q = 1$ ) metric for both standard normal and 419  
standard uniform data. 420

### 2.5.1 Standard normal data

Substituting  $q = 1$  into Eq. 86, we have the following for standard normal data

$$\begin{aligned} \mathbb{E} \left( D_{ij}^{(1*)} \right) &= \frac{\mu_{D_{ij}^{(1)}}}{2\mu_{\max}^{(1)}(m)} \\ &= \frac{p}{\sqrt{\pi}\mu_{\max}^{(1)}(m)}, \end{aligned} \quad (95)$$

where  $\mu_{\max}^{(1)}(m)$  is given by Eq. 80.

Similarly, the variance of  $D_{ij}^{(1*)}$  is given by

$$\begin{aligned} \text{Var} \left( D_{ij}^{(1*)} \right) &= \frac{6\log(m)\sigma_{D_{ij}^{(1)}}^2}{\pi^2 + 24 \left[ \mu_{\max}^{(1)} \right]^2 \log(m)} \\ &= \frac{12p(\pi - 2)\log(m)}{\pi \left( \pi^2 + 24 \left[ \mu_{\max}^{(1)} \right]^2 \log(m) \right)}, \end{aligned} \quad (96)$$

where  $\mu_{\max}^{(1)}(m)$  is given by Eq. 80.

### 2.5.2 Standard uniform data

Substituting  $q = 1$  into Eq. 94, we have the following for standard uniform data

$$\begin{aligned} \mathbb{E} \left( D_{ij}^{(1*)} \right) &= \frac{(m+1)\mu_{D_{ij}^{(1)}}}{m-1} \\ &= \frac{(m+1)p}{3(m-1)}. \end{aligned} \quad (97)$$

Similarly, the variance of  $D_{ij}^{(1*)}$  is given by

$$\begin{aligned} \text{Var} \left( D_{ij}^{(1*)} \right) &= \frac{(m+2)(m+1)^2\sigma_{D_{ij}^{(1)}}^2}{m^3 - m + 2} \\ &= \frac{(m+2)(m+1)^2p}{18(m^3 - m + 2)}. \end{aligned} \quad (98)$$

## 2.6 Normalized Euclidean ( $q = 2$ )

Analogous to the previous section, we demonstrate the usage of Eqs. 86 and 94 for the max-min normalized Euclidean ( $q = 2$ ) metric for both standard normal and standard uniform data.

### 2.6.1 Standard normal data

Substituting  $q = 2$  into Eq. 86, we have the following for standard normal data

$$\begin{aligned} \mathbb{E} \left( D_{ij}^{(2*)} \right) &= \frac{\mu_{D_{ij}^{(2)}}}{2\mu_{\max}^{(1)}(m)} \\ &= \frac{\sqrt{2p-1}}{2\mu_{\max}^{(1)}(m)}, \end{aligned} \quad (99)$$

where  $\mu_{\max}^{(1)}(m)$  is given by Eq. 80. 435

Similarly, the variance of  $D_{ij}^{(2*)}$  is given by 436

$$\begin{aligned} \text{Var} \left( D_{ij}^{(2*)} \right) &= \frac{6\log(m)\sigma_{D_{ij}^{(2)}}^2}{\pi^2 + 24 \left[ \mu_{\max}^{(1)}(m) \right]^2 \log(m)} \\ &= \frac{6\log(m)}{\pi^2 + 24 \left[ \mu_{\max}^{(1)}(m) \right]^2 \log(m)}, \end{aligned} \quad (100)$$

where  $\mu_{\max}^{(1)}(m)$  is given by Eq. 80. 437

### 2.6.2 Standard uniform data 438

Substituting  $q = 2$  into Eq. 94, we have the following for standard uniform data 439

$$\begin{aligned} \text{E} \left( D_{ij}^{(2*)} \right) &= \frac{(m+1)\mu_{D_{ij}^{(2)}}}{m-1} \\ &= \sqrt{\frac{p}{6} - \frac{7}{120}} \left( \frac{m+1}{m-1} \right). \end{aligned} \quad (101)$$

Similarly, the variance of  $D_{ij}^{(2*)}$  is given by 440

$$\begin{aligned} \text{Var} \left( D_{ij}^{(2*)} \right) &= \frac{(m+2)(m+1)^2\sigma_{D_{ij}^{(2)}}^2}{m^3 - m + 2} \\ &= \frac{7(m+2)(m+1)^2}{120(m^3 - m + 2)}. \end{aligned} \quad (102)$$



**Table 1.** Summary of distance distribution derivations for standard normal and standard uniform data. Asymptotic estimates are given for both standard and max-min normalized q-metrics. These estimates are relevant for all  $q \in \mathbb{N}$  and  $p \geq 100$ .

q-Metric	Data	Stat	Formula (Eq. #)
standard (Eq. 2)	$\mathcal{N}(0, 1)$	mean	$\left( \frac{2^q \Gamma(\frac{q+1}{2}) p}{\sqrt{\pi}} \right)^{1/q} \quad (38)$
	$\mathcal{N}(0, 1)$	variance	$\frac{4^q p}{q^2 \left( \frac{2^q \Gamma(\frac{1}{2} q + \frac{1}{2})}{\sqrt{\pi}} p \right)^{2(1-\frac{1}{q})}} \left[ \frac{\Gamma(q+\frac{1}{2})}{\sqrt{\pi}} - \frac{\Gamma^2(\frac{1}{2} q + \frac{1}{2})}{\pi} \right] \quad (38)$
	$\mathcal{U}(0, 1)$	mean	$\left( \frac{2p}{(q+2)(q+1)} \right)^{1/q} \quad (48)$
	$\mathcal{U}(0, 1)$	variance	$\frac{p}{q^2 \left( \frac{2p}{(q+2)(q+1)} \right)^{2(1-\frac{1}{q})}} \left[ \frac{1}{(q+1)(2q+1)} - \left( \frac{2}{(q+2)(q+1)} \right)^2 \right] \quad (48)$
max-min normalized (Eq. 59)	$\mathcal{N}(0, 1)$	mean	$\frac{\mu_{D_{ij}}^{(q)}}{2\mu_{\max}^{(1)}(m)} \quad (93)$ where $\mu_{D_{ij}}^{(q)}$ and $\mu_{\max}^{(1)}(m)$ are given by Eqs. 38 and 87, respectively.
	$\mathcal{N}(0, 1)$	variance	$\frac{6\log(m)\sigma_{D_{ij}}^2}{\pi^2 + 24[\mu_{\max}^{(1)}(m)]^2 \log(m)} \quad (93)$ where $\sigma_{D_{ij}}^2$ and $\mu_{\max}^{(1)}(m)$ are given by Eqs. 38 and 87, respectively.
	$\mathcal{U}(0, 1)$	mean	$\frac{(m+1)\mu_{D_{ij}}^{(q)}}{m-1} \quad (101)$ where $\mu_{D_{ij}}^{(q)}$ is given by Eq. 48
	$\mathcal{U}(0, 1)$	variance	$\frac{(m+2)(m+1)^2 \sigma_{D_{ij}}^2}{m^3 - m + 2} \quad (101)$ where $\sigma_{D_{ij}}^2$ is given by Eq. 48

**Table 2.** Asymptotic estimates for means and variances for the standard  $L_1$  and  $L_2$  distance distributions. Estimates for both standard normal and standard uniform data are given.

$q$ -Metric	Data	Stat	Formula (Eq. #)
standard ( $L_1$ )	$\mathcal{N}(0, 1)$	mean	$\frac{2p}{\sqrt{\pi}} \quad (38)$
		variance	$\frac{2(\pi-2)p}{\pi} \quad (38)$
	$\mathcal{U}(0, 1)$	mean	$\frac{p}{3} \quad (48)$
		variance	$\frac{p}{18} \quad (48)$
standard ( $L_2$ )	$\mathcal{N}(0, 1)$	mean	$\sqrt{2p-1} \quad (38)$
		variance	$1 \quad (38)$
	$\mathcal{U}(0, 1)$	mean	$\sqrt{\frac{p}{6} - \frac{7}{120}} \quad (48)$
		variance	$\frac{7}{120} \quad (48)$

**Table 3.** Asymptotic estimates for means and variances for the max-min normalized  $L_1$  and  $L_2$  distance distributions. Estimates for both standard normal and standard uniform data are given.

$q$ -Metric	Data	Stat	Formula (Eq. #)
max-min normalized ( $L_1$ )	$\mathcal{N}(0, 1)$	mean	$\frac{p}{\sqrt{\pi}\mu_{\max}^{(1)}(m)} \quad (93)$ where $\mu_{\max}^{(1)}(m) = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right)$
		variance	$\frac{12p(\pi-2)\log(m)}{\pi\left(\pi^2+24\left[\mu_{\max}^{(1)}(m)\right]^2\log(m)\right)} \quad (93)$ where $\mu_{\max}^{(1)}(m) = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right)$
	$\mathcal{U}(0, 1)$	mean	$\frac{(m+1)p}{3(m-1)} \quad (101)$
		variance	$\frac{(m+2)(m+1)^2p}{18(m^3-m+2)} \quad (48)$
max-min normalized ( $L_2$ )	$\mathcal{N}(0, 1)$	mean	$\frac{\sqrt{2p-1}}{2\mu_{\max}^{(1)}(m)} \quad (93)$ where $\mu_{\max}^{(1)}(m) = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right)$
		variance	$\frac{6\log(m)}{\pi^2+24\left[\mu_{\max}^{(1)}(m)\right]^2\log(m)} \quad (93)$ where $\mu_{\max}^{(1)}(m) = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right)$
	$\mathcal{U}(0, 1)$	mean	$\sqrt{\frac{p}{6} - \frac{7}{120} \left(\frac{m+1}{m-1}\right)} \quad (101)$
		variance	$\frac{7(m+2)(m+1)^2}{120(m^3-m+2)} \quad (101)$

### 3 GWAS distance distributions

Consider a GWAS data set, which has the following encoding based on minor allele frequency

$$X_{ia} = \begin{cases} 0 & \text{if there are no minor alleles at locus } a, \\ 1 & \text{if there is 1 minor allele at locus } a, \\ 2 & \text{if there are 2 minor alleles at locus } a. \end{cases} \quad (103)$$

A minor allele at a particular locus  $a$  is the least frequent of the two alleles at that particular locus  $a$ . For random GWAS data sets, we can think  $X_{ia}$  as the number of successes in two Bernoulli trials. That is,  $X_{ia} \sim \mathcal{B}(2, f_a)$  where  $f_a$  is the probability of success. The success probability  $f_a$  is the probability of a minor allele occurring at  $a$ . Furthermore, the minor allele probabilities are assumed to be independent and identically distributed according to  $\mathcal{U}(l, u)$ , where  $l$  and  $u$  are the lower and upper bounds, respectively, of the sampling distribution's support. Two commonly known types of distance metrics for GWAS data are the Genotype Mismatch (GM) and Allele Mismatch (AM) metrics. The GM and AM metrics are defined by

$$d_{ij}^{\text{GM}}(a) = \begin{cases} 0 & \text{if } X_{ia} \neq X_{ja}, \\ 1 & \text{otherwise} \end{cases} \quad (104)$$

and

$$d_{ij}^{\text{AM}}(a) = \frac{1}{2} |X_{ia} - X_{ja}|. \quad (105)$$

A more informative metric must take into account whether differences in allele frequency at a particular locus  $a$  result in transitions or transversions. A metric that accounts for transitions (Ti) and transversions (Tv) was introduced in [4]. This metric is given by the following

$$d_{ij}^{\text{TiTv}}(a) = \begin{cases} 0 & \text{if } X_{ia} = X_{ja} \text{ and Ti/Tv,} \\ 1/4 & \text{if } |X_{ia} - X_{ja}| = 1 \text{ and Ti,} \\ 1/2 & \text{if } |X_{ia} - X_{ja}| = 1 \text{ and Tv,} \\ 3/4 & \text{if } |X_{ia} - X_{ja}| = 2 \text{ and Ti,} \\ 1 & \text{if } |X_{ia} - X_{ja}| = 2 \text{ and Tv.} \end{cases} \quad (106)$$

With any of the three metrics given by Eqs. 104 - 106, we compute the pairwise distance between two instances  $i$  and  $j$  using Eq. 1 with  $q = 1$ . Assuming that all data entries  $X_{ia}$  are independent and identically distributed, we have already shown that the distribution of pairwise distances is asymptotically normal regardless of data distribution and value of  $q$ . Therefore, the distance distributions induced by each of the GWAS metrics given by Eqs. 104 - 106 are asymptotically normal. Thus, we will proceed by deriving the mean and variance for each distance distribution induced by these three GWAS metrics.

### 3.1 GM distance distribution

466

The expected value of the GM metric is given by the following

467

$$\begin{aligned}
E \left[ d_{ij}^{\text{GM}}(a) \right] &= \sum_{k=0}^1 k \cdot P \left[ d_{ij}^{\text{GM}}(a) = k \right] \\
&= 0 \cdot P \left[ d_{ij}^{\text{GM}}(a) = 0 \right] + 1 \cdot P \left[ d_{ij}^{\text{GM}}(a) = 1 \right] \\
&= P \left[ d_{ij}^{\text{GM}}(a) = 1 \right] \\
&= 2P[X_{ia} = 0, X_{ja} = 1] + 2P[X_{ia} = 1, X_{ja} = 2] + 2P[X_{ia} = 0, X_{ja} = 2] \\
&= 4(1 - f_a)^3 f_a + 4(1 - f_a) f_a^3 + 2(1 - f_a)^2 f_a^2 \\
&= 2 \left[ 2(1 - f_a)^3 f_a + 2(1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2 \right] \\
&= 2F(a),
\end{aligned} \tag{107}$$

where  $F(a) = 2(1 - f_a)^3 f_a + 2(1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2$ .

468

Then the expected pairwise GM distance between instances  $i$  and  $j$  is computed as follows

469

470

$$\begin{aligned}
E(D_{ij}^{\text{GM}}) &= E \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{GM}}(a) \right) \\
&= \sum_{a \in \mathcal{A}} E \left[ d_{ij}^{\text{GM}}(a) \right] \\
&= 2 \sum_{a \in \mathcal{A}} F(a).
\end{aligned} \tag{108}$$

The second moment about the origin for the GM distance is computed as follows

471

$$\begin{aligned}
E \left[ (D_{ij}^{\text{GM}})^2 \right] &= E \left[ \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{GM}}(a) \right)^2 \right] \\
&= E \left[ \sum_{a \in \mathcal{A}} \left( d_{ij}^{\text{GM}}(a) \right)^2 \right] + 2E \left[ \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} d_{ij}^{\text{GM}}(r) \cdot d_{ij}^{\text{GM}}(s) \right] \\
&= \sum_{a \in \mathcal{A}} \left( \sum_{k=0}^1 k^2 \cdot P \left[ d_{ij}^{\text{GM}}(a) = k \right] \right) \\
&\quad + 2 \sum_{a \in \mathcal{A}} \sum_{s \leq r-1} \left( \sum_{k=0}^1 k \cdot P \left[ d_{ij}^{\text{GM}}(r) = k \right] \right) \cdot \left( \sum_{k=0}^1 k \cdot P \left[ d_{ij}^{\text{GM}}(s) = k \right] \right) \\
&= 2 \sum_{a \in \mathcal{A}} F(a) + 8 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r, s\}} F(\lambda),
\end{aligned} \tag{109}$$

where  $F(a) = 2(1 - f_a)^3 f_a + 2(1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2$ .

472

Using the moments given by Eqs. 108 and 109, the variance is computed as follows 473

$$\begin{aligned}
\text{Var}(D_{ij}^{\text{GM}}) &= \mathbb{E} \left[ (D_{ij}^{\text{GM}})^2 \right] - [\mathbb{E}(D_{ij}^{\text{GM}})]^2 \\
&= 2 \sum_{a \in \mathcal{A}} F(a) + 8 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r,s\}} F(\lambda) - 4 \left( \sum_{a \in \mathcal{A}} F(a) \right)^2 \\
&= 2 \sum_{a \in \mathcal{A}} F(a) - 4 \sum_{a \in \mathcal{A}} F^2(a) \\
&= 2 \sum_{a \in \mathcal{A}} F(a)[1 - 2F(a)],
\end{aligned} \tag{110}$$

where  $F(a) = 2(1 - f_a)^3 f_a + 2(1 - f_a)f_a^3 + (1 - f_a)^2 f_a^2$ . 474

With the mean and variance estimates given by Eqs. 108 and 110, the asymptotic GM distance distribution is given by the following 475  
476

$$D_{ij}^{\text{GM}} \sim \mathcal{N} \left( 2 \sum_{a \in \mathcal{A}} F(a), 2 \sum_{a \in \mathcal{A}} F(a)[1 - 2F(a)] \right), \tag{111}$$

where  $F(a) = 2(1 - f_a)^3 f_a + 2(1 - f_a)f_a^3 + (1 - f_a)^2 f_a^2$ . 477

### 3.2 AM distance distribution 478

The expected value of the AM metric is given by the following 479

$$\begin{aligned}
\mathbb{E} [d_{ij}^{\text{AM}}(a)] &= \sum_{k \in \mathcal{D}} k \cdot \mathbb{P} [d_{ij}^{\text{AM}}(a) = k] \\
&= 0 \cdot \mathbb{P} [d_{ij}^{\text{AM}}(a) = 0] + \frac{1}{2} \cdot \mathbb{P} [d_{ij}^{\text{AM}}(a) = \frac{1}{2}] + 1 \cdot \mathbb{P} [d_{ij}^{\text{AM}}(a) = 1] \\
&= \frac{1}{2} (2\mathbb{P} [X_{ia} = 0, X_{ja} = 1] + 2\mathbb{P} [X_{ia} = 1, X_{ja} = 2]) \\
&\quad + 2\mathbb{P} [X_{ia} = 0, X_{ja} = 2] \\
&= \mathbb{P} [X_{ia} = 0, X_{ja} = 1] + \mathbb{P} [X_{ia} = 1, X_{ja} = 2] + 2\mathbb{P} [X_{ia} = 0, X_{ja} = 2] \\
&= 2(1 - f_a)^3 f_a + 2(1 - f_a)f_a^3 + 2(1 - f_a)^2 f_a^2 \\
&= 2 [(1 - f_a)^3 f_a + (1 - f_a)f_a^3 + (1 - f_a)^2 f_a^2] \\
&= 2F(a),
\end{aligned} \tag{112}$$

where  $F(a) = (1 - f_a)^3 f_a + (1 - f_a)f_a^3 + (1 - f_a)^2 f_a^2$  and  $\mathcal{D} = \{0, \frac{1}{2}, 1\}$ . 480

Then the expected pairwise AM distance between instances  $i$  and  $j$  is computed as follows 481  
482

$$\begin{aligned}
\mathbb{E}(D_{ij}^{\text{AM}}) &= \mathbb{E} \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{AM}}(a) \right) \\
&= \sum_{a \in \mathcal{A}} \mathbb{E} [d_{ij}^{\text{AM}}(a)] \\
&= 2 \sum_{a \in \mathcal{A}} F(a).
\end{aligned} \tag{113}$$

The second moment about the origin for the AM distance is computed as follows

483

$$\begin{aligned}
\mathbb{E} \left[ (D_{ij}^{\text{AM}})^2 \right] &= \mathbb{E} \left[ \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{AM}}(a) \right)^2 \right] \\
&= \mathbb{E} \left[ \sum_{a \in \mathcal{A}} \left( d_{ij}^{\text{AM}}(a) \right)^2 \right] + 2\mathbb{E} \left[ \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} d_{ij}^{\text{AM}}(r) \cdot d_{ij}^{\text{AM}}(s) \right] \\
&= \sum_{a \in \mathcal{A}} \left( \sum_{k \in \mathcal{D}} k^2 \cdot \mathbb{P} \left[ d_{ij}^{\text{AM}}(a) = k \right] \right) \\
&\quad + 2 \sum_{a \in \mathcal{A}} \sum_{s \leq r-1} \left( \sum_{k \in \mathcal{D}} k \cdot \mathbb{P} \left[ d_{ij}^{\text{AM}}(r) = k \right] \right) \cdot \left( \sum_{k \in \mathcal{D}} k \cdot \mathbb{P} \left[ d_{ij}^{\text{AM}}(s) = k \right] \right) \\
&= \sum_{a \in \mathcal{A}} G(a) + 8 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r, s\}} F(\lambda),
\end{aligned} \tag{114}$$

where  $G(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a) + 2(1 - f_a)^2 f_a^2$  and  $F(\lambda) = (1 - f_\lambda)^3 f_\lambda + f_\lambda^3 (1 - f_\lambda) + (1 - f_\lambda)^2 f_\lambda^2$ .

484

485

Using the moments given by Eqs. 113 and 114, the variance is computed as follows

486

$$\begin{aligned}
\text{Var} (D_{ij}^{\text{AM}}) &= \mathbb{E} \left[ (D_{ij}^{\text{AM}})^2 \right] - [\mathbb{E} (D_{ij}^{\text{AM}})]^2 \\
&= \sum_{a \in \mathcal{A}} G(a) + 8 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r, s\}} F(\lambda) - 4 \left( \sum_{a \in \mathcal{A}} F(a) \right)^2 \\
&= \sum_{a \in \mathcal{A}} G(a) - 4 \sum_{a \in \mathcal{A}} F^2(a) \\
&= \sum_{a \in \mathcal{A}} [G(a) - 4F^2(a)],
\end{aligned} \tag{115}$$

where  $G(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a) + 2(1 - f_a)^2 f_a^2$  and  $F(\lambda) = (1 - f_\lambda)^3 f_\lambda + f_\lambda^3 (1 - f_\lambda) + (1 - f_\lambda)^2 f_\lambda^2$ .

487

488

With the mean and variance estimates given by Eqs. 113 and 115, the asymptotic AM distance distribution is given by the following

489

490

$$D_{ij}^{\text{AM}} \sim \mathcal{N} \left( 2 \sum_{a \in \mathcal{A}} F(a), \sum_{a \in \mathcal{A}} [G(a) - 4F^2(a)] \right), \tag{116}$$

where  $G(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a) + 2(1 - f_a)^2 f_a^2$  and  $F(\lambda) = (1 - f_\lambda)^3 f_\lambda + f_\lambda^3 (1 - f_\lambda) + (1 - f_\lambda)^2 f_\lambda^2$ .

491

492

### 3.3 TiTv distance distribution

493

The TiTv metric allows for one to account for both genotype mismatch, allele mismatch, transition, and transversion. However, this added dimension of information requires knowledge of the nucleotide makeup at a particular locus. A sufficient condition to compute the TiTv metric between instances  $i$  and  $j$  is that we know whether the nucleotides associated with a particular locus  $a$  are both purines (PuPu), purine and pyrimidine (PuPy), or both pyrimidines (PyPy). A diagram showing possible transitions and transversions that may occur is given by Fig. 2. Purines (A and G) and pyrimidines (C and T) are shown at the top and bottom, respectively. Transitions occur in the cases of PuPu and PyPy, while transversion occur only with PuPy encoding.

494

495

496

497


498

499

500

501

502



**Fig 2.** Purines (A and G) and pyrimidines (C and T) are shown. Transitions occur when a mutation involves purine-to-purine or pyrimidine-to-pyrimidine insertion. Transversions occur when a purine-to-pyrimidine or pyrimidine-to-purine insertion happens, which is a more extreme case. There are visibly more possibilities for transversions to occur than there are transitions, but there are about twice as many transitions in real data.

503

This information is always given in a particular data set. Let  $\gamma_0$ ,  $\gamma_1$ , and  $\gamma_2$  denote the probabilities of PuPu, PuPy, and PyPy, respectively, for the  $p$  loci of data matrix  $X$ . In real data, there are approximately twice as many transitions as there are transversions. That is, the probability of a transition  $P(\text{Ti})$  is approximately twice the probability of transversion  $P(\text{Tv})$ . It is likely that any particular data set will not satisfy this criterion exactly. In this general case, we have  $P(\text{Ti})$  being equal to some multiple  $\eta$  times  $P(\text{Tv})$ . In order to enforce this general constraint in simulated data, we define the following set of equalities

$$\gamma_0 + \gamma_1 + \gamma_2 = 1, \quad (117)$$

$$P(\text{Ti}) - \eta P(\text{Tv}) = 0. \quad (118)$$

Using this PuPu, PuPy, and PyPy encoding, the probability of a transversion occurring at any fixed locus  $a$  is given by the following

504  
505

$$P(\text{Tv}) = \gamma_1. \quad (119)$$

Using the constraints given by Eqs. 117 and 118, the probability of a transition occurring at locus  $a$  is computed as follows

506  
507

$$P(\text{Ti}) = \gamma_0 + \gamma_2. \quad (120)$$

Also based on the constraints given by Eqs. 117 and 118, it is clear that we have  $P(\text{Tv}) = \frac{1}{\eta+1}$  and  $P(\text{Ti}) = \frac{\eta}{\eta+1}$ . Without loss of generality, we then sample

508  
509

$$\gamma_0 \sim \mathcal{U}\left(\varepsilon, \frac{\eta}{\eta+1} - \varepsilon\right), \quad (121)$$

where  $\varepsilon$  is some small positive real number.

510

Then it immediately follows that we have

511

$$\gamma_2 = \frac{\eta}{\eta+1} - \gamma_0. \quad (122)$$

However, we can derive the mean and variance of the distance distribution induced by the TiTv metric without specifying any relationship between  $\gamma_0$ ,  $\gamma_1$ , and  $\gamma_2$ . We

512  
513



proceed by computing  $P \left[ d_{ij}^{\text{TiTv}}(a) = k \right]$  for each  $k \in \mathcal{D} = \{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}$ . Let  $y$  represent a random sample of size  $p$  from  $\{0, 1, 2\}$ , where

$$y_a = \begin{cases} 0 & \text{if locus } a \text{ is PuPu,} \\ 1 & \text{if locus } a \text{ is PuPy,} \\ 2 & \text{if locus } a \text{ is PyPy.} \end{cases} \quad (123)$$

We derive  $P \left[ d_{ij}^{\text{TiTv}}(a) = 0 \right]$  as follows

$$\begin{aligned} P \left[ d_{ij}^{\text{TiTv}}(a) = 0 \right] &= P[y_a = 0, X_{ia} = X_{ja}] \\ &\quad + P[y_a = 1, X_{ia} = X_{ja}] \\ &\quad + P[y_a = 2, X_{ia} = X_{ja}] \\ &= \gamma_0 [(1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2] \\ &\quad + \gamma_1 [(1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2] \\ &\quad + \gamma_2 [(1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2] \\ &= (\gamma_0 + \gamma_1 + \gamma_2) [(1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2] \\ &= (1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2. \end{aligned} \quad (124)$$

We derive  $P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{1}{4} \right]$  as follows

$$\begin{aligned} P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{1}{4} \right] &= 2P[y_a = 0, X_{ia} = 0, X_{ja} = 1] \\ &\quad + 2P[y_a = 0, X_{ia} = 1, X_{ja} = 2] \\ &\quad + 2P[y_a = 2, X_{ia} = 0, X_{ja} = 1] \\ &\quad + 2P[y_a = 2, X_{ia} = 1, X_{ja} = 2] \\ &= 4\gamma_0(1 - f_a)^3 f_a + 4\gamma_0 f_a^3(1 - f_a) + 4\gamma_2(1 - f_a)^3 f_a \\ &\quad + 4\gamma_2 f_a^3(1 - f_a) \\ &= 4\gamma_0 [(1 - f_a)^3 f_a + f_a^3(1 - f_a)] \\ &\quad + 4\gamma_2 [(1 - f_a)^3 f_a + f_a^3(1 - f_a)] \\ &= 4(\gamma_0 + \gamma_2) [(1 - f_a)^3 f_a + f_a^3(1 - f_a)]. \end{aligned} \quad (125)$$

We derive  $P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{1}{2} \right]$  as follows

$$\begin{aligned} P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{1}{2} \right] &= 2P[y_a = 1, X_{ia} = 0, X_{ja} = 1] \\ &\quad + 2P[y_a = 1, X_{ia} = 1, X_{ja} = 2] \\ &= 4\gamma_1(1 - f_a)^3 f_a + 4\gamma_1 f_a^3(1 - f_a) \\ &= 4\gamma_1 [(1 - f_a)^3 f_a + f_a^3(1 - f_a)]. \end{aligned} \quad (126)$$

We derive  $P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{3}{4} \right]$  as follows

$$\begin{aligned} P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{3}{4} \right] &= 2P[y_a = 0, X_{ia} = 0, X_{ja} = 2] \\ &\quad + 2P[y_a = 2, X_{ia} = 0, X_{ja} = 2] \\ &= 2\gamma_0(1 - f_a)^2 f_a^2 + 2\gamma_2(1 - f_a)^2 f_a^2 \\ &= 2(\gamma_0 + \gamma_2)(1 - f_a)^2 f_a^2. \end{aligned} \quad (127)$$

We derive  $P \left[ d_{ij}^{\text{TiTv}}(a) = 1 \right]$  as follows

520

$$\begin{aligned} P \left[ d_{ij}^{\text{TiTv}}(a) = 1 \right] &= 2P \left[ y_a = 1, X_{ia} = 0, X_{ja} = 2 \right] \\ &= 2\gamma_1(1 - f_a)^2 f_a^2. \end{aligned} \quad (128)$$

Using Eqs. 124 - 128, we compute the expected TiTv distance between instances  $i$  and  $j$  as follows

521

522

$$\begin{aligned} E \left( D_{ij}^{\text{TiTv}} \right) &= \sum_{a \in \mathcal{A}} \left( \sum_{k \in \mathcal{D}} k \cdot P \left[ d_{ij}^{\text{TiTv}}(a) = k \right] \right) \\ &= (\gamma_0 + \gamma_2 + 2\gamma_1) \sum_{a \in \mathcal{A}} \left[ (1 - f_a)^3 f_a + f_a^3 (1 - f_a) \right] \\ &\quad + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} (1 - f_a)^2 f_a^2 \\ &= (\gamma_0 + \gamma_2 + 2\gamma_1) \sum_{a \in \mathcal{A}} F(a) + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G(a), \end{aligned} \quad (129)$$

where  $F(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a)$  and  $G(a) = (1 - f_a)^2 f_a^2$ .

523

The second moment about the origin for the TiTv distance is computed as follows

524

$$\begin{aligned} E \left[ \left( D_{ij}^{\text{TiTv}} \right)^2 \right] &= E \left[ \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{TiTv}}(a) \right)^2 \right] \\ &= E \left[ \sum_{a \in \mathcal{A}} \left( d_{ij}^{\text{TiTv}}(a) \right)^2 \right] + 2E \left[ \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} d_{ij}^{\text{TiTv}}(r) \cdot d_{ij}^{\text{TiTv}}(s) \right] \\ &= \sum_{a \in \mathcal{A}} \left( \sum_{k \in \mathcal{D}} k^2 \cdot P \left[ d_{ij}^{\text{TiTv}}(a) = k \right] \right) \\ &\quad + 2 \sum_{a \in \mathcal{A}} \sum_{s \leq r-1} \left( \sum_{k \in \mathcal{D}} k \cdot P \left[ d_{ij}^{\text{TiTv}}(r) = k \right] \right) \cdot \left( \sum_{k \in \mathcal{D}} k \cdot P \left[ d_{ij}^{\text{TiTv}}(s) = k \right] \right) \\ &= \left[ \frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1 \right] \sum_{a \in \mathcal{A}} F(a) + \left[ \frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G(a) \\ &\quad + 2 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r, s\}} \left( [\gamma_0 + \gamma_2 + 2\gamma_1] F(\lambda) + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] G(\lambda) \right), \end{aligned} \quad (130)$$

where  $F(\lambda) = (1 - f_\lambda)^3 f_\lambda + f_\lambda^3 (1 - f_\lambda)$  and  $G(\lambda) = (1 - f_\lambda)^2 f_\lambda^2$ .

525

Using the moments given by Eqs. 129 and 130, the variance is computed as follows 526

$$\begin{aligned}
\text{Var}(D_{ij}^{\text{TiTv}}) &= \mathbb{E} \left[ (D_{ij}^{\text{TiTv}})^2 \right] - [\mathbb{E}(D_{ij}^{\text{TiTv}})]^2 \\
&= \left[ \frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1 \right] \sum_{a \in \mathcal{A}} F(a) + \left[ \frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G(a) \\
&\quad + 2 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r,s\}} \left( [\gamma_0 + \gamma_2 + 2\gamma_1] F(\lambda) + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] G(\lambda) \right) \\
&\quad - \left( [\gamma_0 + \gamma_2 + 2\gamma_1] \sum_{a \in \mathcal{A}} F(a) + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G(a) \right)^2 \\
&= \left[ \frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1 \right] \sum_{a \in \mathcal{A}} F(a) + \left[ \frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G(a) \\
&\quad - \sum_{a \in \mathcal{A}} \left( [\gamma_0 + \gamma_2 + 2\gamma_1] F(a) + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] G(a) \right)^2,
\end{aligned} \tag{131}$$

where  $F(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a)$  and  $G(a) = (1 - f_a)^2 f_a^2$ . 527

With the mean and variance estimates given by Eqs. 129 and 131, the asymptotic 528  
TiTv distance distribution is given by the following 529

$$\begin{aligned}
D_{ij}^{\text{TiTv}} &\sim \mathcal{N} \left( (\gamma_0 + \gamma_2 + 2\gamma_1) \sum_{a \in \mathcal{A}} F(a) + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G(a), \right. \\
&\quad \left[ \frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1 \right] \sum_{a \in \mathcal{A}} F(a) + \left[ \frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G(a) \\
&\quad \left. - \sum_{a \in \mathcal{A}} \left( [\gamma_0 + \gamma_2 + 2\gamma_1] F(a) + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] G(a) \right)^2 \right),
\end{aligned} \tag{132}$$

where  $F(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a)$  and  $G(a) = (1 - f_a)^2 f_a^2$ . 530


Given upper and lower bounds  $l$  and  $u$ , respectively, of the success probability 531  
sampling interval, the average success probability (or average MAF) is computed as 532  
follows 533

$$\bar{f}_a = \frac{1}{2}(l + u). \tag{133}$$


The maximum TiTv distance occurs at  $\bar{f}_a = 0.5$  for any fixed Ti/Tv ratio  $\eta$  (Eq. 118), 534  
which is the inflection point about which the minor allele changes at locus  $a$  (Fig. 3). 535  
If few minor alleles are present ( $\bar{f}_a \rightarrow 0$ ), the predicted TiTv distance approaches 536  
0. The same is true after the minor allele switches ( $\bar{f}_a \rightarrow 1$ ). We fixed the Ti/Tv 537  
ratio  $\eta$  and generated simulated TiTv distances for  $\bar{f}_a = 0.055, 0.150, 0.250$ , and  $0.350$  538  
(Fig. 4A). For fixed  $\eta$ , TiTv distance increases significantly with increased  $\bar{f}_a$ . We 539  
similarly fixed the average minor allele frequency  $\bar{f}_a$  and generated simulated TiTv 540  
distances for  $\eta = \text{Ti/Tv} = 0.5, 1, 1.5$ , and  $2$  (Fig. 4C). The TiTv distance decreases 541  
slightly with increased  $\eta = \text{Ti/Tv}$ . As  $\eta \rightarrow 0^+$ , the data is approaching all Tv and no 542  
Ti, which means the TiTv distance is larger by definition. On the other hand, the TiTv 543  
distance decreases as  $\eta \rightarrow 2^-$  because the data is approaching approximately twice as 544  
many Ti as there are Tv, which is typical for GWAS data in humans. 545

We also compared theoretical and sample moments as a function of  $\eta = \text{Ti/Tv}$  and  $\bar{f}_a$  546  
for the TiTv distance metric (Fig. 4B and D). We fixed  $\bar{f}_a$  and computed the theoretical 547  
and simulated moments as a function of  $\eta$  (Fig. 4B). Theoretical average TiTv distance, 548  
given by Eq. 129, and simulated TiTv average distance are approximately equal as 549

$\eta$  increases. Theoretical standard deviation, given by Eq. 131, and simulated TiTv standard deviation differ slightly. We also fixed  $\eta$  and computed theoretical and sample moments as a function of  $\bar{f}_a$  (Fig. 4D). In this case, there is approximate agreement with simulated and theoretical moments as  $\bar{f}_a$  increases.



**Fig 3.** Predicted average TiTv distance as a function of average minor allele frequency  $\bar{f}_a$  (see Eq. 133). Success probabilities  $f_a$  were drawn from a sliding window interval from 0.01 to 0.9 in increments of about 0.009. With  $\eta = 0.1$ , where  $\eta$  is the Ti/Tv ratio given by Eq. 117, Tv is ten times more likely than Ti so the distance is large. Increasing to  $\eta = 1$ , Tv and Ti are equally likely so the distance is moderate. In line with real data for  $\eta = 2$ , Tv is half as likely as Ti so the distance is relatively small.



**Fig 4.** Density curves and moments of TiTv distance as a function of average MAF  $\bar{f}_a$ , given by Eq. 133, and Ti/Tv ratio  $\eta$ , given by Eq. 118. **(A)** For fixed  $\eta = 2$ , TiTv distance density is plotted as a function of increasing  $\bar{f}_a = 0.055, 0.150, 0.250$ , and  $0.350$ . TiTv distance increases as  $\bar{f}_a$  approaches a maximum of  $0.5$ , which means that there is about the same frequency of minor alleles as primary alleles at locus  $a$ . **(B)** Simulated and predicted mean  $\pm$  SD are shown as a function of increasing Ti/Tv ratio  $\eta$ . Distance decreases as Tv becomes more frequent than Ti. Theoretical standard deviation is slightly larger than simulated, but the means are approximately the same. **(C)** For fixed  $\bar{f}_a = 0.055$ , TiTv distance density is plotted as a function of increasing  $\eta = 0.5, 1, 1.5$ , and  $2$ . TiTv distance decreases as  $\eta$ , the Ti/Tv ratio, increases. For  $\eta = \text{Ti/Tv} = 0.5$ , there are twice as many transversions as there are transitions. On the other hand,  $\eta = \text{Ti/Tv} = 2$  indicates that there are half as many transversions as there are transitions. Since transversions encode a larger magnitude distance than transitions in Eq. 106, this behavior is expected. **(D)** Simulated and predicted mean  $\pm$  SD are shown as a function of increasing average MAF  $\bar{f}_a$ . Distance increases as the number of minor alleles increases at each locus  $a$ . Theoretical and simulated moments are approximately the same.

**Table 4.** Summary of distance distribution derivations for GWAS data.

GWAS-Metric	Stat	Formula (Eq. #)
GM (Eq. 103)	mean	$\boxed{2 \sum_{a \in \mathcal{A}} F(a)} \quad (110)$ where $F(a) = 2(1 - f_a)^3 f_a + 2f_a^3(1 - f_a) + (1 - f_a)^2 f_a^2$
	variance	$\boxed{2 \sum_{a \in \mathcal{A}} F(a)[1 - 2F(a)]} \quad (110)$ where $F(a) = 2(1 - f_a)^3 f_a + 2f_a^3(1 - f_a) + (1 - f_a)^2 f_a^2$
AM (Eq. 104)	mean	$\boxed{2 \sum_{a \in \mathcal{A}} F(a)} \quad (115)$ where $F(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a) + (1 - f_a)^2 f_a^2$
	variance	$\boxed{\sum_{a \in \mathcal{A}} [G(a) - 4F^2(a)]} \quad (115)$ where $F(a) = 2(1 - f_a)^3 f_a + 2f_a^3(1 - f_a) + (1 - f_a)^2 f_a^2$ and $G(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a) + 2(1 - f_a)^2 f_a^2$
TiTv (Eq. 105)	mean	$\boxed{(\gamma_0 + \gamma_2 + 2\gamma_1) \sum_{a \in \mathcal{A}} F(a) + \left[\frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1\right] \sum_{a \in \mathcal{A}} G(a)} \quad (131)$ where $F(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a)$ and $G(a) = (1 - f_a)^2 f_a^2$
	mean	$\boxed{\left[\frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1\right] \sum_{a \in \mathcal{A}} F(a) + \left[\frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1\right] \sum_{a \in \mathcal{A}} G(a) + \sum_{a \in \mathcal{A}} \left([\gamma_0 + \gamma_2 + 2\gamma_1]F(a) + \left[\frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1\right] G(a)\right)^2} \quad (131)$ where $F(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a)$ and $G(a) = (1 - f_a)^2 f_a^2$

## 4 Time series correlation-based distance distribution

For time series correlation-based data, we consider the case where there are  $m$  correlation matrices  $A^{(p \times p)}$ . In particular, we are focusing on resting-state fMRI (rs-fMRI) data, which falls into this category. The derivations that follow, however, are relevant to all correlation-based data fitting the assumptions we have adopted. The features in rs-fMRI are commonly Regions of Interest (ROIs), which are collections of highly correlated and spatially proximal voxels [19]. These correlations are between different ROIs for a particular brain atlas [20]. Because the features are the ROIs themselves, this leads us to the following metric

$$d_{ij}^{\text{ROI}}(a) = \sum_{k \neq a} |A_{ka}^{(i)} - A_{ka}^{(j)}|. \quad (134)$$

where  $A_{ka}^{(i)}$  and  $A_{ka}^{(j)}$  are the correlations between ROI  $a$  and ROI  $k$  for instances  $i$  and  $j$ , respectively. In order for comparisons between different correlations to be possible, we first perform a Fisher r-to-z transform on the correlations. We then load all of the

transformed correlations into a  $p(p-1) \times m$  matrix  $X$  (see Fig. 5).

567

$$\begin{array}{c}
 \text{ROI}_1 \left\{ \begin{array}{c} \hat{A}_{12}^{(1)} \quad \hat{A}_{12}^{(2)} \quad \hat{A}_{12}^{(3)} \quad \hat{A}_{12}^{(4)} \quad \dots \quad \hat{A}_{12}^{(m)} \\ \hat{A}_{13}^{(1)} \quad \hat{A}_{13}^{(2)} \quad \hat{A}_{13}^{(3)} \quad \hat{A}_{13}^{(4)} \quad \dots \quad \hat{A}_{13}^{(m)} \\ \vdots \quad \vdots \quad \vdots \quad \vdots \quad \dots \quad \vdots \\ \hat{A}_{1p}^{(1)} \quad \hat{A}_{1p}^{(2)} \quad \hat{A}_{1p}^{(3)} \quad \hat{A}_{1p}^{(4)} \quad \dots \quad \hat{A}_{1p}^{(m)} \end{array} \right. \\
 \text{ROI}_2 \left\{ \begin{array}{c} \hat{A}_{21}^{(1)} \quad \hat{A}_{21}^{(2)} \quad \hat{A}_{21}^{(3)} \quad \hat{A}_{21}^{(4)} \quad \dots \quad \hat{A}_{21}^{(m)} \\ \hat{A}_{23}^{(1)} \quad \hat{A}_{23}^{(2)} \quad \hat{A}_{23}^{(3)} \quad \hat{A}_{23}^{(4)} \quad \dots \quad \hat{A}_{23}^{(m)} \\ \vdots \quad \vdots \quad \vdots \quad \vdots \quad \dots \quad \vdots \\ \hat{A}_{2p}^{(1)} \quad \hat{A}_{2p}^{(2)} \quad \hat{A}_{2p}^{(3)} \quad \hat{A}_{2p}^{(4)} \quad \dots \quad \hat{A}_{2p}^{(m)} \end{array} \right. \\
 \vdots \\
 \text{ROI}_p \left\{ \begin{array}{c} \hat{A}_{p1}^{(1)} \quad \hat{A}_{p1}^{(2)} \quad \hat{A}_{p1}^{(3)} \quad \hat{A}_{p1}^{(4)} \quad \dots \quad \hat{A}_{p1}^{(m)} \\ \hat{A}_{p2}^{(1)} \quad \hat{A}_{p2}^{(2)} \quad \hat{A}_{p2}^{(3)} \quad \hat{A}_{p2}^{(4)} \quad \dots \quad \hat{A}_{p2}^{(m)} \\ \vdots \quad \vdots \quad \vdots \quad \vdots \quad \dots \quad \vdots \\ \hat{A}_{p,p-1}^{(1)} \quad \hat{A}_{p,p-1}^{(2)} \quad \hat{A}_{p,p-1}^{(3)} \quad \hat{A}_{p,p-1}^{(4)} \quad \dots \quad \hat{A}_{p,p-1}^{(m)} \end{array} \right.
 \end{array} = X$$

**Fig 5.** Resting-state fMRI transformed subject correlation matrices. Each column corresponds to an instance (or subject)  $I_j$  and each column corresponds to an ROI (or feature). The notation  $\hat{A}_{ka}^{(j)}$  represents the r-to-z transformed correlation between ROIs  $a$  and  $k \neq a$  for instance  $j$ .

568

We further transform the data matrix  $X$  by standardizing so that each of the  $m$  columns has zero mean and unit variance. Therefore, the data in matrix  $X$  are standard normal. Recall from Eqs. 42 and 43, that the mean and variance of the Manhattan ( $q = 1$ ) distance distribution for standard normal data are  $\frac{2p}{\sqrt{\pi}}$  and  $\frac{2(\pi-2)p}{\pi}$ , respectively. This allows us to easily derive the expected pairwise distance between instances  $i$  and  $j$  in rs-fMRI data as follows

569  
570  
571  
572  
573  
574

$$\begin{aligned}
 E(D_{ij}^{\text{fMRI}}) &= E \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{ROI}}(a) \right) \\
 &= E \left( \sum_{a \in \mathcal{A}} \sum_{k \neq a} |\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}| \right) \\
 &= \sum_{a \in \mathcal{A}} \sum_{k \neq a} E \left( |\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}| \right) \\
 &= \sum_{a \in \mathcal{A}} \sum_{k \neq a} \frac{2}{\sqrt{\pi}} \\
 &= \frac{2p(p-1)}{\sqrt{\pi}}.
 \end{aligned} \tag{135}$$

Due to the dependencies that exist between terms in the double sum when computing the rs-fMRI distance, linearity no longer applies to the variance operator. We proceed

575  
576

by writing the form of the variance as follows

577

$$\begin{aligned}
\text{Var}(D_{ij}^{\text{fMRI}}) &= \text{Var} \left( \sum_{a \in \mathcal{A}} \sum_{k \neq a} |\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}| \right) \\
&= \sum_{a=1}^{p-1} \text{Var} \left( \sum_{k=a+1}^p 2|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}| \right) \\
&\quad + 2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|, \sum_{s=r+1}^p 2|\hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)}| \right) \\
&= \sum_{a=1}^{p-1} \sum_{k=a+1}^p \text{Var} \left( 2|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}| \right) \\
&\quad + 2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|, \sum_{s=r+1}^p 2|\hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)}| \right) \quad (136) \\
&= \sum_{a=1}^{p-1} \sum_{k=a+1}^p \frac{4(\pi - 2)}{\pi} \\
&\quad + 2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|, \sum_{s=r+1}^p 2|\hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)}| \right) \\
&= \frac{2p(\pi - 2)(p - 1)}{\pi} \\
&\quad + 2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|, \sum_{s=r+1}^p 2|\hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)}| \right).
\end{aligned}$$

In order to have a formula in terms of the number of ROIs  $p$  only, we must estimate the double sum on the right-hand side of Eq. 136. Through simulation, it can be seen that the difference between the sample variance  $S_{D_{ij}}^2$  and  $\frac{2p(\pi-2)(p-1)}{\pi}$  has a quadratic relationship with  $p$ . More explicitly, we have the following relationship

578  
579  
580  
581

$$S_{D_{ij}}^2 - \frac{2p(\pi - 2)(p - 1)}{\pi} = \beta_1 p^2 + \beta_0 p. \quad (137)$$

The coefficient estimates found through least squares fitting are  $\beta_0 = -\beta_1 \approx 0.08$ . These estimates allow one to infer a functional form for the double sum in the right-hand side of Eq. 136 that is actually proportional to  $\frac{2p(\pi-2)(p-1)}{\pi}$ . That is, we have the following formula for approximating the double sum

582  
583  
584  
585

$$2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|, \sum_{s=r+1}^p 2|\hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)}| \right) = \frac{p(\pi - 2)(p - 1)}{4\pi}. \quad (138)$$

Therefore, the variance of the rs-fMRI distances is approximated well by the following

586

$$\text{Var}(D_{ij}^{\text{fMRI}}) = \frac{9p(\pi - 2)(p - 1)}{4\pi}. \quad (139)$$

With the mean and variance estimates given by Eqs. 135 and 139, we have the following asymptotic distribution for rs-fMRI distances

587  
588

$$D_{ij}^{\text{fMRI}} \sim \mathcal{N} \left( \frac{2p(p - 1)}{\sqrt{\pi}}, \frac{9p(\pi - 2)(p - 1)}{4\pi} \right). \quad (140)$$



Consider the max-min normalized rs-fMRI distance given by the following equation 589

$$D_{ij}^{\text{fMRI}*} = \sum_{a \in \mathcal{A}} \sum_{k \neq a} \frac{|A_{ak}^{(i)} - A_{ak}^{(j)}|}{\max(a) - \min(a)}. \quad (141)$$

Assuming that the data  $X$  has been r-to-z transformed and standardized, we can easily compute the expected attribute range and variance of the attribute range. The expected maximum of a given attribute in data matrix  $X$  is estimated by the following 590  
591  
592

$$\mathbb{E}(X_a^{\max} - X_a^{\min}) = 2\mu_{\max}^{(1)}(m, p) = 2 \left[ \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m(p-1)}\right)} - \Phi^{-1}\left(\frac{1}{m(p-1)}\right) \right]. \quad (142)$$

The variance can be esimated with the following 593

$$\text{Var}(X_a^{\max} - X_a^{\min}) = \frac{\pi^2}{6 \log[m(p-1)]}. \quad (143)$$

Let  $\mu_{D_{ij}^{\text{fMRI}}}$  and  $\sigma_{D_{ij}^{\text{fMRI}}}^2$  denote the mean and variance of the rs-fMRI distance distribution given by Eqs. 135 and 139. Using the formulas for the mean and variance of the max-min normalized distance distribution given in Eq. 86, we have the following asymptotic distribution for the max-min normalized rs-fMRI distances 594  
595  
596  
597

$$D_{ij}^{\text{fMRI}*} \sim \mathcal{N} \left( \frac{\mu_{D_{ij}^{\text{fMRI}}}}{2\mu_{\max}^{(1)}(m, p)}, \frac{6\sigma_{D_{ij}^{\text{fMRI}}}^2 \log[m(p-1)]}{\pi^2 + 24 \left[ \mu_{\max}^{(1)}(m, p) \right]^2 \log[m(p-1)]} \right). \quad (144)$$

#### 4.1 Normalized Manhattan ( $q = 1$ ) for rs-fMRI 598

Substituting the non-normalized mean given by Eq. 135 into Eq. 144 for the mean of the max-min normalized rs-fMRI metric, we have the following 599  
600

$$\begin{aligned} \mathbb{E}(D_{ij}^{\text{fMRI}*}) &= \frac{\mu_{D_{ij}^{\text{fMRI}}}}{2\mu_{\max}^{(1)}(m, p)} \\ &= \frac{p(p-1)}{\sqrt{\pi} \mu_{\max}^{(1)}(m, p)}, \end{aligned} \quad (145)$$

where  $\mu_{\max}^{(1)}(m, p)$  is given in Eq. 142. 601

Similarly, the variance of  $D_{ij}^{\text{fMRI}*}$  is given by 602

$$\begin{aligned} \text{Var}(D_{ij}^{\text{fMRI}*}) &= \frac{6\sigma_{D_{ij}^{\text{fMRI}}}^2 \log[m(p-1)]}{\pi^2 + 24 \left[ \mu_{\max}^{(1)}(m, p) \right]^2 \log[m(p-1)]} \\ &= \frac{27(\pi - 2) \log[m(p-1)](p-1)p}{2\pi \left( \pi^2 + 24 \left[ \mu_{\max}^{(1)}(m, p) \right]^2 \log[m(p-1)] \right)}, \end{aligned} \quad (146)$$

where  $\mu_{\max}^{(1)}(m, p)$  is given in Eq. 142. 603

**Table 5.** Summary of distance distribution derivations for rs-fMRI data.

rs-fMRI - Metric	Stat	Formula (Eq. #)
standard (Eq. 132)	mean	$\frac{2p(p-1)}{\sqrt{\pi}} \quad (139)$
	variance	$\frac{9p(\pi-2)(p-1)}{4\pi} \quad (139)$
max-min normalized (Eq. 140)	mean	$\frac{\mu_{D_{ij}}}{2\mu_{\max}^{(1)}(m, p)} \quad (143)$ <p>where <math>\mu_{D_{ij}}</math> and <math>\mu_{\max}^{(1)}(m, p)</math> are given by Eqs. 140 and 142</p>
	variance	$\frac{6\sigma_{D_{ij}}^2 \log[m(p-1)]}{\pi^2 + 24 \left[ \mu_{\max}^{(1)}(m, p) \right]^2 \log[m(p-1)]} \quad (143)$ <p>where <math>\sigma_{D_{ij}}^2</math> and <math>\mu_{\max}^{(1)}(m, p)</math> are given by Eqs. 140 and 142</p>

## 5 Effects of correlation on distances

All of the derivations presented in previous sections are for the cases where there is no correlation between instances or features. We assumed that any pair  $(X_{ia}, X_{ja})$  of data points for instances  $i$  and  $j$  and fixed feature  $a$  were independent and identically distributed. This was done in order to determine asymptotic estimates in null data. That is, data with no main effects, interaction effects, or pairwise correlations between features. Within this highly simplified context, our asymptotic formulas for distributional moments are reliable. However, correlations do exist between features and instances in real data. There are a multitude of different statistical effects that impact distance distributional properties. Ultimately, divergence from normality is caused primarily by large magnitude pairwise correlation between features. Pairwise feature correlation can be the result of main effects, where features have different within-group means. On the other hand, there could be an underlying interaction network in which there are strong associations between features. If features are differentially correlated between phenotype groups, then interactions exist that change affect the distance distribution. In the following few sections, we consider particular cases of the  $L_q$  metric for continuous and discrete data under the effects of pairwise feature correlation.

### 5.1 Continuous data

Without loss of generality, suppose that we have  $X^{(m \times p)}$  where  $X_{ia} \sim \mathcal{N}(0, 1)$  for all  $i = 1, 2, \dots, m$  and  $a = 1, 2, \dots, p$ , and let  $m = p = 100$  and consider only the  $L_2$  (Euclidean) metric (Eq. 1,  $q = 2$ ). We explore the effects of correlation on these distances by generating simulated data sets with increasing degree of pairwise feature correlation and then plotting the density curve of the induced distances (Fig. 6A). Divergence from normality in distances is directly related to the average absolute pairwise correlation

that exists in the simulated data. This measure is given by

$$\bar{r}_{\text{abs}} = \frac{2}{p(p-1)} \sum_{i=1}^{p-1} \sum_{j>i} r_{ij} \quad (147)$$

where  $r_{ij}$  is the correlation between features  $i, j \in \mathcal{I}$  across all instances  $m$ . Distances generated on data without correlation closely approximate a Gaussian. The mean and variance of the uncorrelated distance distribution are given by Eqs. 48 and 47, respectively, by substituting  $p = 100$  for the mean. As  $\bar{r}_{\text{abs}}$  increases, we very quickly see positive skewness and increased variability in distances. The predicted and sample means, however, are approximately the same between correlated and uncorrelated distances due to linearity of the expectation operator. Because of the dependencies between features, the predicted variance of 1 for  $L_2$  on standard normal data obviously no longer holds.

In order to introduce a controlled level of correlation between features, we created correlation matrices based on a random graph with specified connection probability, where features correspond to the vertices in each graph. We assigned high correlations to connected features from the random graph and low correlations to all non-connections. Using the upper-triangular cholesky factor  $U$  for uncorrelated data matrix  $X$ , we computed the following product to create correlated data matrix  $X^{\text{corr}}$

$$X^{\text{corr}} = XU^T. \quad (148)$$


The new data matrix given by Eq. 148 has approximately the same correlation structure as the randomly generated correlation matrix created from a random graph. The cholesky method is a standard approach in creating correlated data sets.

## 5.2 GWAS data

In analogy to the previous section, we explore the effects of pairwise feature correlation in the context of GWAS data. Without loss of generality, we let  $m = p = 100$  and consider only the TiTv metric, which is given by combining Eqs. 106 and 1 with  $q = 1$ . To create correlated GWAS data, we first generated standard normal data with random correlation structure, just as in the previous section. We then applied the standard normal cumulative distribution function (CDF) to this correlated data in order transform the correlated standard normal variates into uniform data with preserved correlation structure. We then subsequently applied the inverse binomial CDF to the correlated uniform data with random success probabilities  $f_a$  for all  $a \in \mathcal{A}$ . Each feature  $a \in \mathcal{A}$  corresponds to an individual SNP in the data matrix. The resulting GWAS data set is binomial with  $n = 2$  trials and has roughly the same correlation matrix as the original correlated standard normal data with which we started. Average absolute pairwise correlation  $\bar{r}_{\text{abs}}$  induces positive skewness in GWAS data at lower levels than in correlated standard normal data (Fig. 6B). This could have important implications in nearest neighborhoods in NPDR and other similar methods.

## 5.3 Correlation-based data

For our correlation data-based metric given by Eqs. 134 and 1 with  $q = 1$ , we consider additional effects of correlation between features. Without loss of generality, we let  $m = 100$  and  $p = 30$ . We show an illustration of the effects of correlated features in this context (Fig. 6C). Based on the correlated distance densities, it appears that correlation between features introduces positive skewness at lower values of  $\bar{r}_{\text{abs}}$ . We introduced correlation to the transformed data matrix (Fig. 5) with the cholesky method used previously.



**Fig 6.** Ridgeline plots of distance densities from uncorrelated vs correlated bioinformatics data. **(A)** Euclidean distance densities for random normal data with and without correlation. Correlated data was created by multiplying random normal data by upper-triangular cholsky factor from randomly generated correlation matrix. We created correlated data for average absolute pairwise correlation (Eq. 147)  $\bar{r}_{\text{abs}} = 0.105, 0.263, 0.458$ , and  $0.612$ . **(B)** TiTv distance densities for random binomial data with and without correlation. Correlated data was created by first generating correlated standard normal data using the cholsky method from (A). Then we applied the standard normal CDF to create correlated uniformly distributed data, which was then transformed by the inverse binomial CDF with  $n = 2$  trials and success probabilities  $f_a$  for all  $a \in \mathcal{A}$ . The result is correlated random binomial data with correlation matrix approximately the same as the original correlated standard normal data. **(C)** Time series correlation-based distance densities for random rs-fMRI data (Fig. 5) with and without additional pairwise feature correlation. Correlation was added to the transformed rs-fMRI data matrix (Fig. 5) using the cholsky algorithm from (A).

## 6 Discussion

Nearest-neighbor distance-based feature selection is class of methods that are relatively simple to implement, intuitive in nature, and perform surprisingly well in detecting interaction effects in high dimensional data. However, there has been little work done to understand how the limiting behavior of distance distributions can aid in determining how to properly parameterize these methods for feature selection. Furthermore, little has been done in the way of optimizing the choice of distance metric. Most often, distance-based feature selection methods use the  $L_q$  metric given by Eq. 1 with  $q = 1$  or  $q = 2$ . However, these two realizations of the  $L_q$  metric have considerably different expressions for the mean and variance of their respective limiting distributions. For instance, the expected distance for  $L_1$  and  $L_2$  on standard normal data is on the order of  $p$  (see Eq. 42) and  $\sqrt{p}$  (see Eq. 46), respectively. In addition,  $L_1$  and  $L_2$  on standard normal data have asymptotic variances on the order of  $p$  and 1, respectively. Considering whether one should choose  $L_1$  or  $L_2$  in this context may depend on motivation. For instance, distances become harder to distinguish from one another in high dimensions, which is one of the curses of dimensionality. In the case of  $L_2$ , unit variance in the limit distribution means that distances will be almost completely contained within a ball of radius 1. The limiting  $L_2$  distribution can therefore be thought of simply as a positive translation of the standard normal distribution. On the other hand, the  $L_1$  distances become more dispersed due to the fact that the variance of the limiting distribution is proportional to the feature dimension  $p$ . This could actually be more desirable when determining nearest neighbors because instances may be easier to distinguish with this metric. If using  $L_1$ , then it may be best to use a fixed-k algorithm instead of fixed-radius.

This is because fixed-radius neighborhood order could vary quite a bit considering the  $L_1$  variance is proportional to feature dimension  $p$ , which in turn could affect the quality of selected features. If  $L_2$  is being used, then perhaps either fixed-k or fixed-radius may perform equally well because most distances will be within 1 standard deviation away from the mean.

In any neighborhood selection algorithm, it is important to know what the average distance is and how dispersed these distances become as the feature dimension  $p$  grows. In our analysis, we have derived distance asymptotics for some of the most commonly used metrics in nearest-neighbor distance-based feature selection, as well as two new metrics for GWAS and time series correlation-based data like resting-state fMRI. Using extreme value theory, we have derived limiting distributions for the sample maximum and minimum of a fixed feature  $a$ . This has allowed us to determine the expected value and variance of the max-min normalized  $L_q$  distance in standard normal and standard uniform data, which is a new result to the best of our knowledge. Our derivations provide an important reference for individuals that are using nearest-neighbor feature selection methods in typical bioinformatics data.

In this work, we have expanded nearest-neighbor distance-based feature selection into the context of time series correlation-based data. Our motivation for this is partly based on the fact that these methods have not yet been applied to resting-state fMRI data. In order for this to be possible, we had to create a metric (see Eq. 134) that could allow us to have regions of interest (ROIs) as features. Not all ROIs will be relevant to a particular phenotype in case-control studies, so it could be important to use a nearest-neighbor feature selection method to determine which ROIs are important. This could allow us to detect interactions to help elucidate the network structure of the brain as it relates to the phenotype of interest.

The recently introduced transition-transversion metric given by Eq. 106 provides an additional dimension to the commonly used discrete metrics in GWAS nearest-neighbor distance-based feature selection. In this work, we have provided the asymptotic mean and variance of the limiting TiTv distance distribution. This novel result, as well as asymptotic estimates for the GM (see Eq. 104) and AM (see Eq. 105) metrics, provides an important reference to aid in neighborhood parameter selection in this context. We have also shown how the Ti/TV ratio  $\eta$  (see Eq. 118) and minor allele frequency (or success probability)  $f_a$  affects these discrete distances. For the GM and AM metrics, the distance is solely determined by the minor allele frequencies because the genotype encoding is not taken into account. We showed how both minor allele frequency and Ti/Tv ratio uniquely affects the TiTv distance (Figs. 4A and 4C). Because transversions are more drastic forms of mutation than transitions, this additional dimension of information is important to consider, which is why we have provided asymptotic results for this metric.

Correlations exist between features and instances in real data. Because of this, there can be rather drastic divergence from the asymptotic results for uncorrelated data we have derived in this work. To illustrate this behavior, we showed how strong correlations lead to positive skewness in the distance distribution of random normal, binomial, and rs-fMRI data (Figs. 6A, 6B, and 6C). Pairwise correlation between features does not change the average distance, so our asymptotic results for uncorrelated data also apply when features are not independent. In contrast, the sample variance of distances diverges from the uncorrelated case substantially as the average absolute pairwise feature-feature correlation increases (see Eq. 147). For fixed-radius neighborhood methods, this increases the probability of neighborhood inclusion for a fixed instance. The increased variability in distances on correlated data may provide further motivation for optimizing the choice of metric in nearest-neighbor feature selection. This most certainly motivates a discussion on optimal choices of neighborhood selection parameters, which we will address in future work.

There are many different distance metrics that can be used in place of those we have considered for bioinformatics data, but we have derived results for those that are the most commonly used in practice. Our work brings together many important aspects of nearest-neighbor distance-based feature selection, which also serves as a guide to other researchers that may be interested in a different choice of metric for a similar analysis. In future work, we will consider how pairwise feature correlation, as well as a mixture of main and interaction effects, changes the optimal choice of neighborhood selection parameters like fixed-k and fixed-radius.

## References

1. Ryan J. Urbanowicz, Randal S. Olson, Peter Schmitt, Melissa Meeker, and Jason H. Moore. Benchmarking relief-based feature selection methods for bioinformatics data mining. *Journal of Biomedical Informatics*, 85:168–188, 2018.
2. Ryan J. Urbanowicz, Melissa Meeker, William La Cava, Randal S. Olson, and Jason H. Moore. Relief-based feature selection: Introduction and review. *Journal of Biomedical Informatics*, 2018.
3. Marko Robnik Šikonja and Igor Kononenko. Theoretical and Empirical Analysis of ReliefF and RReliefF. *Machine Learning*, 53:23 – 69, February 2003.
4. M. Arabnejad, B. A. Dawkins, W. S. Bush, B. C. White, A. R. Harkness, and B. A. McKinney. Transition-transversion encoding and genetic relationship metfic in ReliefF feature selection improves pathway enrichment in GWAS. *BioData Mining*, 11(23), 2018.
5. Archana Venkataraman, Marek Kubicki, Carl-Fredrik Westin, and Polina Golland. Robust Feature Selection in Resting-State fMRI Connectivity Based on Population Studies. *Conf Comput Vis Pattern Recognit Workshops*, pages 63–70, 2010.
6. Etay Hay, Petra Ritter, Nancy J. Lobaugh, and Anthony R. McIntosh. Multiregional integration in the brain during resting-state fMRI activity. *PLOS Computational Biology*, March 2017.
7. Benedikt Sundermann, Mona Olde lütke Beverborg, and Bettina Pfleiderer. Toward literature-based feature selection for diagnostic classification: a meta-analysis of resting-state fMRI in depression. *Frontiers in Human Neuroscience*, September 2014.
8. Svyatoslav Vergun, Alok S. Deshpande, Timothy B. Meier, Jie Song, Dana L. Tudorascu, Veena A. Nair, Vikas Singh, Bharat B. Biswal, M. Elizabeth Meverand, Rasmus M. Birn, and Vivek Prabhakaran. Characterizing functional connectivity differences in aging adults using machine learning on resting state fMRI data. *Frontiers in Computational Neuroscience*, April 2013.
9. Stephen J. Gotts, W. Kyle Simmons, Lydia A. Milbury, Gregory L. Wallace, Robert W. Cox, and Alex Martin. Fractionation of social brain circuits in autism spectrum disorders. *Brain*, 135:2711–2725, 2012.
10. Trang T Le, Ryan J Urbanowicz, Jason H Moore, and Brett A McKinney. Statistical inference relief (stir) feature selection. *Bioinformatics*, page bty788, 2018.
11. Brett A McKinney, Bill C White, Diane E Grill, Peter W Li, Richard B Kennedy, Gregory A Poland, and Ann L Oberg. ReliefSeq: a gene-wise adaptive-K nearest-neighbor feature selection tool for finding gene-gene interactions and main effects in mRNA-Seq gene expression data. *PLoS one*, 8(12):e81527, 2013.

12. Larry Wasserman. *All of Statistics: A Concise Course in Statistical Inference*. 790  
Springer, New York, NY, 2004. 791
13. Irwin Miller and Marylees Miller. *John E. Freund's Mathematical Statistics with* 792  
*Applications*. Pearson Prentice Hall, 7 edition, 2004. 793
14. Alvis Brazma and Jaak Vilo. Gene expression data analysis. *FEBS Letters*, 794  
480:17–24, June 2000. 795
15. Dongfang Wang and Jin Gu. VASC: Dimension Reduction and Visualization of 796  
Single-cell RNA-seq Data by Deep Variational Autoencoder. *Genomics Proteomics* 797  
*Bioinformatics*, 16:320–331, December 2018. 798
16. E. J. Gumbel. The Distribution of the Range. *The Annals of Mathematical* 799  
*Statistics*, 18(3):384–412, September 1947. 800
17. Sourav Chatterjee. *Superconcentration and Related Topics*. 1439-7382. Springer 801  
International Publishing, 1 edition, 2014. 802
18. Harald Cramér. *Mathematical Methods of Statistics*, volume 1. Princeton University 803  
Press, reprint, revised edition, 1999. 804
19. Megan H. Lee, Christopher D. Smyser, and Joshua S. Shimony. Resting state 805  
fMRI: A review of methods and clinical applications. *AJNR Am J Neuroradiol.*, 806  
34(10):1866–1872, October 2013. 807
20. David Alexander Dickie, Susan D. Shenkin, Devasuda Anblagan, Juyoung Lee, 808  
Manuel Blesa Cabez, David Rodriguez, James P. Boardman, Adam Waldman, 809  
Dominic E. Job, and Joanna M. Wardlaw. Whole Brain Magnetic Resonance 810  
Image Atlases: A Systematic Review of Existing Atlases and Caveats for Use in 811  
Population Imaging. *Frontiers in Neuroinformatics*, January 2017. 812