

RESEARCH

# Theoretical properties of nearest-neighbor distance distributions and novel metrics for high dimensional bioinformatics data

Bryan A Dawkins<sup>1</sup>, Trang T Le<sup>2</sup> and Brett A McKinney<sup>1,3\*</sup>

\*Correspondence:

[brett-mckinney@utulsa.edu](mailto:brett-mckinney@utulsa.edu)

<sup>3</sup>Tandy School of Computer

Science, University of Tulsa,

74104 Tulsa, United States

Full list of author information is  
available at the end of the article

## Abstract

The performance of nearest-neighbor feature selection and prediction methods depends on the metric for computing neighborhoods and the distribution type of the underlying data. The effects of the distribution and metric, as well as the presence of correlation and interactions, are reflected in the expected moments of the distribution of pairwise distances. We derive general analytical expressions for the mean and variance of pairwise distances for  $L_q$  metrics for Gaussian and uniform data with  $p$  attributes and  $m$  instances. We use extreme value theory to derive results for metrics that are normalized by the range of each attribute (max - min). In addition to these expressions applicable to the analysis of continuous data such as gene expression, we derive similar analytical expressions for a new metric for genetic variant (categorical) data in genome-wide association studies (GWAS). The distance distributions account for minor allele frequency and transition/transversion ratio. We introduce a new metric for resting-state fMRI data that is applicable to correlation-based predictors derived from time series data. Derivations assume independent data, but empirically we also consider the effect of correlation.

**Keywords:** Nearest neighbor methods; Distance distributions; Feature selection; Relief-based algorithms; High-dimensional data; Gene expression; Genome-wide association studies; Resting-state fMRI

## 1 Background

Statistical models can deviate from expected behavior depending on whether certain properties of the underlying data are satisfied, such as being normally dis-

tributed. The expected behavior of nearest neighbor models is further influenced<sup>1</sup>  
 by the choice of metric, such as Euclidean or Manhattan. For random normal data<sup>2</sup>  
 $(\mathcal{N}(0, 1))$ , for example, the variance of the pairwise distances of a Manhattan metric<sup>3</sup>  
 is proportional to the number of attributes ( $p$ ) whereas the variance is constant for a<sup>4</sup>  
 Euclidean metric. Relief methods [1–3] and nearest-neighbor projected distance re-<sup>5</sup>  
 gression (NDPR) [4] use nearest neighbors to compute attribute importance scores<sup>6</sup>  
 and often use adaptive neighborhoods that rely on the mean and variance of the dis-<sup>7</sup>  
 tance distribution. The ability of this class of methods to identify association effects,<sup>8</sup>  
 like main effects or interaction effects, depends on parameters such as neighborhood<sup>9</sup>  
 radii or number of neighbors  $k$  [5, 6]. Thus, knowledge of the expected values for a<sup>10</sup>  
 given metric and data distribution may improve the performance of these feature<sup>11</sup>  
 selection methods by informing the selection of neighborhood parameters.<sup>12</sup>

For continuous data, the metrics most commonly used in nearest neighbor meth-<sup>13</sup>  
 ods are  $L_q$  with  $q = 1$  (Manhattan) or  $q = 2$  (Euclidean). For data from standard<sup>14</sup>  
 normal  $(\mathcal{N}(0, 1))$  or standard uniform  $(\mathcal{U}(0, 1))$  distributions, the asymptotic be-<sup>15</sup>  
 havior of the  $L_q$  metrics is known. However, detailed derivations of these distance<sup>16</sup>  
 distribution asymptotics are not readily available in the literature. We provide de-<sup>17</sup>  
 tailed derivations of generalized expressions parameterized by metric  $q$ , attributes<sup>18</sup>  
 $p$ , and samples  $m$ , and we extend the derivations to  $L_q$  metrics normalized by the<sup>19</sup>  
 range of the attributes using Extreme Value Theory (EVT). These range (max-min)<sup>20</sup>  
 normalized metrics are often used in Relief-based algorithms [3].<sup>21</sup>

For genome-wide association study (GWAS) data, which is categorical, various<sup>22</sup>  
 metrics have been developed for feature selection and for computing similarity be-<sup>23</sup>  
 tween individuals based on shared genetic variation. We build on the mathematical<sup>24</sup>  
 formalism for continuous data to derive the asymptotic properties of various cate-<sup>25</sup>  
 gorical data metrics for GWAS. We derive asymptotic formulas for the mean and<sup>26</sup>  
 variance for three recently introduced GWAS metrics [7]. These metrics were de-<sup>27</sup>  
 veloped for Relief-based feature selection to account for binary genotype differences<sup>28</sup>  
 (two levels), allelic differences (three levels), and transition/transversion differences<sup>29</sup>  
 (five levels). The mean and variance expressions we derive for these multi-level<sup>30</sup>  
 categorical data types are parameterized by the minor allele frequency and the<sup>31</sup>  
 transition/transversion ratio.<sup>32</sup>

Resting-state fMRI (rs-fMRI) is a growing application area for machine learning<sup>1</sup> and feature selection [8–11], which involves correlation data derived from time-series<sup>2</sup> brain activity. For a given subject, a correlation or similar matrix is computed be-<sup>3</sup> tween brain regions of interest (ROIs) from their time series. Each time series rep-<sup>4</sup> resents functional activity of the ROI while the subject is not performing any task,<sup>5</sup> and the ROI typically corresponds to a region with known function for emotion<sup>6</sup> or cognition. Thus, the dataset consists of pairwise ROI correlations for each of<sup>7</sup> the  $m$  subjects. Nearest-neighbor based feature selection was applied to rs-fMRI<sup>8</sup> in the private evaporative cooling method [12], where the predictors were pairwise<sup>9</sup> correlations between ROIs. The use of pairwise correlation predictors is a common<sup>10</sup> practice because of convenience and differential connectivity between brain regions<sup>11</sup> may be of biological importance [13]. However, one may be interested in the im-<sup>12</sup> portance of attributes at the ROI level. Thus, in the current study we introduce<sup>13</sup> a new metric to be used in NPDR [4] with resting state correlation matrices that<sup>14</sup> provides attribute importance for ROIs. This metric is applicable to general time<sup>15</sup> series derived correlation data, and we derive asymptotic estimates for the mean<sup>16</sup> and variance of distance distributions for our new ts-corr based metric.<sup>17</sup>

In Section 2, we introduce preliminary notation and apply the Central Limit The-<sup>18</sup> orem (CLT) and the Delta Method to derive asymptotics for pairwise distances. In<sup>19</sup> Section 3, we present general derivations for continuously distributed data sets with<sup>20</sup>  $m$  instances and  $p$  attributes. We begin with the cases of standard normal ( $\mathcal{N}(0, 1)$ )<sup>21</sup> and standard uniform ( $\mathcal{U}(0, 1)$ ) data distributions, but we derive analytical expres-<sup>22</sup> sions parameterized by  $q$ ,  $p$ , and  $m$ . In Section 3.4 we use Extreme Value Theory<sup>23</sup> (EVT) to derive attribute range-normalized (max-min) versions of  $L_q$  metrics. In<sup>24</sup> Section 4, we extend the derivations to categorical data with a binomial distribution<sup>25</sup> for GWAS data with multiple metric types. In Section 5, we present a new time se-<sup>26</sup> ries correlation-based distance metric, with a particular emphasis on rs-fMRI data,<sup>27</sup> and we derive the corresponding asymptotic distance distribution results. Lastly, in<sup>28</sup> Section 7, we demonstrate the effect of correlation in the attribute space on distance<sup>29</sup> distributional properties.<sup>30</sup>

31

31

32

32

33

33

## 12 Limit distribution for $L_q$ on null data

In the application of nearest-neighbor distance-based methods to continuous data, the distance between instances  $(i, j \in \mathcal{I}, |\mathcal{I}| = m)$  in the data set  $X^{m \times p}$  of  $m^3$  instances and  $p$  attributes (or features) is calculated in the space of all attributes  $(a \in \mathcal{A}, |\mathcal{A}| = p)$  using a metric such as

$$D_{ij}^{(q)} = \left( \sum_{a \in \mathcal{A}} |d_{ij}(a)|^q \right)^{1/q}, \quad (1)$$

which is typically Manhattan ( $q = 1$ ) in Relief-based methods but may also be Euclidean ( $q = 2$ ). We use the terms “feature” and “attribute” interchangeably for the remainder of this work. The quantity  $d_{ij}(a)$ , known as a “diff” in Relief literature, is the projection of the distance between instances  $i$  and  $j$  onto the attribute  $a_{12}$  dimension. The function  $d_{ij}(a)$  supports any type of attributes (e.g., numeric and categorical). For example, the projected difference between two instances  $i$  and  $j$  for a continuous numeric ( $d^{\text{num}}$ ) attribute  $a$  may be

$$\begin{aligned} d_{ij}^{\text{num}}(a) &= \text{diff}(a, (i, j)) \\ &= |\hat{X}_{ia} - \hat{X}_{ja}|, \end{aligned} \quad (2)$$

where  $\hat{X}$  represents the standardized data matrix  $X$ . We use a simplified  $d_{ij}(a)$  notation in place of the  $\text{diff}(a, (i, j))$  notation that is customary in Relief-based methods. In NPDR, we omit the division by  $\max(a) - \min(a)$  used by Relief to constrain scores to the interval from  $-1$  to  $1$ , where  $\max(a) = \max_{k \in \mathcal{I}} \{X_{ka}\}$  and  $\min(a) = \min_{k \in \mathcal{I}} \{X_{ka}\}$ . The numeric  $d_{ij}^{\text{num}}(a)$  projection is simply the absolute difference between row elements  $i$  and  $j$  of the data matrix  $X^{m \times p}$  for the attribute column  $a$ .

All derivations in the following sections are applicable to nearest-neighbor distance-based methods in general, which includes not only NPDR, but also Relief-based algorithms. Each of these methods uses a distance metric (Eq. 1) to compute neighbors for each instance  $i \in \mathcal{I}$ . Therefore, our derivations of asymptotic distance distributions are applicable to all methods that compute neighbors in order to weight attributes. The predictors used by NPDR, however, are the one-dimensional projected distances between two instances  $i, j \in \mathcal{I}$  (Eq. 2). Hence, all asymptotic estimates we derive for diff metrics (Eq. 2) are particularly relevant to NPDR. Since

<sup>1</sup>the standard distance metric (Eq. 1) is a function of the one-dimensional projection<sup>1</sup>  
<sup>2</sup>(Eq. 2), asymptotic estimates derived for this projection (Eq. 2) are implicitly rele-<sup>2</sup>  
<sup>3</sup>vant to older nearest-neighbor distance-based methods like Relief-based algorithms.<sup>3</sup>  
<sup>4</sup>We proceed in the following section by applying the Classical Central Limit Theo-<sup>4</sup>  
<sup>5</sup>rem and the Delta Method to derive the limit distribution of pairwise distances on<sup>5</sup>  
<sup>6</sup>any data distribution that is induced by the standard distance metric (Eq. 1).<sup>6</sup>

## 82.1 Asymptotic normality of pairwise distances

<sup>9</sup>Suppose that

<sup>10</sup> $X_{ia}, X_{ja} \stackrel{iid}{\sim} \mathcal{F}_X(\mu_X, \sigma_X^2)$  for two fixed and distinct instances  $i, j \in \mathcal{I}$  and a fixed<sup>10</sup>  
<sup>11</sup>attribute  $a \in \mathcal{A}$ .  $\mathcal{F}_X$  represents any data distribution with mean  $\mu_X$  and variance<sup>11</sup>  
<sup>12</sup> $2\sigma_X^2$ .<sup>12</sup>

<sup>13</sup>It is clear that  $|X_{ia} - X_{ja}|^q = |d_{ij}(a)|^q$  is another random variable. Let  $Z_a^q \sim$ <sup>13</sup>  
<sup>14</sup> $\mathcal{F}_{Z_a^q}(\mu_{z_a^q}, \sigma_{z_a^q}^2)$  be the random variable such that<sup>14</sup>

$$Z_a^q = |d_{ij}(a)|^q = |X_{ia} - X_{ja}|^q, \quad a \in \mathcal{A}. \quad (3)$$

<sup>17</sup>Furthermore, the collection  $\{Z_a^q | a \in \mathcal{A}\}$  is a random sample of size  $p$  of mutually<sup>17</sup>  
<sup>18</sup>independent random variables. Hence, the sum of  $Z_a^q$  over all  $a \in \mathcal{A}$  is asymptoti-<sup>18</sup>  
<sup>19</sup>cally normal by the Classical Central Limit Theorem (CCLT). More explicitly, this<sup>19</sup>  
<sup>20</sup>implies that<sup>20</sup>

$$\left(D_{ij}^{(q)}\right)^q = \sum_{a \in \mathcal{A}} |d_{ij}(a)|^q = \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q = \sum_{a \in \mathcal{A}} Z_a^q \sim \mathcal{N}(\mu_{z_a^q} p, \sigma_{z_a^q}^2 p). \quad (4)$$

<sup>24</sup>Consider the smooth function  $g(z) = z^{1/q}$  that is continuously differentiable for<sup>24</sup>  
<sup>25</sup> $z > 0$ . Assuming that  $\mu_{z_a^q} > 0$ , the Delta Method [14] can be applied to show that<sup>25</sup>

$$\begin{aligned} g\left(\left(D_{ij}^{(q)}\right)^q\right) &= g\left(\sum_{a \in \mathcal{A}} Z_a^q\right) \\ &= \left(\sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q\right)^{1/q} \\ &= D_{ij}^{(q)} \sim \mathcal{N}\left(g(\mu_{z_a^q} p), [g'(\mu_{z_a^q} p)]^2 \sigma_{z_a^q}^2 p\right) \\ \Rightarrow D_{ij}^{(q)} &\sim \mathcal{N}\left((\mu_{z_a^q} p)^{1/q}, \frac{\sigma_{z_a^q}^2 p}{q^2 (\mu_{z_a^q} p)^{2(1-\frac{1}{q})}}\right). \end{aligned} \quad (5)$$


<sup>1</sup> Therefore, the distance between two fixed, distinct instances  $i$  and  $j$  given by<sup>1</sup>  
<sup>2</sup>Eq. 1 is asymptotically normal. Specifically, when  $q = 2$ , the distribution of  $D_{ij}^{(2)}$ <sup>2</sup>  
<sup>3</sup>asymptotically approaches  $\mathcal{N}\left(\sqrt{\mu_{z_a^2} p}, \frac{\sigma_{z_a^2}^2}{4\mu_{z_a^2}}\right)$ . When  $p$  is small, however, we observe<sup>3</sup>  
<sup>4</sup>empirically that a closer estimate of the sample mean is<sup>4</sup>

$$\begin{aligned} \text{E}\left(D_{ij}^{(2)}\right) &= \sqrt{\text{E}\left[\left(D_{ij}^{(2)}\right)^2\right] - \text{Var}\left(D_{ij}^{(2)}\right)} \\ &= \sqrt{\mu_{z_a^2} p - \frac{\sigma_{z_a^2}^2}{4\mu_{z_a^2}}}. \end{aligned} \quad (6)$$

<sup>9</sup>We estimate rate of convergence to normality for Euclidean ( $q = 2$ ) and Manhat-<sup>9</sup>  
<sup>10</sup>tan ( $q = 1$ ) metrics by comparing the distribution of pairwise distances in simu-<sup>10</sup>  
<sup>11</sup>lated data to a Gaussian (Fig. 1). We compute the distance between all pairs of<sup>11</sup>  
<sup>12</sup>instances in simulated datasets of uniformly distributed random data. We simu-<sup>12</sup>  
<sup>13</sup>late data with fixed  $m = 100$  instances, and, by varying the number of attributes<sup>13</sup>  
<sup>14</sup>( $p = 10, 100, 10000$ ), we observe rapid convergence to Gaussian. For  $p$  as low as 10<sup>14</sup>  
<sup>15</sup>attributes, Gaussian is a good approximation. The number of attributes in bioin-<sup>15</sup>  
<sup>16</sup>formatics data is typically quite large, at least on the order of  $10^3$ . The Euclidean<sup>16</sup>  
<sup>17</sup>metric has stronger convergence to a Gaussian than Manhattan. This may be due<sup>17</sup>  
<sup>18</sup>to Euclidean's use of the square root, which is a common transformation of data in<sup>18</sup>  
<sup>19</sup>statistics. Normality was assessed using the Shapiro-Wilk test.<sup>19</sup>

<sup>20</sup>To show asymptotic normality of distances, we did not specify whether the data<sup>20</sup>  
<sup>21</sup>distribution  $\mathcal{F}_X$  was discrete or continuous. This is because asymptotic normality<sup>21</sup>  
<sup>22</sup>is a general phenomenon in high attribute dimension  $p$  for any data distribution<sup>22</sup>  
<sup>23</sup> $\mathcal{F}_X$  satisfying the assumptions we have made. Therefore, the simulated distances<sup>23</sup>  
<sup>24</sup>we have shown (Fig. 1) has an analogous representation for discrete data, as well<sup>24</sup>  
<sup>25</sup>as all other continuous data distributions.<sup>25</sup>

<sup>26</sup>For distance based learning methods, all pairwise distances are used to determine<sup>26</sup>  
<sup>27</sup>relative importances for attributes. The collection of all distances above the diagonal<sup>27</sup>  
<sup>28</sup>in an  $m \times m$  distance matrix does not satisfy the independence assumption used in<sup>28</sup>  
<sup>29</sup>the previous derivations. This is because of the redundancy that is inherent to the<sup>29</sup>  
<sup>30</sup>distance matrix calculation. However, this collection is still asymptotically normal<sup>30</sup>  
<sup>31</sup>with mean and variance approximately equal to those we have previously given<sup>31</sup>  
<sup>32</sup>(Eq. 5). In the next section, we assume actual data distributions in order to define<sup>32</sup>  
<sup>33</sup>more specific general formulas for standard  $L_q$  and max-min normalized  $L_q$  metrics.<sup>33</sup>



**Figure 1** Convergence to Gaussian for Manhattan and Euclidean distances for simulated standard uniform data with  $m = 100$  instances and  $p = 10, 100$ , and  $10000$  attributes.

Convergence to Gaussian occurs rapidly with increasing  $p$ , and Gaussian is a good approximation for  $p$  as low as 10 attributes. The number of attributes in bioinformatics data is typically much larger, at least on the order of  $10^3$ . The Euclidean metric has stronger convergence to normal than Manhattan. P values from Shapiro-Wilk test, where the null hypothesis is a Gaussian distribution.

We also derive asymptotic moments for a new discrete metric in GWAS data and a new metric for time series correlation-based data, such as, resting-state fMRI.

### 3 $L_q$ metric moments for continuous data distributions

In this section, we begin by deriving general formulas for asymptotic means and variances of the  $L_q$  distance (Eq. 1) for standard normal and standard uniform data.

With our general formulas for continuous data, we compute moments associated with Manhattan ( $L_1$ ) and Euclidean ( $L_2$ ) metrics. We then consider the max-min normalized version of the  $L_q$  distance, where the magnitude difference (Eq. 2) is

<sup>1</sup>divided by the range of each attribute  $a$ . Using Extreme Value Theory (EVT), we<sup>1</sup>  
<sup>2</sup>derive formulas for the moments of attribute range in standard normal and standard<sup>2</sup>  
<sup>3</sup>uniform data. Transitioning into discrete data distributions relevant to GWAS, we<sup>3</sup>  
<sup>4</sup>derive asymptotic moments for two well known metrics and one new metric. In<sup>4</sup>  
<sup>5</sup>addition, we derive distance asymptotics for time series correlation-based data, such<sup>5</sup>  
<sup>6</sup>as, resting-state fMRI. 6

### <sup>7</sup>3.1 Distribution of $|d_{ij}(a)|^q = |X_{ia} - X_{ja}|^q$ 7

<sup>8</sup>Suppose that  $X_{ia}, X_{ja} \stackrel{iid}{\sim} \mathcal{F}_X(\mu_x, \sigma_x^2)$  and define  $Z_a^q = |d_{ij}(a)|^q = |X_{ia} - X_{ja}|^q$ ,<sup>8</sup>  
<sup>9</sup>where  $a \in \mathcal{A}$  and  $|\mathcal{A}| = p$ . In order to find the distribution of  $Z_a^q$ , we will use the<sup>9</sup>  
<sup>10</sup>following theorem given in [15]. 10

<sup>11</sup>**Theorem 3.1** *Let  $f(x)$  be the value of the probability density of the continuous*  
<sup>12</sup>*random variable  $X$  at  $x$ . If the function given by  $y = u(x)$  is differentiable and either*  
<sup>13</sup>*increasing or decreasing for all values within the range of  $X$  for which  $f(x) \neq 0$ ,*  
<sup>14</sup>*then, for these values of  $x$ , the equation  $y = u(x)$  can be uniquely solved for  $x$  to give*  
<sup>15</sup> *$x = w(y)$ , and for the corresponding values of  $y$  the probability density of  $Y = u(X)$*   
<sup>16</sup>*is given by* 16

$$g(y) = f[w(y)] \cdot |w'(y)| \quad \text{provided } u'(x) \neq 0$$

<sup>17</sup>Elsewhere,  $g(y) = 0$ . 17

<sup>18</sup>We have the following cases that result from solving for  $X_{ja}$  in the equation given<sup>18</sup>  
<sup>19</sup>by  $Z_a^q = |X_{ia} - X_{ja}|^q$ : 19

<sup>20</sup>(i) Suppose that  $X_{ja} = X_{ia} - (Z_a^q)^{1/q}$ . Based on the iid assumption for  $X_{ia}$  and<sup>20</sup>  
<sup>21</sup> $X_{ja}$ , it follows from Thm. 3.1 that the joint density function  $g^{(1)}$  of  $X_{ia}$  and<sup>21</sup>  
<sup>22</sup> $Z_a^q$  is given by 22

$$\begin{aligned} g^{(1)}(x_{ia}, z_a) &= f_X(x_{ia}, x_{ja}) \left| \frac{\partial x_{ja}}{\partial z_a} \right| \\ &= f_X(x_{ia}) f_X(x_{ja}) \left| \frac{-1}{q} (z_a^q)^{\frac{1}{q}-1} \right| \\ &= \frac{1}{q (z_a^q)^{1-\frac{1}{q}}} f_X(x_{ia}) f_X(x_{ia} - (z_a^q)^{1/q}), \quad z_a > 0 \end{aligned} \quad (7)$$



The density function  $f_{Z_a^q}^{(1)}$  of  $Z_a^q$  is then defined as

$$\begin{aligned} f_{Z_a^q}^{(1)}(z_a^q) &= \int_{-\infty}^{\infty} g^{(1)}(x_{ia}, z_a^q) dx_{ia} \\ &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{\infty} f_X(x_{ia}) f_X\left(x_{ia} - (z_a^q)^{1/q}\right) dx_{ia}, \quad z_a > 0. \end{aligned} \quad (8)$$

(ii) Suppose that  $X_{ja} = X_{ia} + (Z_a^q)^{1/q}$ . Based on the iid assumption for  $X_{ia}$  and  $X_{ja}$ , it follows from Thm. 3.1 that the joint density function  $g^{(2)}$  of  $X_{ia}$  and  $Z_a$  is given by

$$\begin{aligned} g^{(2)}(x_{ia}, z_a) &= f_X(x_{ia}, x_{ja}) \left| \frac{\partial x_{ja}}{\partial z_a} \right| \\ &= f_X(x_{ia}) f_X(x_{ja}) \left| \frac{1}{q} (z_a^q)^{\frac{1}{q}-1} \right| \\ &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} f_X(x_{ia}) f_X\left(x_{ia} - (z_a^q)^{1/q}\right), \quad z_a > 0. \end{aligned} \quad (9)$$

The density function  $f_{Z_a^q}^{(2)}$  of  $Z_a^q$  is then defined as

$$\begin{aligned} f_{Z_a^q}^{(2)}(z_a^q) &= \int_{-\infty}^{\infty} g^{(2)}(x_{ia}, z_a^q) dx_{ia} \\ &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{\infty} f_X(x_{ia}) f_X\left(x_{ia} + (z_a^q)^{1/q}\right) dx_{ia}, \quad z_a > 0. \end{aligned} \quad (10)$$

Let  $F_{Z_a^q}$  denote the distribution function of the random variable  $Z_a^q$ . Furthermore, we define the events  $E^{(1)}$  and  $E^{(2)}$  as

$$E^{(1)} = \{|X_{ia} - X_{ja}|^q \leq z_a^q : X_{ja} = X_{ia} - (Z_a^q)^{1/q}\} \quad (11)$$

and

$$E^{(2)} = \{|X_{ia} - X_{ja}|^q \leq z_a^q : X_{ja} = X_{ia} + (Z_a^q)^{1/q}\}. \quad (12)$$

Then it follows from fundamental rules of probability that

$$\begin{aligned}
 F_{Z_a^q}(z_a^q) &= \mathbb{P}[Z_a^q \leq z_a^q] \\
 &= \mathbb{P}[|X_{ia} - X_{ja}|^q \leq z_a^q] \\
 &= \mathbb{P}[E^{(1)} \cup E^{(2)}] \\
 &= \mathbb{P}[E^{(1)}] + \mathbb{P}[E^{(2)}] - \mathbb{P}[E^{(1)} \cap E^{(2)}] \\
 &= \mathbb{P}[E^{(1)}] + \mathbb{P}[E^{(2)}] \\
 &= \int_{-\infty}^{z_a^q} f_{Z_a^q}^{(1)}(t) dt + \int_{-\infty}^{z_a^q} f_{Z_a^q}^{(2)}(t) dt \\
 &= \int_{-\infty}^{z_a^q} (f_{Z_a^q}^{(1)}(t) + f_{Z_a^q}^{(2)}(t)) dt \\
 &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{z_a^q} \left\{ \left( \int_{-\infty}^{\infty} f_X(x_{ia}) [f_X(x_{ia} - t) \right. \right. \\
 &\quad \left. \left. + f_X(x_{ia} + t)] dx_{ia} \right) \right\} dt,
 \end{aligned} \tag{13}$$

where  $z_a > 0$ .

It follows directly from the previous result (Eq. 13) that the density function of the random variable  $Z_a^q$  is given by

$$\begin{aligned}
 f_{Z_a^q}(z_a^q) &= \frac{\partial}{\partial z_a^q} F_{Z_a^q}(z_a^q) \\
 &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{\infty} \left\{ f_X(x_{ia}) \left[ f_X(x_{ia} - (z_a^q)^{1/q}) \right. \right. \\
 &\quad \left. \left. + f_X(x_{ia} + (z_a^q)^{1/q}) \right] \right\} dx_{ia},
 \end{aligned} \tag{14}$$

where  $z_a > 0$ .

Using the previous result (Eq. 14), we can compute the mean and variance of the random variable  $Z_a^q$  as

$$\mu_{z_a^q} = \int_{-\infty}^{\infty} z_a^q f_{Z_a^q}(z_a^q) dz_a^q \tag{15}$$

and

$$\sigma_{z_a^q}^2 = \int_{-\infty}^{\infty} (z_a^q)^2 f_{Z_a^q}(z_a^q) dz_a^q - \mu_{z_a^q}^2. \tag{16}$$

<sup>1</sup> It follows immediately from the mean (Eq. 15) and variance (Eq. 16) and the<sup>1</sup>  
<sup>2</sup>Classical Central Limit Theorem (CCLT) that<sup>2</sup>

$$\left(D_{ij}^{(q)}\right)^q = \sum_{a \in \mathcal{A}} Z_a^q = \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q \sim \mathcal{N}(\mu_{z_a^q} p, \sigma_{z_a^q}^2 p). \quad (17)$$

Applying the convergence result we derived previously (Eq. 5), the distribution<sup>3</sup>  
of  $D_{ij}^{(q)}$  is given by<sup>4</sup>

$$D_{ij}^{(q)} \sim \mathcal{N}\left((\mu_{z_a^q} p)^{1/q}, \frac{\sigma_{z_a^q}^2 p}{q^2 (\mu_{z_a^q} p)^{2(1-\frac{1}{q})}}\right), \quad \mu_{z_a^q} > 0, \quad (18)$$

where we have an improved estimate of the mean for  $q = 2$  (Eq. 6).

### 3.1.1 Standard normal data

If  $X_{ia}, X_{ja} \stackrel{iid}{\sim} \mathcal{N}(0, 1)$ , then the marginal density functions with respect to  $X$  for<sup>5</sup>  
 $X_{ia}$ ,  $X_{ia} - (Z_a^q)^{1/q}$ , and  $X_{ia} + (Z_a^q)^{1/q}$  are defined as<sup>6</sup>

$$f_X(x_{ia}) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_{ia}^2}, \quad (19)$$

$$f_X\left(x_{ia} - (z_a^q)^{1/q}\right) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(x_{ia} - (z_a^q)^{1/q})^2}, \quad z_a > 0, \text{ and} \quad (20)$$

$$f_X\left(x_{ia} + (z_a^q)^{1/q}\right) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(x_{ia} + (z_a^q)^{1/q})^2}, \quad z_a > 0. \quad (21)$$

Substituting these marginal densities (Eqs. 19-21) into the general density func-  
tion for  $Z_a^q$  (Eq. 14) and completing the square on  $x_{ia}$  in the exponents, we have

$$\begin{aligned}
f_{Z_a^q}(z_a^q) &= \frac{1}{2q\pi (z_a^q)^{1-\frac{1}{q}}} e^{-\frac{1}{4}(z_a^q)^{2/q}} \int_{-\infty}^{\infty} \left( e^{-\frac{1}{2}[\sqrt{2}x_{ia} - \frac{\sqrt{2}}{2}(z_a^q)^{1/q}]^2} \right. \\
&\quad \left. + e^{-\frac{1}{2}[\sqrt{2}x_{ia} + \frac{\sqrt{2}}{2}(z_a^q)^{1/q}]^2} \right) dx_{ia} \\
&= \frac{1}{2q\sqrt{\pi} (z_a^q)^{1-\frac{1}{q}}} e^{-\frac{1}{4}(z_a^q)^{2/q}} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \left( e^{-\frac{1}{2}u^2} + e^{-\frac{1}{2}u^2} \right) du \\
&= \frac{1}{2q\sqrt{\pi} (z_a^q)^{1-\frac{1}{q}}} e^{-\frac{1}{4}(z_a^q)^{2/q}} (1+1) \\
&= \frac{1}{q\sqrt{\pi}} (z_a^q)^{\frac{1}{q}-1} e^{-\frac{1}{4}(z_a^q)^{2/q}} \\
&= \frac{\frac{2}{q}}{(2^q)^{1/q} \Gamma\left(\frac{1}{2} + \frac{1}{q}\right)} (z_a^q)^{\frac{1}{q}-1} e^{-\left(\frac{z_a^q}{2^q}\right)^{2/q}}.
\end{aligned} \tag{22}$$

The density function given previously (Eq. 22) is a Generalized Gamma density with parameters  $b = \frac{2}{q}$ ,  $c = 2^q$ , and  $d = \frac{1}{q}$ . This distribution has mean and variance given by

$$\begin{aligned}
\mu_{z_a^q} &= \frac{c\Gamma\left(\frac{d+1}{b}\right)}{\Gamma\left(\frac{d}{b}\right)} \\
&= \frac{2^q\Gamma\left(\frac{q+1}{2}\right)}{\sqrt{\pi}}
\end{aligned} \tag{23}$$

and

$$\begin{aligned}
\sigma_{z_a^q}^2 &= c^2 \left[ \frac{\Gamma\left(\frac{d+2}{b}\right)}{\Gamma\left(\frac{d}{b}\right)} - \left( \frac{\Gamma\left(\frac{d+1}{b}\right)}{\Gamma\left(\frac{d}{b}\right)} \right)^2 \right] \\
&= 4^q \left[ \frac{\Gamma\left(q + \frac{1}{2}\right)}{\sqrt{\pi}} - \frac{\Gamma^2\left(\frac{1}{2}q + \frac{1}{2}\right)}{\pi} \right].
\end{aligned} \tag{24}$$

By linearity of the expected value and variance operators under the iid assumption, the mean (Eq. 23) and variance (Eq. 24) of the random variable  $Z_a^q$  allow the  $p$ -dimensional mean and variance of the  $\left(D_{ij}^{(q)}\right)^q$  distribution to be computed directly as

$$\begin{aligned}
\mu_{\left(D_{ij}^{(q)}\right)^q} &= \mathbb{E} \left[ \left( D_{ij}^{(q)} \right)^q \right] = \mathbb{E} \left( \sum_{a \in \mathcal{A}} Z_a^q \right) = \sum_{a \in \mathcal{A}} \mathbb{E} (Z_a^q) = \sum_{a \in \mathcal{A}} \frac{2^q \Gamma\left(\frac{q+1}{2}\right)}{\sqrt{\pi}} \\
&= \frac{2^q \Gamma\left(\frac{q+1}{2}\right)}{\sqrt{\pi}} p
\end{aligned} \tag{25}$$

<sup>1</sup>and

$$\begin{aligned}
 \sigma^2_{(D_{ij}^{(q)})^q} &= \text{Var} \left[ \left( D_{ij}^{(q)} \right)^q \right] = \text{Var} \left( \sum_{a \in \mathcal{A}} Z_a^q \right) \\
 &= \sum_{a \in \mathcal{A}} \text{Var} (Z_a^q) \\
 &= \sum_{a \in \mathcal{A}} 4^q \left[ \frac{\Gamma(q + \frac{1}{2})}{\sqrt{\pi}} - \frac{\Gamma^2(\frac{1}{2}q + \frac{1}{2})}{\pi} \right] \\
 &= 4^q \left[ \frac{\Gamma(q + \frac{1}{2})}{\sqrt{\pi}} - \frac{\Gamma^2(\frac{1}{2}q + \frac{1}{2})}{\pi} \right] p.
 \end{aligned} \tag{26}$$

<sup>10</sup> Therefore, the asymptotic distribution of  $D_{ij}^{(q)}$  for standard normal data is

$$\begin{aligned}
 \mathcal{N} \left( \left( 2^q \frac{\Gamma(\frac{q+1}{2})}{\sqrt{\pi}} p \right)^{1/q}, \right. \\
 \left. \frac{4^q p}{q^2 \left( \frac{2^q \Gamma(\frac{1}{2}q + \frac{1}{2})}{\sqrt{\pi}} p \right)^{2(1-\frac{1}{q})}} \left[ \frac{\Gamma(q + \frac{1}{2})}{\sqrt{\pi}} - \frac{\Gamma^2(\frac{1}{2}q + \frac{1}{2})}{\pi} \right] \right).
 \end{aligned} \tag{27}$$

<sup>19</sup> We provide a summary table of the moment estimates (Eq. 27) for the  $L_q$  metric on standard normal data (Table 1). The summary is organized by data type, type of statistic (mean or variance), and corresponding asymptotic formula. In the next section, we derive the  $L_q$  distance distribution on standard uniform data in a similar fashion.

### <sup>24</sup>3.1.2 Standard uniform data

<sup>25</sup>If  $X_{ia}, X_{ja} \stackrel{iid}{\sim} \mathcal{U}(0, 1)$ , then the marginal density functions with respect to  $X$  for  $X_{ia}, X_{ia} - (Z_a^q)^{1/q}$ , and  $X_{ia} + (Z_a^q)^{1/q}$  are defined as

$$f_X(x_{ia}) = 1, \quad 0 \leq x_{ia} \leq 1 \tag{28}$$

$$f_X \left( x_{ia} - (z_a^q)^{1/q} \right) = 1, \quad 0 \leq x_{ia} - (z_a^q)^{1/q} \leq 1, \text{ and} \tag{29}$$

$$f_X \left( x_{ia} + (z_a^q)^{1/q} \right) = 1, \quad 0 \leq x_{ia} + (z_a^q)^{1/q} \leq 1. \quad (30)$$

Substituting these marginal densities (Eqs. 28-30) into the more general density function for  $Z_a^q$  (Eq. 14), we have

$$\begin{aligned} f_{Z_a^q}(z_a^q) &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_{-\infty}^{\infty} \left\{ f_X(x_{ia}) \left[ f_X \left( x_{ia} - (z_a^q)^{1/q} \right) \right. \right. \\ &\quad \left. \left. + f_X \left( x_{ia} + (z_a^q)^{1/q} \right) \right] \right\} dx_{ia} \\ &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_0^1 \left[ f_X(x_{ia} - (z_a^q)^{1/q}) + f_X \left( x_{ia} + (z_a^q)^{1/q} \right) \right] dx_{ia} \\ &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} \int_{(z_a^q)}^1 1 dx_{ia} + \int_0^{1-(z_a^q)^{1/q}} 1 dx_{ia} \\ &= \frac{1}{q(z_a^q)^{1-\frac{1}{q}}} [(1 - (z_a^q)^{1/q}) + (1 - (z_a^q)^{1/q})] \\ &= \frac{1}{q} \cdot 2 (z_a^q)^{\frac{1}{q}-1} [1 - (z_a^q)^{1/q}]^{2-1}, \end{aligned} \quad (31)$$

where  $0 < z_a \leq 1$ .

The previous density (Eq. 31) is a Kumaraswamy density with parameters  $b = \frac{1}{q}$  and  $c = 2$  with moment generating function (MGF) given by

$$\begin{aligned} M_n &= \frac{c \Gamma \left( 1 + \frac{n}{b} \right) \Gamma(c)}{\Gamma \left( 1 + c + \frac{n}{b} \right)} \\ &= \frac{2}{(nq + 2)(nq + 1)}. \end{aligned} \quad (32)$$

Using this MGF (Eq. 32), the mean and variance of  $Z_a^q$  are computed as

$$\mu_{z_a^q} = \frac{2}{(q + 2)(q + 1)} \quad (33)$$

and

$$\sigma_{z_a^q}^2 = \frac{1}{(q + 1)(2q + 1)} - \left( \frac{2}{(q + 2)(q + 1)} \right)^2. \quad (34)$$

By linearity of the expected value and variance operators under the iid assumption, the mean (Eq. 33) and variance (Eq. 34) of the random variable  $Z_a^q$  allow the  $p$ -dimensional mean and variance of the  $\left( D_{ij}^{(q)} \right)^q$  distribution to be computed

directly as

$$\begin{aligned}
 \mu_{(D_{ij}^{(q)})^q} &= \mathbb{E} \left[ (D_{ij}^{(q)})^q \right] = \mathbb{E} \left( \sum_{a \in \mathcal{A}} Z_a^q \right) \\
 &= \sum_{a \in \mathcal{A}} \mathbb{E}(Z_a^q) \\
 &= \sum_{a \in \mathcal{A}} \frac{2}{(q+2)(q+1)} \\
 &= \frac{2p}{(q+2)(q+1)}
 \end{aligned} \tag{35}$$

and

$$\begin{aligned}
 \sigma_{(D_{ij}^{(q)})^q}^2 &= \text{Var} \left[ (D_{ij}^{(q)})^q \right] = \text{Var} \left( \sum_{a \in \mathcal{A}} Z_a^q \right) \\
 &= \sum_{a \in \mathcal{A}} \text{Var}(Z_a^q) \\
 &= \sum_{a \in \mathcal{A}} \left[ \frac{1}{(q+1)(2q+1)} - \left( \frac{2}{(q+2)(q+1)} \right)^2 \right] \\
 &= \left[ \frac{1}{(q+1)(2q+1)} - \left( \frac{2}{(q+2)(q+1)} \right)^2 \right] p.
 \end{aligned} \tag{36}$$

Therefore, the asymptotic distribution of  $D_{ij}^{(q)}$  for standard uniform data is

$$\begin{aligned}
 \mathcal{N} \left( \left( \frac{2p}{(q+2)(q+1)} \right)^{1/q}, \right. \\
 \left. \frac{p}{q^2 \left( \frac{2p}{(q+2)(q+1)} \right)^{2(1-\frac{1}{q})}} \left[ \frac{1}{(q+1)(2q+1)} - \left( \frac{2}{(q+2)(q+1)} \right)^2 \right] \right).
 \end{aligned} \tag{37}$$

We provide a summary table of the moment estimates (Eq. 37) for the  $L_q$  metric on standard uniform data (Table 1). The summary is organized by data type, type of statistic (mean or variance), and corresponding asymptotic formula. In the next section, we use our general  $L_q$  distance distribution derivations to provide Manhattan ( $q = 1$ ) and Euclidean ( $q = 2$ ) asymptotic moments on both standard normal and standard uniform data. These are the most commonly applied metrics in the context of nearest-neighbor feature selection, so they are of particular interest.

### <sup>1</sup>3.2 Manhattan ( $L_1$ ) <sup>1</sup>

<sup>2</sup>With our general formulas for the asymptotic <sup>2</sup>

<sup>3</sup>mean and variance (Eqs. 27 and 37) for any value of  $q \in \mathbb{N}$ , we can simply substitute<sup>3</sup>

<sup>4</sup>a particular value of  $q$  in order to determine the asymptotic distribution of the<sup>4</sup>

<sup>5</sup>corresponding distance  $L_q$  metric. We demonstrate this with the example of the<sup>5</sup>

<sup>6</sup>Manhattan metric ( $L_1$ ) for standard normal and standard uniform data (Eq. 1,<sup>6</sup>

<sup>7</sup> $q = 1$ ). <sup>7</sup>

<sup>8</sup> <sup>8</sup>

#### <sup>9</sup>3.2.1 Standard normal data <sup>9</sup>

<sup>10</sup>Substituting  $q = 1$  into the asymptotic formula for the mean  $L_q$  distance (Eq. 27),<sup>10</sup>

<sup>11</sup>we have the following for expected  $L_1$  distance between two independently sampled<sup>11</sup>

<sup>12</sup>instances  $i, j \in \mathcal{I}$  in standard normal data <sup>12</sup>

$$\begin{aligned} \text{E} \left( D_{ij}^{(1)} \right) &= \left( 2 \frac{\Gamma \left( \frac{1+1}{2} \right)}{\sqrt{\pi}} p \right)^{1/1} \\ &= \frac{2p}{\sqrt{\pi}}. \end{aligned} \quad (38) \quad \begin{matrix} \text{13} \\ \text{14} \\ \text{15} \\ \text{16} \end{matrix}$$

<sup>17</sup> We see in the formula for the expected Manhattan distance (Eq. 38) that  $D_{ij}^{(1)} \sim p^{17}$

<sup>18</sup>in the limit, which implies that this distance is unbounded as attribute dimension<sup>18</sup>

<sup>19</sup> $p$  increases. <sup>19</sup>

<sup>20</sup> Substituting  $q = 1$  into the formula for the asymptotic variance of  $D_{ij}^{(1)}$  (Eq. 27)<sup>20</sup>

<sup>21</sup>leads to the following <sup>21</sup>

$$\begin{aligned} \text{Var} \left( D_{ij}^{(1)} \right) &= \frac{4^1 p}{1^2 \left( \frac{2^1 \Gamma \left( \frac{1}{2}(1) + \frac{1}{2} \right)}{\sqrt{\pi}} p \right)^{2(1 - \frac{1}{1})}} \left[ \frac{\Gamma \left( 1 + \frac{1}{2} \right)}{\sqrt{\pi}} - \frac{\Gamma^2 \left( \frac{1}{2}(1) + \frac{1}{2} \right)}{\pi} \right] \\ &= \frac{2(\pi - 2)p}{\pi}. \end{aligned} \quad (39) \quad \begin{matrix} \text{22} \\ \text{23} \\ \text{24} \\ \text{25} \\ \text{26} \end{matrix}$$

<sup>27</sup> Similar to the mean (Eq. 38), the limiting variance of  $D_{ij}^{(1)}$  (Eq. 39) grows on the<sup>27</sup>

<sup>28</sup>order of attribute dimension  $p$ , which implies that points become more dispersed<sup>28</sup>

<sup>29</sup>as the dimension increases. The moment estimates given in this section (Eqs. 38,<sup>29</sup>

<sup>30</sup>and 39) are summarized in a table that is organized by metric, data type, statistic<sup>30</sup>

<sup>31</sup>(mean or variance), and asymptotic formula (Table 2). <sup>31</sup>

<sup>32</sup> <sup>32</sup>

<sup>33</sup> <sup>33</sup>



### 3.2.2 Standard uniform data

Substituting  $q = 1$  into the asymptotic formula of the mean (Eq. 37), we have the

following for the expected  $L_1$  distance between two independently sampled instances

$i, j \in \mathcal{I}$  in standard uniform data

$$\begin{aligned} \mathbb{E}\left(D_{ij}^{(1)}\right) &= \left(\frac{2p}{(1+2)(1+1)}\right)^{1/1} \\ &= \frac{p}{3}. \end{aligned} \quad (40)$$

Once again, we see that the mean of  $D_{ij}^{(1)}$  (Eq. 40) grows on the order of  $p$  just as in the case of standard normal data.

Substituting  $q = 1$  into the formula of the asymptotic variance of  $D_{ij}^{(1)}$  (Eq. 37) leads to the following

$$\begin{aligned} \text{Var}\left(D_{ij}^{(1)}\right) &= \frac{p}{1^2 \left(\frac{2p}{(1+2)(1+1)}\right)^{2(1-\frac{1}{1})}} \left\{ \left[ \frac{1}{(1+1)(2(1)+1)} \right. \right. \\ &\quad \left. \left. - \left( \frac{2}{(1+2)(1+1)} \right)^2 \right] \right\} \\ &= \frac{p}{18}. \end{aligned} \quad (41)$$

As in the case of the  $L_1$  metric on standard normal data, we have a variance (Eq. 41) that grows on the order of  $p$ . The distances between points in high-dimensional uniform data become more widely dispersed with this metric. The moment estimates given in this section (Eqs. 40 and 41) are summarized in a table that is organized by metric, data type, statistic (mean or variance), and asymptotic formula (Table 2).

### 3.2.3 Distribution of one-dimensional projected pairwise distance onto an attribute

In nearest-neighbor distance-based feature selection like NPDR and Relief-based algorithms, the one-dimensional projection of the pairwise distance onto an attribute (Eq. 2) is particularly fundamental to attribute quality for association with an outcome. For instance, this distance projection is the predictor used to determine beta coefficients in NPDR. In particular, understanding distributional properties of the projected distances is necessary for defining pseudo P values for NPDR. In this section, we summarize the exact distribution of the one-dimensional projected dis-

ance onto an attribute  $a \in \mathcal{A}$ . These results apply to continuous data, such as gene<sup>1</sup>  
expression.<sup>2</sup>

<sup>3</sup> In previous sections, we derived the exact density function (Eq. 14) and moments<sup>3</sup>  
<sup>4</sup>(Eqs. 15 and 16) for the distribution of  $Z_a^q = |X_{ia} - X_{ja}|^q$ . We then derived the<sup>4</sup>  
<sup>5</sup>exact density (Eq. 22) and moments (Eqs. 23 and 24) for standard normal data.<sup>5</sup>  
<sup>6</sup>Analogously, we formulated the exact density (Eq. 31) and moments (Eqs. 33 and<sup>6</sup>  
<sup>7</sup>34) for standard uniform data. From these exact densities and moments, we simply<sup>7</sup>  
<sup>8</sup>substitute  $q = 1$  to define the distribution of the one-dimensional projected distance<sup>8</sup>  
onto an attribute  $a \in \mathcal{A}$ .<sup>9</sup>

<sup>10</sup> Assuming data is standard normal, we substitute  $q = 1$  into the density function<sup>10</sup>  
<sup>11</sup>of  $Z_a^q$  (Eq. 22) to arrive at the following<sup>11</sup>

$$\begin{aligned} f_{Z_a^1}(z_a^1) &= \frac{\frac{2}{1}}{(2^1)^{1/1} \Gamma\left(\frac{1}{2}\right)} (z_a^1)^{1/1-1} e^{-\left(\frac{z_a^1}{2^1}\right)^{2/1}}, \quad z_a > 0 \\ &= \frac{1}{\sqrt{\pi}} z_a e^{-\frac{1}{4} z_a^2}, \quad z_a > 0. \end{aligned} \quad (42)$$

<sup>16</sup> The mean corresponding to this Generalized Gamma density is computed by<sup>16</sup>  
<sup>17</sup>substituting  $q = 1$  into the formula for the mean of  $Z_a^q$  (Eq. 23). This result is given<sup>17</sup>  
by<sup>18</sup>

$$\begin{aligned} \mu_{Z_a^1} &= \frac{2^1 \Gamma\left(\frac{1+1}{2}\right)}{\sqrt{\pi}} \\ &= \frac{2}{\sqrt{\pi}}. \end{aligned} \quad (43)$$

<sup>23</sup> Substituting  $q = 1$  into Eq. 24 for the variance, we have the following<sup>23</sup>

$$\begin{aligned} \sigma_{Z_a^1}^2 &= 4^1 \left[ \frac{\Gamma\left(1 + \frac{1}{2}\right)}{\sqrt{\pi}} - \frac{\Gamma^2\left(\frac{1}{2} \cdot 1 + \frac{1}{2}\right)}{\pi} \right] \\ &= \frac{2(\pi - 2)}{\pi}. \end{aligned} \quad (44)$$

<sup>28</sup> These last few results (Eqs. 42-44) provide us with the distribution for NPDR<sup>28</sup>  
predictors when the data is from the standard normal distribution.<sup>29</sup>

<sup>30</sup> If we have standard uniform data, we substitute  $q = 1$  into the density function<sup>30</sup>  
of  $Z_a^q$  (Eq. 31) to obtain the following<sup>31</sup>

$$\begin{aligned} f_{Z_a^1} &= \frac{1}{1} \cdot 2 (z_a^1)^{1/1-1} \left[ 1 - (z_a^1)^{1/1} \right]^{2-1}, \quad 0 < z_a \leq 1 \\ &= 2z_a(1 - z_a), \quad 0 < z_a \leq 1. \end{aligned} \quad (45)$$

<sup>1</sup> The mean corresponding to this Kumaraswamy density is computed by substitut-<sup>1</sup>  
<sup>2</sup>ing  $q = 1$  into the formula for the mean of  $Z_a^q$  (Eq. 33). After substitution, we have<sup>2</sup>  
<sup>3</sup>the following result<sup>3</sup>

$$\begin{aligned} \mu_{Z_a^1} &= \frac{2}{(1+2)(1+1)} \\ &= \frac{1}{3}. \end{aligned} \quad (46)$$

<sup>4</sup> Substituting  $q = 1$  into the formula for the variance of  $Z_a^q$  (Eq. 34), we have the<sup>4</sup>  
<sup>5</sup>following<sup>5</sup>

$$\begin{aligned} \sigma_{Z_a^1}^2 &= \frac{1}{(1+1)(2 \cdot 1 + 1)} - \left( \frac{2}{(1+2)(1+1)} \right)^2 \\ &= \frac{1}{18}. \end{aligned} \quad (47)$$

<sup>6</sup> In the event that the data distribution is standard uniform, the density function<sup>6</sup>  
<sup>7</sup>(Eq. 45), the mean (Eq. 46), and the variance (Eq. 47) sufficiently define the distri-<sup>7</sup>  
<sup>8</sup>bution for NPDR predictors. The means (Eqs. 43 and 46) and variances (Eqs. 44<sup>8</sup>  
<sup>9</sup>and 47) come from the exact distribution of pairwise distances with respect to a<sup>9</sup>  
<sup>10</sup>single attribute  $a \in \mathcal{A}$ . This is the distribution of the so-called “projection” of the<sup>10</sup>  
<sup>11</sup>pairwise distance onto a single attribute to which we have been referring, which is<sup>11</sup>  
<sup>12</sup>a direct implication from our more general derivations. In a similar manner, one<sup>12</sup>  
<sup>13</sup>can substitute any value of  $q \geq 2$  into the general densities of  $Z_a^q$  for standard<sup>13</sup>  
<sup>14</sup>normal (Eq. 22) and standard uniform (Eq. 31) to derive the associated density of<sup>14</sup>  
<sup>15</sup> $Z_a^q = |X_{ia} - X_{ja}|^q$  for the given data type.<sup>15</sup>

### <sup>16</sup>3.3 Euclidean ( $L_2$ )<sup>16</sup>

<sup>17</sup>Moment estimates for the Euclidean metric are obtained by substituting  $q = 2$  into<sup>17</sup>  
<sup>18</sup>the asymptotic moment formulas for standard normal data (Eq. 27) and standard<sup>18</sup>  
<sup>19</sup>uniform data (Eq. 37). As in the case of the Manhattan metric in the previous<sup>19</sup>  
<sup>20</sup>sections, we initially proceed by deriving Euclidean distance moments in standard<sup>20</sup>  
<sup>21</sup>normal data.<sup>21</sup>

#### <sup>22</sup>3.3.1 Standard normal data<sup>22</sup>

<sup>23</sup>Substituting  $q = 2$  into the asymptotic formula of the mean (Eq. 27), we have the<sup>23</sup>  
<sup>24</sup>following for expected  $L_2$  distance between two independently sampled instances<sup>24</sup>  
<sup>25</sup><sup>25</sup>

<sup>1</sup> $i, j \in \mathcal{I}$  in standard normal data

$$\begin{aligned} \text{E} \left( D_{ij}^{(2)} \right) &= \left( 2 \frac{\Gamma \left( \frac{2+1}{2} \right)}{\sqrt{\pi}} p \right)^{1/2} \\ &= \sqrt{2p}. \end{aligned} \quad (48)$$

<sup>6</sup> In the case of  $L_2$  on standard normal data, we see that the mean of  $D_{ij}^{(2)}$  (Eq. 48)<sup>6</sup>

<sup>7</sup>grows on the order of  $\sqrt{p}$ . Hence, the Euclidean distance does not increase as quickly<sup>7</sup>

<sup>8</sup>as the Manhattan distance on standard normal data.

<sup>9</sup> Substituting  $q = 2$  into the formula for the asymptotic variance of  $D_{ij}^{(2)}$  (Eq. 27)<sup>9</sup>  
<sup>10</sup>leads to the following

$$\begin{aligned} \text{Var} \left( D_{ij}^{(2)} \right) &= \frac{4^2 p}{2^2 \left( \frac{2^2 \Gamma \left( \frac{1}{2}(2) + \frac{1}{2} \right)}{\sqrt{\pi}} p \right)^{2(1-\frac{1}{2})}} \left[ \frac{\Gamma \left( 2 + \frac{1}{2} \right)}{\sqrt{\pi}} - \frac{\Gamma^2 \left( \frac{1}{2}(2) + \frac{1}{2} \right)}{\pi} \right] \\ &= 1. \end{aligned} \quad (49)$$

Surprisingly, the asymptotic variance (Eq. 49) is just 1. Regardless of data dimensions  $m$  and  $p$ , the variance of Euclidean distances on standard normal data tends to 1. Therefore, most instances are contained within a ball of radius 1 about the mean in high attribute dimension  $p$ . This means that the Euclidean distance distribution on standard normal data is simply a horizontal shift to the right of the standard normal distribution.

For the case in which the number of attributes  $p$  is small, we have an improved estimate of the mean (Eq. 6). The lower dimensional estimate of the mean is given by

$$\begin{aligned} \text{E} \left( D_{ij}^{(2)} \right) &= \left( 2 \frac{\Gamma \left( \frac{2+1}{2} \right)}{\sqrt{\pi}} p - 1 \right)^{1/2} \\ &= \sqrt{2p - 1}. \end{aligned} \quad (50)$$

For high dimensional data sets like gene expression [16, 17], which typically contain thousands of genes (or attributes), it is clear that the magnitude of  $p$  will be sufficient to use the standard asymptotic estimate (Eq. 48) since  $\sqrt{2p} \approx \sqrt{2p - 1}$  in that case. The moment estimates given in this section (Eqs. 50 and 49) are summarized in a table that is organized by metric, data type, statistic (mean or variance), and asymptotic formula (Table 2).

### 3.3.2 Standard uniform data

Substituting  $q = 2$  into the asymptotic formula of the mean (Eq. 37), we have the

following for expected  $L_2$  distance between two independently sampled instances

$i, j \in \mathcal{I}$  in standard uniform data

$$\begin{aligned} \mathbb{E} \left( D_{ij}^{(2)} \right) &= \left( \frac{2p}{(2+2)(2+1)} \right)^{1/2} \\ &= \sqrt{\frac{p}{6}}. \end{aligned} \quad (51)$$

As in the case of standard normal data, the expected value of  $D_{ij}^{(2)}$  (Eq. 51) grows on the order of  $\sqrt{p}$ .

Substituting  $q = 2$  into the formula for the asymptotic variance of  $D_{ij}^{(2)}$  (Eq. 37) leads to the following

$$\begin{aligned} \text{Var} \left( D_{ij}^{(2)} \right) &= \frac{p}{2^2 \left( \frac{2p}{(2+2)(2+1)} \right)^{2(1-\frac{1}{2})}} \left\{ \left[ \frac{1}{(2+1)(2(2)+1)} \right. \right. \\ &\quad \left. \left. - \left( \frac{2}{(2+2)(2+1)} \right)^2 \right] \right\} \\ &= \frac{7}{120}. \end{aligned} \quad (52)$$

Once again, the variance of Euclidean distance surprisingly approaches a constant.

For the case in which the number of attributes  $p$  is small, we have an improved estimate of the mean (Eq. 6). The lower dimensional estimate of the mean is given by

$$\begin{aligned} \mathbb{E} \left( D_{ij}^{(2)} \right) &= \left( \frac{2p}{(2+2)(2+1)} - \frac{7}{120} \right)^{1/2} \\ &= \sqrt{\frac{p}{6} - \frac{7}{120}}. \end{aligned} \quad (53)$$

We summarize the moment estimates given in this section for standard  $L_q$  metrics (Eqs. 53 and 52) organized by metric, data type, statistic (mean or variance), and asymptotic formula (Table 2). In the next section, we extend these results for the standard  $L_q$  metric to derive asymptotics for the attribute range-normalized (max-min)  $L_q$  metric used frequently in Relief-based algorithms [1, 3] for scoring attributes. These derivations use extreme value theory to handle the maximum and minimum attributes for standard normal and standard uniform data.

### 3.4 Distributions of attribute range-normalized $L_q$ metric

For Relief-based methods [1, 3], the standard numeric diff metric is normalized by

the range of each attribute  $a$ :

$$d_{ij}^{\text{num}}(a) = \text{diff}(a, (i, j)) = \frac{|X_{ia} - X_{ja}|}{\max(a) - \min(a)}, \quad (54)$$

where  $X$  is the subject attribute numeric data matrix,  $\max(a) = \max_{k \in \mathcal{I}} \{X_{ka}\}$ ,  $\min(a) = \min_{k \in \mathcal{I}} \{X_{ka}\}$ , and  $\mathcal{I} = \{1, 2, \dots, m\}$ .

The pairwise distance using this max-min range-normalized diff metric is then computed as

$$\begin{aligned} D_{ij}^{(q*)} &= \left( \sum_{a \in \mathcal{A}} |d_{ij}(a)|^q \right)^{1/q} \\ &= \left( \sum_{a \in \mathcal{A}} \left( \frac{|X_{ia} - X_{ja}|}{\max(a) - \min(a)} \right)^q \right)^{1/q}. \end{aligned} \quad (55)$$

In order to determine moments of asymptotic max-min normalized distance (Eq. 54) distributions, we will first derive the asymptotic extreme value distributions of the attribute maximum and minimum. Although the exact distribution of the maximum or minimum requires an assumption about the data distribution, the Fisher-Tippett-Gnedenko Theorem is an important result that allows one to generally categorize the extreme value distribution for a collection of independent and identically distributed random variables into one of three distributional families. This theorem does not, however, tell us the exact distribution of the maximum that we require in order to determine asymptotic results for the max-min normalized distance (Eq. 55). We mention this theorem simply to provide some background on convergence of extreme values. Before stating the theorem, we first need the following definition

**Definition 3.1** A distribution  $\mathcal{F}_X$  is said to be **degenerate** if its density function  $f_X$  is the Dirac delta  $\delta(x - c_0)$  centered at a constant  $c_0 \in \mathbb{R}$ , with corresponding distribution function  $F_X$  defined as

$$F_X(x) = \begin{cases} 1, & x \geq c_0, \\ 0, & x < c_0. \end{cases}$$

<sup>1</sup>**Theorem 3.2** (Fisher-Tippett-Gnedenko) *Let  $X_{ia} \stackrel{iid}{\sim} \mathcal{F}_X(\mu_x, \sigma_x^2)$  for all  $i \in \mathcal{I}^1$*

<sup>2</sup>*and let  $X_a^{max} = \max_{k \in \mathcal{I}} \{X_{ka}\}$ . If there exists two non-random sequences  $b_m > 0$  and<sup>2</sup>*

<sup>3</sup> $c_m$  such that

$$\lim_{m \rightarrow \infty} P\left(\frac{X_a^{max} - c_m}{b_m} \leq x\right) = G_X(x),$$

<sup>6</sup>where  $G_X$  is a non-degenerate distribution function, then the limiting distribution<sup>7</sup>

<sup>8</sup> $G_X$  is in the Gumbel, Fréchet, or Weibull family.

<sup>9</sup> The three distribution families given in Theorem 3.2 are actually special cases<sup>9</sup>  
<sup>10</sup>of the Generalized Extreme Value Distribution. In the context of extreme values,<sup>10</sup>  
<sup>11</sup>Theorem 3.2 is analogous to the Central Limit Theorem for the distribution of<sup>11</sup>  
<sup>12</sup>sample mean. Although we will not explicitly invoke this theorem, it does tell us<sup>12</sup>  
<sup>13</sup>something very important about the asymptotic behavior of sample extremes under<sup>13</sup>  
<sup>14</sup>certain necessary conditions. For illustration of this general phenomenon of sample<sup>14</sup>  
<sup>15</sup>extremes, we derive the distribution of the maximum for standard normal data to<sup>15</sup>  
<sup>16</sup>show that the limiting distribution is in the Gumbel family, which is a well known<sup>16</sup>  
<sup>17</sup>result. In the case of standard uniform data, we will derive the distribution of the<sup>17</sup>  
<sup>18</sup>maximum and minimum directly. Regardless of data type, the distribution of the<sup>18</sup>  
<sup>19</sup>sample maximum can be derived as follows

$$\begin{aligned} P[X_a^{max} \leq x] &= P\left[\max_{k \in \mathcal{I}} \{X_{ka}\} \leq x\right] \\ &= P[X_{1a} \leq x, X_{2a} \leq x, \dots, X_{ma} \leq x] \\ &= \prod_{k=1}^m P[X_{ka} \leq x] \\ &= \prod_{k=1}^m F_X(x) \\ &= [F_X(x)]^m. \end{aligned} \tag{56}$$

<sup>28</sup> Using more precise notation, the distribution function of the sample maximum in<sup>28</sup>  
<sup>29</sup>standard normal data is

$$F_{\max}(x) = [F_X(x)]^m, \tag{57}$$

<sup>32</sup>where  $m$  is the size of the sample from which the maximum is derived and  $F_X$  is<sup>32</sup>  
<sup>33</sup>the distribution function corresponding to the data sample. This means that the<sup>33</sup>

<sup>1</sup>distribution of the sample maximum relies only on the distribution function of the <sup>1</sup>  
<sup>2</sup>data from which extremes are drawn  $F_X$  and the size of the sample  $m$ . <sup>2</sup>

<sup>3</sup> Differentiating the distribution function (Eq. 57) gives us the following density <sup>3</sup>  
<sup>4</sup>function for the distribution of the maximum <sup>4</sup>

$$\begin{aligned}
 f_{\max}(x) &= \frac{d}{dx} F_{\max}(x) \\
 &= \frac{d}{dx} [F_X(x)]^m \\
 &= m[F_X(x)]^{m-1} f_X(x),
 \end{aligned}
 \tag{58}$$

<sup>9</sup> where  $m$  is the size of the sample from which the maximum is derived,  $F_X$  is <sup>9</sup>  
<sup>10</sup>the distribution function corresponding to the data sample, and  $f_X$  is the density <sup>10</sup>  
<sup>11</sup>function corresponding to the data sample. Similar to the distribution function for <sup>11</sup>  
<sup>12</sup>the sample maximum (Eq. 57), the density function (Eq 58) relies only on the <sup>12</sup>  
<sup>13</sup>distribution and density function of the data from which extremes are derived. <sup>13</sup>

<sup>14</sup> The distribution of the sample minimum,  $X_a^{\min}$ , can be derived as follows <sup>14</sup>

$$\begin{aligned}
 P[X_a^{\min} \leq x] &= 1 - P[X_a^{\min} \geq x] \\
 &= 1 - P\left[\min_{k \in \mathcal{I}} \{X_{ka}\} \geq x\right] \\
 &= 1 - P[X_{1a} \geq x, X_{2a} \geq x, \dots, X_{ma} \geq x] \\
 &= 1 - \prod_{k=1}^m P[X_{ka} \geq x] \\
 &= 1 - [P[X_{1a} \geq x]]^m \\
 &= 1 - [1 - P[X_{1a} \leq x]]^m \\
 &= 1 - [1 - F_X(x)]^m,
 \end{aligned}
 \tag{59}$$

<sup>25</sup> where  $m$  is the size of the sample from which the maximum is derived and  $F_X$  is the <sup>25</sup>  
<sup>26</sup>distribution function corresponding to the data sample. Therefore, the distribution <sup>26</sup>  
<sup>27</sup>of sample minimum also relies only on the distribution function of the data from <sup>27</sup>  
<sup>28</sup>which extremes are derived. <sup>28</sup>

<sup>29</sup> With more precise notation, we have the following expression for the distribution <sup>29</sup>  
<sup>30</sup>function of the minimum <sup>30</sup>

$$F_{\min}(x) = 1 - [1 - F_X(x)]^m. \tag{60}$$



<sup>1</sup>where  $m$  is the size of the sample from which the minimum is derived and  $F_X$  is <sup>1</sup>  
<sup>2</sup>the distribution function corresponding to the data sample. <sup>2</sup>

<sup>3</sup> Differentiating the distribution function (Eq. 60) gives us the following density <sup>3</sup>  
<sup>4</sup>function for the distribution of sample minimum <sup>4</sup>

$$\begin{aligned}
 f_{\min}(x) &= \frac{d}{dx} F_{\min}(x) \\
 &= \frac{d}{dx} (1 - [1 - F_X(x)]^m) \\
 &= m [1 - F_X(x)]^{m-1} f_X(x),
 \end{aligned}
 \tag{61}$$

where  $m$  is the size of the sample from which the minimum is derived,  $F_X$  is the <sup>9</sup>  
<sup>10</sup>distribution function corresponding to the data sample, and  $f_X$  is the density func- <sup>10</sup>  
<sup>11</sup>tion corresponding to the data sample. As in the case of the density function for <sup>11</sup>  
<sup>12</sup>sample maximum (Eq. 58), the density function for sample minimum relies only on <sup>12</sup>  
<sup>13</sup>the distribution  $F_X$  and density  $f_X$  functions of the data from which extremes are <sup>13</sup>  
<sup>14</sup>derived and the sample size  $m$ . <sup>14</sup>

Given the densities of the distribution of sample maximum and minimum, we can <sup>15</sup>  
<sup>16</sup>easily compute the raw moments and variance. The first moment about the origin <sup>16</sup>  
<sup>17</sup>of the distribution of sample maximum is given by the following <sup>17</sup>

$$\begin{aligned}
 \mu_{\max}^{(1)}(m) &= E(X_a^{\max}) = \int_{-\infty}^{\infty} x f_{\max}(x) dx \\
 &= \int_{-\infty}^{\infty} x (m [F_X(x)]^{m-1} f_X(x)) dx \\
 &= m \int_{-\infty}^{\infty} x f_X(x) [F_X(x)]^{m-1} dx,
 \end{aligned}
 \tag{62}$$

where  $m$  is the sample size,  $F_X$  is the distribution function, and  $f_X$  is the density <sup>23</sup>  
<sup>24</sup>function of the data from which the maximum is derived. <sup>24</sup>

The second raw moment of the distribution of sample maximum is derived simi- <sup>25</sup>  
<sup>26</sup>larly as follows <sup>26</sup>

$$\begin{aligned}
 \mu_{\max}^{(2)}(m) &= E[(X_a^{\max})^2] = \int_{-\infty}^{\infty} x^2 f_{\max}(x) dx \\
 &= \int_{-\infty}^{\infty} x^2 (m [F_X(x)]^{m-1} f_X(x)) dx \\
 &= m \int_{-\infty}^{\infty} x^2 f_X(x) [F_X(x)]^{m-1} dx
 \end{aligned}
 \tag{63}$$

where  $m$  is the sample size,  $F_X$  is the distribution function, and  $f_X$  is the density <sup>32</sup>  
<sup>33</sup>function of the data from which the maximum is derived. <sup>33</sup>

<sup>1</sup> Using the first (Eq. 62) and second (Eq. 63) raw moments of the distribution of<sup>1</sup>  
<sup>2</sup>sample maximum, the variance is given by<sup>2</sup>

$$\sigma_{\max}^2(m) = \mu_{\max}^{(2)}(m) - \left[ \mu_{\max}^{(1)}(m) \right]^2, \quad (64)$$

<sup>5</sup> where  $m$  is the sample size of the data from which the maximum is derived and<sup>5</sup>  
<sup>6</sup> $\mu_{\max}^{(1)}(m)$  and  $\mu_{\max}^{(2)}$  are the first and second raw moments, respectively, of the dis-<sup>6</sup>  
<sup>7</sup>tribution of sample maximum.<sup>7</sup>

<sup>8</sup> Moving on to the distribution of sample minimum, the first raw moment is given<sup>8</sup>  
<sup>9</sup>by the following<sup>9</sup>

$$\begin{aligned} \mu_{\min}^{(1)}(m) &= E(X_a^{\min}) = \int_{-\infty}^{\infty} x f_{\min}(x) dx \\ &= \int_{-\infty}^{\infty} x (m[1 - F_X(x)]^{m-1} f_X(x)) dx \\ &= m \int_{-\infty}^{\infty} x f_X(x) [1 - F_X(x)]^{m-1} dx, \end{aligned} \quad (65)$$

<sup>15</sup> where  $m$  is the sample size,  $F_X$  is the distribution function, and  $f_X$  is the density<sup>15</sup>  
<sup>16</sup>function of the data from which the minimum is derived.<sup>16</sup>

<sup>17</sup> Similarly, the second raw moment of the distribution of sample minimum is given<sup>17</sup>  
<sup>18</sup>by the following<sup>18</sup>

$$\begin{aligned} \mu_{\min}^{(2)}(m) &= E[(X_a^{\min})^2] = \int_{-\infty}^{\infty} x^2 f_{\min}(x) dx \\ &= \int_{-\infty}^{\infty} x^2 (m[1 - F_X(x)]^{m-1} f_X(x)) dx \\ &= m \int_{-\infty}^{\infty} x^2 f_X(x) [1 - F_X(x)]^{m-1} dx, \end{aligned} \quad (66)$$

<sup>24</sup> where  $m$  is the sample size,  $F_X$  is the distribution function, and  $f_X$  is the density<sup>24</sup>  
<sup>25</sup>function of the data from which the minimum is derived.<sup>25</sup>

<sup>26</sup> Using the first (Eq. 65) and second (Eq. 66) raw moments of the distribution of<sup>26</sup>  
<sup>27</sup>sample minimum, the variance is given by<sup>27</sup>

$$\sigma_{\min}^2(m) = \mu_{\min}^{(2)}(m) - \left[ \mu_{\min}^{(1)}(m) \right]^2, \quad (67)$$

<sup>31</sup>where  $m$  is the sample size of the data from which the maximum is derived and<sup>31</sup>  
<sup>32</sup> $\mu_{\min}^{(1)}(m)$  and  $\mu_{\min}^{(2)}$  are the first and second raw moments, respectively, of the distri-<sup>32</sup>  
<sup>33</sup>bution of sample minimum.<sup>33</sup>

<sup>1</sup> Using the expected attribute maximum (Eq. 62) and minimum (Eq. 65) for sample  
<sup>2</sup>size  $m$ , the following expected attribute range results from linearity of the expecta-  
<sup>3</sup>tion operator 3

$$\begin{aligned} \text{E}(X_a^{\max} - X_a^{\min}) &= \text{E}(X_a^{\max}) - \text{E}(X_a^{\min}) \\ &= \mu_{\max}^{(1)}(m) - \mu_{\min}^{(1)}(m). \end{aligned} \quad (68)$$

<sup>7</sup> where  $\mu_{\max}^{(1)}(m)$  is the expected sample maximum (Eq. 62) and  $\mu_{\min}^{(1)}(m)$  is the  
<sup>8</sup>expected sample minimum. 8

<sup>9</sup> For a data distribution whose density is an even function, the expected attribute  
<sup>10</sup>range (Eq. 68) can be simplified to the following expression 10

$$\text{E}(X_a^{\max} - X_a^{\min}) = 2\mu_{\max}^{(1)}(m), \quad (69)$$

<sup>14</sup>where  $m$  is the size of the sample from which the maximum is derived. Hence, the  
<sup>15</sup>expected attribute range is simply twice the expected attribute maximum (Eq. 62).  
<sup>16</sup>This result naturally applies to standard normal data, which is symmetric about  
<sup>17</sup>its mean at 0 and without any skewness. 17

<sup>18</sup> For large samples ( $m \gg 1$ ) from an exponential type distribution that has in-  
<sup>19</sup>finite support and all moments, the covariance between the sample maximum and  
<sup>20</sup>minimum is approximately zero [18]. In this case, the variance of the attribute range  
<sup>21</sup>of a sample of size  $m$  is given by the following 21

$$\begin{aligned} \text{Var}(X_a^{\max} - X_a^{\min}) &\approx \text{Var}(X_a^{\max}) + \text{Var}(X_a^{\min}) \\ &= \sigma_{\max}^2(m) + \sigma_{\min}^2(m). \end{aligned} \quad (70)$$

<sup>25</sup> Under the assumption of zero skewness, infinite support and even density function,  
<sup>26</sup>sufficiently large sample size  $m$ , and distribution of an exponential type for all  
<sup>27</sup>moments, the variance of attribute range (Eq. 70) simplifies to the following 27

$$\begin{aligned} \text{Var}(X_a^{\max} - X_a^{\min}) &= 2\text{Var}(X_a^{\max}) \\ &= 2\sigma_{\max}^2. \end{aligned} \quad (71)$$

<sup>31</sup>Let  $\mu_{D_{ij}^{(q)}}$  and  $\sigma_{D_{ij}^{(q)}}^2$  (Eq. 18) denote the mean and variance of the standard  $L_q$  dis-  
<sup>32</sup>tance metric (Eq. 1). Then the expected value of the max-min normalized distance 32

<sup>33</sup> 33

<sup>1</sup>(Eq. 55) distribution is given by the following

$$\begin{aligned}
 \mu_{D_{ij}^{(q*)}} &= \mathbb{E} \left[ \left( \sum_{a \in \mathcal{A}} \left( \frac{|X_{ia} - X_{ja}|}{X_a^{\max} - X_a^{\min}} \right)^q \right)^{1/q} \right] \\
 &\approx \frac{1}{\mathbb{E}(X_a^{\max} - X_a^{\min})} \mathbb{E} \left[ \left( \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q \right)^{1/q} \right] \\
 &= \frac{\mu_{D_{ij}^{(q)}}}{\mathbb{E}(X_a^{\max}) - \mathbb{E}(X_a^{\min})} \\
 &= \frac{\mu_{D_{ij}^{(q)}}}{\mu_{\max}^{(1)}(m) - \mu_{\min}^{(1)}(m)},
 \end{aligned} \tag{72}$$

<sup>10</sup>where  $m$  is the size of the sample from which extremes are derived,  $\mu_{\max}^{(1)}(m)$  is the <sup>10</sup>  
<sup>11</sup>expected value of the sample maximum (Eq. 62), and  $\mu_{\min}^{(1)}$  is the expected value of <sup>11</sup>  
<sup>12</sup>the sample minimum.

<sup>13</sup>The variance of the max-min normalized distance (Eq. 55) distribution is given <sup>13</sup>  
<sup>14</sup>by the following

$$\begin{aligned}
 \sigma_{D_{ij}^{(q*)}}^2 &= \text{Var} \left[ \left( \sum_{a \in \mathcal{A}} \left( \frac{|X_{ia} - X_{ja}|}{X_a^{\max} - X_a^{\min}} \right)^q \right)^{1/q} \right] \\
 &= \mathbb{E} \left[ \left( \sum_{a \in \mathcal{A}} \left( \frac{|X_{ia} - X_{ja}|}{X_a^{\max} - X_a^{\min}} \right)^q \right)^{2/q} \right] \\
 &\quad - \left( \mathbb{E} \left[ \left( \sum_{a \in \mathcal{A}} \left( \frac{|X_{ia} - X_{ja}|}{X_a^{\max} - X_a^{\min}} \right)^q \right)^{1/q} \right] \right)^2 \\
 &\approx \frac{\mathbb{E} \left[ \left( \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q \right)^{2/q} \right]}{\mathbb{E}[(X_a^{\max} - X_a^{\min})^2]} - \frac{\left( \mathbb{E} \left[ \left( \sum_{a \in \mathcal{A}} |X_{ia} - X_{ja}|^q \right)^{1/q} \right] \right)^2}{\mathbb{E}[(X_a^{\max} - X_a^{\min})^2]} \\
 &= \frac{\sigma_{D_{ij}^{(q)}}^2 + \mu_{D_{ij}^{(q)}}^2}{\mathbb{E}[(X_a^{\max} - X_a^{\min})^2]} - \frac{\mu_{D_{ij}^{(q)}}^2}{\mathbb{E}[(X_a^{\max} - X_a^{\min})^2]} \\
 &= \frac{\sigma_{D_{ij}^{(q)}}^2}{\mathbb{E}[(X_a^{\max} - X_a^{\min})^2]} \\
 &= \frac{\sigma_{D_{ij}^{(q)}}^2}{\mathbb{E}[(X_a^{\max})^2] - 2\mathbb{E}(X_a^{\max})\mathbb{E}(X_a^{\min}) + \mathbb{E}(X_a^{\min})^2} \\
 &= \frac{\sigma_{D_{ij}^{(q)}}^2}{\mu_{\max}^{(2)}(m) - 2\mu_{\max}^{(1)}(m)\mu_{\min}^{(1)}(m) + \mu_{\min}^{(2)}(m)},
 \end{aligned} \tag{73}$$

<sup>1</sup>where  $m$  is the size of the sample from which extremes are derived,  $\mu_{\max}^{(1)}(m)$  is the <sup>1</sup>  
<sup>2</sup>expected value of the sample maximum (Eq. 62), and  $\mu_{\min}^{(1)}$  is the expected value of <sup>2</sup>  
<sup>3</sup>the sample minimum. <sup>3</sup>

<sup>4</sup> With the mean (Eq. 72) and variance (Eq. 73) of the max-min normalized distance <sup>4</sup>  
<sup>5</sup>(Eq. 55), we have the following generalized estimate for the asymptotic distribution <sup>5</sup>  
<sup>6</sup>of the max-min normalized distance distribution <sup>6</sup>

$$D_{ij}^{(q*)} \sim \mathcal{N} \left( \frac{\mu_{D_{ij}^{(q)}}}{\mu_{\max}^{(1)}(m) - \mu_{\min}^{(1)}(m)}, \frac{\sigma_{D_{ij}^{(q)}}^2}{\mu_{\max}^{(2)}(m) - 2\mu_{\max}^{(1)}(m)\mu_{\min}^{(1)}(m) + \mu_{\min}^{(2)}(m)} \right), \quad (74)$$

<sup>7</sup>where  $m$  is the size of the sample from which extremes are derived,  $\mu_{\max}^{(1)}(m)$  is the <sup>7</sup>  
<sup>8</sup>expected value of the sample maximum (Eq. 62), and  $\mu_{\min}^{(1)}$  is the expected value of <sup>8</sup>  
<sup>9</sup>the sample minimum. <sup>9</sup>

<sup>10</sup> For data with a density function that is even, the expected sample maximum is <sup>10</sup>  
<sup>11</sup>the additive inverse of the expected sample minimum. This allows us to express the <sup>11</sup>  
<sup>12</sup>expected range-normalized pairwise distance (Eq. 72) exclusively in terms of the <sup>12</sup>  
<sup>13</sup>expected sample maximum and results in the mean of Eq. 74 becoming <sup>13</sup>

$$\mu_{D_{ij}^{(q*)}} \approx \frac{\mu_{D_{ij}^{(q)}}}{2\mu_{\max}^{(1)}(m)}, \quad (75)$$

<sup>14</sup>where  $m$  is the size of the sample from which the maximum is derived and  $\mu_{\max}^{(1)}(m)$  <sup>14</sup>  
<sup>15</sup>is the expected value of the sample maximum (Eq. 62). <sup>15</sup>

<sup>16</sup> A similar substitution gives us the following expression for the variance of the <sup>16</sup>  
<sup>17</sup>max-min normalized distance distribution <sup>17</sup>

$$\sigma_{D_{ij}^{(q*)}}^2 \approx \frac{\sigma_{D_{ij}^{(q)}}^2}{2\mu_{\max}^{(2)}(m) + 2[\mu_{\max}^{(1)}(m)]^2} = \frac{\sigma_{D_{ij}^{(q)}}^2}{2\left(\sigma_{\max}^2(m) + [\mu_{\max}^{(1)}(m)]^2\right)}, \quad (76)$$

<sup>18</sup>where  $m$  is the size of the sample from which extremes are derived,  $\mu_{\max}^{(1)}(m)$  is the <sup>18</sup>  
<sup>19</sup>expected value of the sample maximum (Eq. 62), and  $\sigma_{\max}^2(m)$  is the variance of <sup>19</sup>  
<sup>20</sup>the sample maximum (Eq. 64). <sup>20</sup>

<sup>1</sup> Therefore, the asymptotic distribution of the max-min normalized distance dis-<sup>1</sup>  
<sup>2</sup>tribution (Eq. 74) becomes <sup>2</sup>

$$D_{ij}^{(q*)} \sim \mathcal{N} \left( \frac{\mu_{D_{ij}^{(q)}}}{2\mu_{\max}^{(1)}(m)}, \frac{\sigma_{D_{ij}^{(q)}}^2}{2 \left( \sigma_{\max}^2(m) + [\mu_{\max}^{(1)}(m)]^2 \right)} \right), \quad (77)$$

<sup>3</sup> where  $m$  is the size of the sample from which extremes are derived,  $\mu_{\max}^{(1)}(m)$  is the<sup>7</sup>  
<sup>8</sup>expected value of the sample maximum (Eq. 62), and  $\sigma_{\max}^2(m)$  is the variance of<sup>8</sup>  
<sup>9</sup>the sample maximum (Eq. 64). <sup>9</sup>

<sup>10</sup> We have now derived asymptotic estimates of the moments of the max-min nor-<sup>10</sup>  
<sup>11</sup>malized  $L_q$  distance metric (Eq. 55) for any continuous data distribution. In the<sup>11</sup>  
<sup>12</sup>next two sections, we examine the max-min normalized  $L_q$  distance on standard<sup>12</sup>  
<sup>13</sup>normal and standard uniform data. As in previous sections in which we analyzed<sup>13</sup>  
<sup>14</sup>the standard  $L_q$  metric (Eq. 1), we will use the more general results for the max-min<sup>14</sup>  
<sup>15</sup> $L_q$  metric to derive asymptotic estimates for normalized Manhattan ( $q = 1$ ) and<sup>15</sup>  
<sup>16</sup>Euclidean ( $q = 2$ ). <sup>16</sup>

### <sup>17</sup> 3.4.1 Standard normal data <sup>17</sup>

<sup>18</sup> The standard normal distribution has zero skewness, even density function, infinite <sup>18</sup>  
<sup>19</sup>support, and all moments. This implies that the corresponding mean and variance <sup>19</sup>  
<sup>20</sup>of the distribution of sample range can be expressed exclusively in terms of the sam-<sup>20</sup>  
<sup>21</sup>ple maximum. Given the nature of the density function of the sample maximum for <sup>21</sup>  
<sup>22</sup>sample size  $m$ , the integration required to determine the moments (Eqs. 62 and 63) <sup>22</sup>  
<sup>23</sup>is not possible. These moments can either be approximated numerically or we can <sup>23</sup>  
<sup>24</sup>use extreme value theory to determine the form of the asymptotic distribution of <sup>24</sup>  
<sup>25</sup>the sample maximum. Using the latter method, we will show that the asymptotic <sup>25</sup>  
<sup>26</sup>distribution of the sample maximum for standard normal data is in the Gumbel <sup>26</sup>  
<sup>27</sup>family. Let  $c_m = -\Phi^{-1} \left( \frac{1}{m} \right)$  and  $b_m = \frac{1}{c_m}$ , where  $\Phi$  is the standard normal cumula-<sup>27</sup>  
<sup>28</sup>tive distribution function. Using Taylor's Theorem, we have the following expansion <sup>28</sup>  
<sup>29</sup>

$$\begin{aligned} \log \Phi(-c_m - b_m x) &= \log \Phi(-c_m) - b_m x \frac{\phi(-c_m)}{\Phi(-c_m)} + \mathcal{O}(b_m^2 x^2) \\ &= \log \left( \frac{1}{m} \right) - x \frac{\phi(-c_m)}{c_m \Phi(-c_m)} + \mathcal{O}(b_m^2 x^2), \end{aligned} \quad (78)$$

<sup>1</sup>where  $m$  is the size of the sample from which the maximum is derived.

<sup>2</sup>In order to simplify the right-hand side of this expansion (Eq. 78), we will use the<sup>2</sup>

<sup>3</sup>well known Mills Ratio Bounds [19] given by the following

$$1 \leq \frac{\phi(x)}{x\Phi(-x)} \leq 1 + \frac{1}{x^2}, \quad x > 0, \quad (79)$$

<sup>6</sup>where  $\Phi$  and  $\phi$  once again represent the cumulative distribution function and density<sup>7</sup>  
<sup>8</sup>function, respectively, of the standard normal distribution.

<sup>9</sup>The inequalities given above (Eq. 79) show that

$$\frac{\phi(x)}{x\Phi(-x)} \rightarrow 1 \text{ as } x \rightarrow \infty.$$

<sup>12</sup>This further implies that

$$\frac{\phi(c_m)}{c_m\Phi(-c_m)} \rightarrow 1 \text{ as } m \rightarrow \infty$$

<sup>15</sup>since

$$c_m = -\Phi^{-1}\left(\frac{1}{m}\right) \rightarrow \infty \text{ as } m \rightarrow \infty.$$

<sup>19</sup>This gives us the following approximation of the right-hand side of the expansion<sup>19</sup>  
<sup>20</sup>(Eq. 78) given previously

$$\begin{aligned} \log\Phi(-c_m - b_mx) &\approx \log\left(\frac{1}{m}\right) - x + \mathcal{O}(b_m^2x^2) \\ \Rightarrow \Phi(-c_m - b_mx) &\approx \frac{1}{m}e^{-x+\mathcal{O}(b_m^2x^2)} \\ \Rightarrow \Phi(c_m + b_mx) &\approx 1 - \frac{1}{m}e^{-x+\mathcal{O}(b_m^2x^2)}, \end{aligned} \quad (80)$$

<sup>25</sup>where  $m$  is the size of the sample from which the maximum is derived.

<sup>1</sup> Using the approximation of the expansion given previously (Eq. 80), we now derive<sup>1</sup>

<sup>2</sup>the limit distribution for the sample maximum in standard normal data as follows<sup>2</sup>

$$\begin{aligned}
 & \mathbb{P}\left(\frac{X_a^{\max} - c_m}{b_m} \leq x\right) = \mathbb{P}(X_a^{\max} \leq c_m + b_m x) \\
 & = \Phi^m(c_m + b_m x) \\
 & \approx \left(1 - \frac{1}{m} e^{-x + \mathcal{O}(b_m^2 x^2)}\right)^m \\
 & = \left(1 - \frac{1}{m} e^{-x + \mathcal{O}\left(\frac{1}{c_m^2} x^2\right)}\right)^m \\
 & \approx \left(1 - \frac{1}{m} e^{-x}\right)^m \\
 \Rightarrow \lim_{m \rightarrow \infty} \mathbb{P}\left(\frac{X_a^{\max} - c_m}{b_m} \leq x\right) &= \lim_{m \rightarrow \infty} \left(1 - \frac{1}{m} e^{-x}\right)^m \\
 &= e^{-e^{-x}},
 \end{aligned} \tag{81}$$

<sup>14</sup>which is the cumulative distribution function of the standard Gumbel distribution.<sup>14</sup>

<sup>15</sup>The mean of this distribution is given by the following<sup>15</sup>

$$\mathbb{E}(X_a^{\max}) = \mu_{\max}^{(1)} = -\Phi^{-1}\left(\frac{1}{m}\right) - \frac{\gamma}{\Phi^{-1}\left(\frac{1}{m}\right)}, \tag{82}$$

<sup>18</sup>where  $m$  is the size of the sample from which the maximum is derived and  $\gamma$  is the<sup>18</sup>

<sup>19</sup>well known Euler-Mascheroni constant. This constant has many equivalent defini-<sup>19</sup>

<sup>20</sup>tions, one of which is given by<sup>20</sup>

$$\gamma = \lim_{m \rightarrow \infty} \left( -\log(m) + \sum_{k=1}^m \frac{1}{k} \right).$$

<sup>24</sup>Perhaps a more convenient definition of the Euler-Mascheroni constant is simply<sup>24</sup>

$$\gamma = -\Gamma'(1) = \frac{d}{dt} \left( \int_0^\infty z^{t-1} e^{-z} dz \right) \Big|_{t=1},$$

<sup>27</sup>which is just the additive inverse of the first derivative of the gamma function<sup>27</sup>

<sup>28</sup>evaluated at 1.<sup>28</sup>

<sup>29</sup> The median of the distribution of the maximum for standard normal data is given<sup>29</sup>

<sup>30</sup>by<sup>30</sup>

$$\tilde{\mu}_{\max} = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right), \tag{83}$$

<sup>33</sup>where  $m$  is the size of the sample from which the maximum is derived.<sup>33</sup>



Finally, the variance of the asymptotic distribution of the sample maximum is<sup>1</sup>  
given by<sup>2</sup>

$$\text{Var}(X_a^{\max}) = \frac{\pi^2}{6} \left( \frac{1}{-\Phi^{-1}\left(\frac{1}{m}\right)} \right)^2, \quad (84)$$

where  $m$  is the size of the sample from which the maximum is derived.

For typical sample sizes  $m$  in high-dimensional spaces, the variance estimate<sup>7</sup>  
(Eq. 84) exceeds the variance of the sample maximum significantly. Using the fact<sup>8</sup>  
that<sup>9</sup>

$$-\Phi^{-1}\left(\frac{1}{m}\right) \sim \sqrt{2\log(m)} \quad [20]$$

and<sup>12</sup>

$$\frac{1}{2\log(m)} \leq \left( \frac{1}{-\Phi^{-1}\left(\frac{1}{m}\right)} \right)^2, \quad m \geq 2,$$

we can get a more accurate approximation of the variance with the following

$$\begin{aligned} \sigma_{\max}^2(m) = \text{Var}(X_a^{\max}) &\approx \frac{\pi^2}{6} \left( \frac{1}{\sqrt{2\log(m)}} \right)^2 \\ &= \frac{\pi^2}{12\log(m)}. \end{aligned} \quad (85)$$

Therefore, the mean of the range of  $m$  iid standard normal random variables is<sup>21</sup>  
given by<sup>22</sup>

$$\text{E}(X_a^{\max} - X_a^{\min}) = 2\mu_{\max}^{(1)}(m) = 2 \left[ -\Phi^{-1}\left(\frac{1}{m}\right) - \frac{\gamma}{\Phi^{-1}\left(\frac{1}{m}\right)} \right], \quad (86)$$

where  $\gamma$  is the Euler-Mascheroni constant.

It is well known that the sample extremes from the standard normal distribution<sup>26</sup>  
are approximately uncorrelated for large sample size  $m$  [18]. This implies that we can<sup>27</sup>  
approximate the variance of the range of  $m$  iid standard normal random variables<sup>28</sup>

<sup>1</sup>with the following result

$$\begin{aligned}
 & \text{Var}(X_a^{\max} - X_a^{\min}) \approx \text{Var}(X_a^{\max}) + \text{Var}(X_a^{\min}) \\
 & = \sigma_{\max}^2(m) + \sigma_{\min}^2(m) \\
 & = 2\sigma_{\max}^2(m) \\
 & \approx 2 \left( \frac{\pi^2}{12\log(m)} \right) \\
 & = \frac{\pi^2}{6\log(m)}.
 \end{aligned} \tag{87}$$

<sup>9</sup> For the purpose of approximating the mean and variance of the max-min normal-  
<sup>10</sup>ized distance distribution, we observe empirically that the formula for the median<sup>10</sup>  
<sup>11</sup>of the distribution of the attribute maximum (Eq. 83) yields more accurate results.<sup>11</sup>  
<sup>12</sup>More precisely, the approximation of the expected maximum (Eq. 82) overestimates<sup>12</sup>  
<sup>13</sup>the sample maximum slightly. The formula for the median of the sample maximum<sup>13</sup>  
<sup>14</sup>(Eq. 83) provides a more accurate estimate of this sample extreme. Therefore, the<sup>14</sup>  
<sup>15</sup>following estimate for the mean of the attribute range will be used instead

$$\text{E}(X_a^{\max} - X_a^{\min}) = 2\mu_{\max}^{(1)}(m) \approx 2 \left[ \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right) \right], \tag{88}$$

<sup>18</sup>where  $m$  is the size of the sample from which extremes are derived.

<sup>19</sup> We have already determined the mean and variance (Eq. 27) for the  $L_q$  metric  
<sup>20</sup>(Eq. 1) on standard normal data. Using the expected value of the sample maximum  
<sup>21</sup>(Eq. 88), the variance of the sample maximum (Eq. 87), and the general formulas for  
<sup>22</sup>the mean and variance of the max-min normalized distance distribution (Eq. 77),  
<sup>23</sup>this leads us to the following asymptotic estimate for the distribution of the max-  
<sup>24</sup>min normalized distances for standard normal data

$$D_{ij}^{(q*)} \sim \mathcal{N} \left( \frac{\mu_{D_{ij}^{(q)}}}{2\mu_{\max}^{(1)}(m)}, \frac{6\log(m)\sigma_{D_{ij}^{(q)}}^2}{\pi^2 + 24 \left[ \mu_{\max}^{(1)}(m) \right]^2 \log(m)} \right). \tag{89}$$

<sup>28</sup>where  $m$  is the size of the sample from which the maximum is derived,  $\mu_{\max}^{(1)}$  is  
<sup>29</sup>the median of the sample maximum (Eq. 83),  $\mu_{D_{ij}^{(q)}}$  is the expected  $L_q$  pairwise  
<sup>30</sup>distance (Eq. 25), and  $\sigma_{D_{ij}^{(q)}}^2$  is the variance of the  $L_q$  pairwise distance (Eq. 26).  
<sup>31</sup>The moments of the max-min normalized  $L_q$  distance metric in standard normal  
<sup>32</sup>data (Eq. 89) are summarized in a table that is organized by metric, data type,  
<sup>33</sup>statistic (mean or variance), and asymptotic formula (Table 1).

### <sup>1</sup>3.4.2 *Standard uniform data* <sup>1</sup>

<sup>2</sup>Standard uniform data does not have an even density function. Due to the simplicity<sup>2</sup>  
<sup>3</sup>of the density function, however, we can derive the distribution of the maximum and<sup>3</sup>  
<sup>4</sup>minimum of a sample of size  $m$  explicitly. Using the general forms of the distribution<sup>4</sup>  
<sup>5</sup>functions of the maximum (Eq. 57) and minimum (Eq. 60), we have the following<sup>5</sup>  
<sup>6</sup>distribution functions for standard uniform data <sup>6</sup>

$$\text{<sup>8</sup> } F_{\max}(x) = x^m \text{ <sup>8</sup> (90)}$$

<sup>9</sup> and <sup>9</sup>

<sup>10</sup> <sup>10</sup>

$$\text{<sup>11</sup> } F_{\min}(x) = 1 - (1 - x)^m, \text{ <sup>11</sup> (91)}$$

<sup>12</sup> <sup>12</sup>

<sup>13</sup>where  $m$  is the size of the sample from which extremes are derived. <sup>13</sup>

<sup>14</sup> Using the general forms of the density functions of the maximum (Eq. 58) and<sup>14</sup>

<sup>15</sup>minimum (Eq. 61), we have the following density functions for standard uniform<sup>15</sup>

<sup>16</sup>data <sup>16</sup>

<sup>17</sup> <sup>17</sup>

$$\text{<sup>18</sup> } f_{\max}(x) = mx^{m-1} \text{ <sup>18</sup> (92)}$$

<sup>19</sup> and <sup>19</sup>

<sup>20</sup> <sup>20</sup>

$$\text{<sup>21</sup> } f_{\min}(x) = m(1 - x)^{m-1}, \text{ <sup>21</sup> (93)}$$

<sup>22</sup> <sup>22</sup>

<sup>23</sup>where  $m$  is the size of the sample from which extremes are derived. <sup>23</sup>

<sup>24</sup> Then the expected maximum and minimum are computed through <sup>24</sup>

<sup>25</sup>straightforward integration as follows <sup>25</sup>

<sup>26</sup> <sup>26</sup>

$$\text{<sup>27</sup> } E(X_a^{\max}) = \mu_{\max}^{(1)}(m) = \int_0^1 x f_{\max}(x) dx \text{ <sup>27</sup> }$$

$$\text{<sup>28</sup> } = \int_0^1 x [mx^{m-1}] dx \text{ <sup>28</sup> (94)}$$

$$\text{<sup>29</sup> } = \frac{m}{m+1} \text{ <sup>29</sup> }$$

<sup>30</sup> <sup>30</sup>

<sup>31</sup> <sup>31</sup>

<sup>32</sup> <sup>32</sup>

<sup>33</sup> <sup>33</sup>

<sup>1</sup>and

$$\begin{aligned}
 \text{E}(X_a^{\min}) &= \mu_{\min}^{(1)}(m) = \int_0^1 x f_{\min}(x) dx \\
 &= \int_0^1 x [m(1-x)^{m-1}] dx \\
 &= \frac{1}{m+1},
 \end{aligned}
 \tag{95}$$

<sup>7</sup>where  $m$  is the size of the sample from which extremes are derived.

<sup>8</sup> We can compute the second moment about the origin of the sample range as<sup>8</sup>  
<sup>9</sup>follows

$$\begin{aligned}
 \text{E}[(X_a^{\max} - X_a^{\min})^2] &= \text{E}[(X_a^{\max})^2 - 2X_a^{\max}X_a^{\min} + (X_a^{\min})^2] \\
 &= \text{E}[(X_a^{\max})^2] - 2\text{E}(X_a^{\max})\text{E}(X_a^{\min}) + \text{E}[(X_a^{\min})^2] \\
 &= \mu_{\max}^{(2)}(m) - 2\mu_{\max}^{(1)}(m)\mu_{\min}^{(1)}(m) + \mu_{\min}^{(2)}(m) \\
 &= \int_0^1 x^2 [mx^{m-1}] dx - 2 \left( \frac{m}{m+1} \right) \left( \frac{1}{m+1} \right) \\
 &\quad + \int_0^1 x^2 [m(1-x)^{m-1}] dx \\
 &= \frac{m}{m+2} - \frac{2m}{(m+1)^2} + \frac{2}{(m+1)(m+2)} \\
 &= \frac{m^3 - m + 2}{(m+2)(m+1)^2},
 \end{aligned}
 \tag{96}$$

<sup>20</sup>where  $m$  is the size of the sample from which extremes are derived.

<sup>21</sup> Using the general asymptotic distribution of max-min normalized distances for<sup>21</sup>  
<sup>22</sup>any data type (Eq. 74) and the mean and variance (Eq. 37) of the standard  $L_q$ <sup>22</sup>  
<sup>23</sup>distance metric (Eq. 1), we have the following asymptotic estimate for the max-min<sup>23</sup>  
<sup>24</sup>normalized distance distribution for standard uniform data

$$D_{ij}^{(q*)} \sim \mathcal{N} \left( \frac{(m+1)\mu_{D_{ij}^{(q)}}}{m-1}, \frac{(m+2)(m+1)^2\sigma_{D_{ij}^{(q)}}^2}{m^3 - m + 2} \right),
 \tag{97}$$

<sup>28</sup> where  $m$  is the size of the sample from which extremes are derived,  $\mu_{D_{ij}^{(q)}}$  is the<sup>28</sup>  
<sup>29</sup>expected value (Eq. 35) of the  $L_q$  metric (Eq. 1) in standard uniform data, and<sup>29</sup>  
<sup>30</sup> $\sigma_{D_{ij}^{(q)}}^2$  is the variance (Eq. 36) of the  $L_q$  metric (Eq. 1) in standard uniform data.<sup>30</sup>  
<sup>31</sup>The moments of the max-min normalized  $L_q$  distance metric in standard uniform<sup>31</sup>  
<sup>32</sup>data (Eq. 89) are summarized in a table that is organized by metric, data type,<sup>32</sup>  
<sup>33</sup>statistic (mean or variance), and asymptotic formula (Table 1).<sup>33</sup>

### 3.5 Range-Normalized Manhattan ( $q = 1$ )

Using the general asymptotic results for mean and variance of max-min normalized distances in standard normal and standard uniform data (Eqs. 89 and 97) for any value of  $q \in \mathbb{N}$ , we can substitute a particular value of  $q$  in order to determine a more specified distribution for the normalized distance ( $D^{(q*)}$ , Eq. 55). The following results are for the max-min normalized Manhattan ( $q = 1$ ),  $D^{(1*)}$ , metric for both standard normal and standard uniform data.

#### 3.5.1 Standard normal data

Substituting  $q = 1$  into the asymptotic formula for the expected max-min normalized distance (Eq. 89), we derive the expected normalized Manhattan distance in standard normal data as follows

$$\begin{aligned} E\left(D_{ij}^{(1*)}\right) &= \frac{\mu_{D_{ij}^{(1)}}}{2\mu_{\max}^{(1)}(m)} \\ &= \frac{p}{\sqrt{\pi}\mu_{\max}^{(1)}(m)}, \end{aligned} \quad (98)$$

where  $\mu_{\max}^{(1)}(m)$  is the expected attribute maximum (Eq. 83),  $m$  is the size of the sample from which the maximum is derived, and  $p$  is the total number of attributes.

Similarly, the variance of  $D_{ij}^{(1*)}$  is given by

$$\begin{aligned} \text{Var}\left(D_{ij}^{(1*)}\right) &= \frac{6\log(m)\sigma_{D_{ij}^{(1)}}^2}{\pi^2 + 24\left[\mu_{\max}^{(1)}\right]^2\log(m)} \\ &= \frac{12p(\pi - 2)\log(m)}{\pi\left(\pi^2 + 24\left[\mu_{\max}^{(1)}\right]^2\log(m)\right)}, \end{aligned} \quad (99)$$

where  $\mu_{\max}^{(1)}(m)$  is the expected attribute maximum (Eq. 83),  $m$  is the size of the sample from which the maximum is derived, and  $p$  is the total number of attributes.

Similar to the variance of the standard Manhattan distance, the variance of the max-min normalized Manhattan distance is on the order of  $p$  for fixed instance dimension

$m$ . For fixed  $p$ , the variance (Eq. 99) vanishes as  $m$  grows without bound. If we fix  $m$ ,

the same variance increases monotonically with increasing  $p$ . The moments derived

in this section (Eqs. 98 and 99) are summarized in a table that is organized by

metric, data type, statistic (mean or variance), and asymptotic formula (Table 3).

### 3.5.2 Standard uniform data

Substituting  $q = 1$  into the asymptotic formula for the expected max-min pairwise distance (Eq. 97), we derive the expected normalized Manhattan distance in standard uniform data as

$$\begin{aligned} \mathbb{E}\left(D_{ij}^{(1*)}\right) &= \frac{(m+1)\mu_{D_{ij}^{(1)}}}{m-1} \\ &= \frac{(m+1)p}{3(m-1)}, \end{aligned} \quad (100)$$

where  $m$  is the size of the sample from which extremes are derived and  $p$  is the total number attributes.

Similarly, the variance of  $D_{ij}^{(1*)}$  is given by

$$\begin{aligned} \text{Var}\left(D_{ij}^{(1*)}\right) &= \frac{(m+2)(m+1)^2\sigma_{D_{ij}^{(1)}}^2}{m^3 - m + 2} \\ &= \frac{(m+2)(m+1)^2p}{18(m^3 - m + 2)}, \end{aligned} \quad (101)$$

where  $m$  is the size of the sample from which extremes are derived and  $p$  is the total number of attributes. Interestingly, the variance of the max-min normalized Manhattan distance in standard uniform data approaches  $p/18$  as  $m$  increases without bound for a fixed number of attributes  $p$ . This is the same asymptotic value to which the variance of the standard Manhattan distance (Eq. 40) converges. Therefore, large sample sizes make the variance of the normalized Manhattan distance approach the variance of the standard Manhattan distance in standard uniform data. The moments derived in this section (Eqs. 100 and 101) are summarized in a table that is organized by metric, data type, statistic (mean or variance), and asymptotic formula (Table 3).

### 3.6 Range-Normalized Euclidean ( $q = 2$ )

Analogous to the previous section, we use the asymptotic moment estimates for the max-min normalized metric ( $D^{(q*)}$ , Eq. 55) for standard normal (Eq. 89) and standard uniform (Eq. 97) data but specific to a range-normalized Euclidean metric ( $q = 2$ ).

### 3.6.1 Standard normal data

Substituting  $q = 2$  into the asymptotic formula for the expected max-min normalized pairwise distance (Eq. 89), we derive the expected normalized Euclidean distance in standard normal data as

$$\begin{aligned} E\left(D_{ij}^{(2*)}\right) &= \frac{\mu_{ij}^{D^{(2)}}}{2\mu_{\max}^{(1)}(m)} \\ &= \frac{\sqrt{2p-1}}{2\mu_{\max}^{(1)}(m)}, \end{aligned} \quad (102)$$

where  $\mu_{\max}^{(1)}(m)$  is the expected attribute maximum (Eq. 83),  $m$  is the size of the sample from which the maximum is derived, and  $p$  is the total number of attributes.

Similarly, the variance of  $D_{ij}^{(2*)}$  is given by

$$\begin{aligned} \text{Var}\left(D_{ij}^{(2*)}\right) &= \frac{6\log(m)\sigma_{D_{ij}^{(2)}}^2}{\pi^2 + 24\left[\mu_{\max}^{(1)}(m)\right]^2\log(m)} \\ &= \frac{6\log(m)}{\pi^2 + 24\left[\mu_{\max}^{(1)}(m)\right]^2\log(m)}, \end{aligned} \quad (103)$$

where  $\mu_{\max}^{(1)}(m)$  is the expected attribute maximum (Eq. 83) and  $m$  is the size of the sample from which the maximum is derived. It is interesting to note that the variance (Eq. 103) vanishes as the sample size  $m$  increases without bound, which means that all distances will be tightly clustered about the mean (Eq. 102). This is different than the variance of the standard  $L_2$  metric (Eq. 49), which is asymptotically equal to 1. This could imply that any two pairwise distances computed with the max-min normalized Euclidean metric in a large sample space  $m$  may be indistinguishable, which is another curse of dimensionality. The moments derived in this section (Eqs. 102 and 103) are summarized in a table that is organized by metric, data type, statistic (mean or variance), and asymptotic formula (Table 3).

### 3.6.2 Standard uniform data

Substituting  $q = 2$  into the asymptotic formula for the expected max-min normalized pairwise distance (Eq. 97), we derive the expected normalized Euclidean

distance in standard uniform data as

$$\begin{aligned} \mathbb{E}\left(D_{ij}^{(2*)}\right) &= \frac{(m+1)\mu_{D_{ij}^{(2)}}}{m-1} \\ &= \sqrt{\frac{p}{6} - \frac{7}{120}} \left(\frac{m+1}{m-1}\right). \end{aligned} \quad (104)$$

where  $m$  is the size of the sample from which extremes are derived and  $p$  is the total number of attributes.

Similarly, the variance of  $D_{ij}^{(2*)}$  is given by

$$\begin{aligned} \text{Var}\left(D_{ij}^{(2*)}\right) &= \frac{(m+2)(m+1)^2\sigma_{D_{ij}^{(2)}}^2}{m^3 - m + 2} \\ &= \frac{7(m+2)(m+1)^2}{120(m^3 - m + 2)}. \end{aligned} \quad (105)$$

where  $m$  is the size of the sample from which extremes are derived. Similar to the variance of max-min normalized Manhattan distances in standard uniform data (Eq. 101), the variance of normalized Euclidean distances approaches the variance of the standard Euclidean distances in uniform data (Eq. 52) as  $m$  increases without bound. That is, the variance of the max-min normalized Euclidean distance (Eq. 105) approaches  $7/120$  as  $m$  grows larger. The moments derived in this section (Eqs. 104 and 105) are summarized in a table that is organized by metric, data type, statistic (mean or variance), and asymptotic formula (Table 3).

We summarize moment estimates in tables (Tables 1-3) that contain all of our asymptotic results for both standard and max-min normalized  $L_q$  metrics in each data type we have considered. This includes our most general results for any combination of sample size  $m$ , number of attributes  $p$ , type of metric  $L_q$ , and data type (Table 1). From these more general derivations, we show the results of the standard  $L_1$  and  $L_2$  metrics for any combination of sample size  $m$ , number of attributes  $p$ , and data type (Table 2). Our last set of summarized results show asymptotics for the max-min normalized  $L_1$  and  $L_2$  metrics for any combination of sample size  $m$ , number of attributes  $p$ , and data type (Table 3). For both standard and max-min normalized  $L_2$  metrics (Tables 2 and 3), the low-dimensional improved estimates of sample means (Eqs. 50 and 53) are used because they perform well at both low and high attribute dimension  $p$ .



<sup>1</sup> In the next section, we make a transition into discrete GWAS data. We will discuss<sup>1</sup>  
<sup>2</sup>some commonly known metrics and then a relatively new metric, which will lead<sup>2</sup>  
<sup>3</sup>us into novel asymptotic results for this data type. <sup>3</sup>

<sup>4</sup>

<sup>4</sup>

#### <sup>5</sup>4 GWAS distance distributions <sup>5</sup>

<sup>6</sup>In genome-wide association studies (GWAS), data is encoded by the minor allele<sup>6</sup>  
<sup>7</sup>at a particular locus  $a$ , which is just the second most common allele (adenine-A,<sup>7</sup>  
<sup>8</sup>thymine-T, cytosine-C, or guanine-G) associated with a given attribute  $a$  in the<sup>8</sup>  
<sup>9</sup>data set. Attributes in GWAS data are single nucleotide polymorphisms (SNPs),<sup>9</sup>  
<sup>10</sup>or mutations involving the substitution, deletion, or insertion of one nucleotide at<sup>10</sup>  
<sup>11</sup>some point in the DNA sequence of an organism. These are common mutations<sup>11</sup>  
<sup>12</sup>that can affect how an individual reacts to certain pathogens or the susceptibility<sup>12</sup>  
<sup>13</sup>for certain diseases. Feature selection in GWAS is typically concerned with finding<sup>13</sup>  
<sup>14</sup>interacting SNPs that are associated with disease susceptibility [21]. <sup>14</sup>

<sup>15</sup> Consider a GWAS data set, which has the following encoding based on minor<sup>15</sup>  
<sup>16</sup>allele frequency <sup>16</sup>

$$\begin{aligned} & \supset 0 \quad \text{if there are no minor alleles at locus } a, \\ & X_{ia} = \begin{cases} 1 & \text{if there is 1 minor allele at locus } a, \\ 2 & \text{if there are 2 minor alleles at locus } a. \end{cases} \quad (106) \end{aligned}$$

<sup>21</sup> A minor allele at a particular locus  $a$  is the least frequent of the two alleles at that <sup>21</sup>  
<sup>22</sup>particular locus  $a$ . For random GWAS data sets, we can think  $X_{ia}$  as the number of <sup>22</sup>  
<sup>23</sup>successes in two Bernoulli trials. That is,  $X_{ia} \sim \mathcal{B}(2, f_a)$  where  $f_a$  is the probability <sup>23</sup>  
<sup>24</sup>of success. The success probability  $f_a$  is the probability of a minor allele occurring <sup>24</sup>  
<sup>25</sup>at  $a$ . Furthermore, the minor allele probabilities are assumed to be independent and <sup>25</sup>  
<sup>26</sup>identically distributed according to  $\mathcal{U}(l, u)$ , where  $l$  and  $u$  are the lower and upper <sup>26</sup>  
<sup>27</sup>bounds, respectively, of the sampling distribution's support. Two commonly known <sup>27</sup>  
<sup>28</sup>types of distance metrics for GWAS data are the Genotype Mismatch (GM) and <sup>28</sup>  
<sup>29</sup>Allele Mismatch (AM) metrics. The GM and AM metrics are defined by <sup>29</sup>

$$\begin{aligned} & \supset 0 \quad \text{if } X_{ia} \neq X_{ja}, \\ & d_{ij}^{\text{GM}}(a) = \begin{cases} 1 & \text{otherwise} \end{cases} \quad (107) \end{aligned}$$

and

$$d_{ij}^{\text{AM}}(a) = \frac{1}{2} |X_{ia} - X_{ja}|. \quad (108)$$

More informative metrics may include differences at the nucleotide level for each allele by considering differences in the rates of transition and transversion mutations (Fig. 2). One such discrete metric that accounts for transitions (Ti) and transversions (Tv) was introduced in [7] and can be written as

$$d_{ij}^{\text{TiTv}}(a) = \begin{cases} 0 & \text{if } X_{ia} = X_{ja} \text{ and Ti/Tv,} \\ 1/4 & \text{if } |X_{ia} - X_{ja}| = 1 \text{ and Ti,} \\ 1/2 & \text{if } |X_{ia} - X_{ja}| = 1 \text{ and Tv,} \\ 3/4 & \text{if } |X_{ia} - X_{ja}| = 2 \text{ and Ti,} \\ 1 & \text{if } |X_{ia} - X_{ja}| = 2 \text{ and Tv.} \end{cases} \quad (109)$$

With these GWAS distance metrics, we then compute the pairwise distance between two instances  $i, j \in \mathcal{I}$  with

$$D_{ij}^{\text{GM}}(a) = \sum_{a \in \mathcal{A}} d_{ij}^{\text{GM}}(a), \quad (110)$$

$$D_{ij}^{\text{AM}}(a) = \sum_{a \in \mathcal{A}} d_{ij}^{\text{AM}}(a), \text{ or} \quad (111)$$

$$D_{ij}^{\text{TiTv}}(a) = \sum_{a \in \mathcal{A}} d_{ij}^{\text{TiTv}}(a). \quad (112)$$

Assuming that all data entries  $X_{ia}$  are independent and identically distributed, we have already shown that the distribution of pairwise distances is asymptotically normal regardless of data distribution and value of  $q$ . Therefore, the distance distributions induced by each of the GWAS metrics (Eqs. 107-109) are asymptotically normal. With this Gaussian limiting behavior, we will proceed by deriving the mean and variance for each distance distribution induced by these three GWAS metrics.

#### 4.1 GM distance distribution

The simplest distance metric in nearest-neighbor feature selection in GWAS data<sup>2</sup> is the genotype-mismatch (GM) distance metric (Eq. 110). The GM attribute diff<sup>3</sup> (Eq. 107) indicates only whether two genotypes are the same or not. There are many<sup>4</sup> ways two genotypes could differ, but this metric does not record this information.<sup>5</sup> We will now derive the moments for the GM distance (Eq. 110), which are sufficient<sup>6</sup> for defining its corresponding asymptotic distribution.<sup>7</sup>

The expected value of the GM attribute diff metric (Eq. 107) is given by the<sup>8</sup> following<sup>9</sup>

$$\begin{aligned}
 E \left[ d_{ij}^{\text{GM}}(a) \right] &= \sum_{k=0}^1 k \cdot P \left[ d_{ij}^{\text{GM}}(a) = k \right] \\
 &= 0 \cdot P \left[ d_{ij}^{\text{GM}}(a) = 0 \right] + 1 \cdot P \left[ d_{ij}^{\text{GM}}(a) = 1 \right] \\
 &= P \left[ d_{ij}^{\text{GM}}(a) = 1 \right] \\
 &= 2P[X_{ia} = 0, X_{ja} = 1] + 2P[X_{ia} = 1, X_{ja} = 2] \\
 &\quad + 2P[X_{ia} = 0, X_{ja} = 2] \\
 &= 4(1 - f_a)^3 f_a + 4(1 - f_a) f_a^3 + 2(1 - f_a)^2 f_a^2 \\
 &= 2 \left[ 2(1 - f_a)^3 f_a + 2(1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2 \right] \\
 &= 2F^{\text{GM}}(a),
 \end{aligned} \tag{113}$$

where  $F^{\text{GM}}(a) = 2(1 - f_a)^3 f_a + 2(1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2$  and  $f_a$  is the probability<sup>21</sup> of a minor allele occurring at locus  $a$ .<sup>22</sup>

Then the expected pairwise GM distance between instances  $i, j \in \mathcal{I}$  is given by<sup>23</sup>

$$\begin{aligned}
 E \left( D_{ij}^{\text{GM}} \right) &= E \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{GM}}(a) \right) \\
 &= \sum_{a \in \mathcal{A}} E \left[ d_{ij}^{\text{GM}}(a) \right] \\
 &= 2 \sum_{a \in \mathcal{A}} F^{\text{GM}}(a),
 \end{aligned} \tag{114}$$

where  $F^{\text{GM}}(a) = 2(1 - f_a)^3 f_a + 2(1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2$  and  $f_a$  is the probability<sup>30</sup> of a minor allele occurring at locus  $a$ . We see that the expected GM pairwise distance<sup>31</sup> (Eq. 114) relies only on the minor allele probabilities  $f_a$  for all  $a \in \mathcal{A}$ . In real data,<sup>32</sup> we can easily determine these probabilities by dividing the total number of minor<sup>33</sup>

<sup>1</sup>alleles at locus  $a$  by the twice the number of instances  $m$ . To be more explicit, this<sup>1</sup>  
<sup>2</sup>is just<sup>2</sup>

$$f_a = \frac{1}{2m} \sum_{i \in \mathcal{I}} X_{ia}, \quad \text{for all } a \in \mathcal{A},$$

<sup>3</sup>where  $m$  is the number of instances (or sample size). This is because each instance<sup>6</sup>  
<sup>7</sup>has two alleles, the minor and major alleles, at each locus. Therefore, the total<sup>7</sup>  
<sup>8</sup>number of alleles at locus  $a$  is  $2m$ .<sup>8</sup>

<sup>9</sup>The second moment about the origin for the GM distance is computed as follows<sup>9</sup>

$$\begin{aligned} \mathbb{E} \left[ (D_{ij}^{\text{GM}})^2 \right] &= \mathbb{E} \left[ \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{GM}}(a) \right)^2 \right] \\ &= \mathbb{E} \left[ \sum_{a \in \mathcal{A}} \left( d_{ij}^{\text{GM}}(a) \right)^2 \right] + 2\mathbb{E} \left[ \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} d_{ij}^{\text{GM}}(r) \cdot d_{ij}^{\text{GM}}(s) \right] \\ &= \sum_{a \in \mathcal{A}} \left( \sum_{k=0}^1 k^2 \cdot \mathbb{P} \left[ d_{ij}^{\text{GM}}(a) = k \right] \right) \\ &\quad + 2 \sum_{a \in \mathcal{A}} \sum_{s \leq r-1} \left( \sum_{k=0}^1 k \cdot \mathbb{P} \left[ d_{ij}^{\text{GM}}(r) = k \right] \right) \cdot \left( \sum_{k=0}^1 k \cdot \mathbb{P} \left[ d_{ij}^{\text{GM}}(s) = k \right] \right) \\ &= 2 \sum_{a \in \mathcal{A}} F^{\text{GM}}(a) + 8 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r, s\}} F^{\text{GM}}(\lambda), \end{aligned} \tag{115}$$

<sup>21</sup>where  $F^{\text{GM}}(a) = 2(1 - f_a)^3 f_a + 2(1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2$  and  $f_a$  is the probability<sup>21</sup>  
<sup>22</sup>of a minor allele occurring at locus  $a$ .<sup>22</sup>

<sup>23</sup>Using the first (Eq. 114) and second (Eq. 115) raw moments of the GM distance,<sup>23</sup>  
<sup>24</sup>the variance is given by<sup>24</sup>

$$\begin{aligned} \text{Var} (D_{ij}^{\text{GM}}) &= \mathbb{E} \left[ (D_{ij}^{\text{GM}})^2 \right] - [\mathbb{E} (D_{ij}^{\text{GM}})]^2 \\ &= 2 \sum_{a \in \mathcal{A}} F^{\text{GM}}(a) + 8 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r, s\}} F^{\text{GM}}(\lambda) \\ &\quad - 4 \left( \sum_{a \in \mathcal{A}} F^{\text{GM}}(a) \right)^2 \\ &= 2 \sum_{a \in \mathcal{A}} F^{\text{GM}}(a) - 4 \sum_{a \in \mathcal{A}} [F^{\text{GM}}(a)]^2 \\ &= 2 \sum_{a \in \mathcal{A}} F^{\text{GM}}(a) [1 - 2F^{\text{GM}}(a)], \end{aligned} \tag{116}$$

<sup>1</sup>where  $F^{\text{GM}}(a) = 2(1 - f_a)^3 f_a + 2(1 - f_a)f_a^3 + (1 - f_a)^2 f_a^2$  and  $f_a$  is the probability<sup>1</sup>  
<sup>2</sup>of a minor allele occurring at locus  $a$ . Hence, the variance of the asymptotic GM<sup>2</sup>  
<sup>3</sup>distance distribution also just depends on the minor allele probabilities  $f_a$  for all<sup>3</sup>  
<sup>4</sup> $a \in \mathcal{A}$ . This implies that the limiting GM distance distribution is fully determined<sup>4</sup>  
<sup>5</sup>by the minor allele probabilities, which are known in real data.<sup>5</sup>

<sup>6</sup> With the mean and variance estimates (Eqs. 114 and 116), the asymptotic GM<sup>6</sup>  
<sup>7</sup>distance distribution is given by the following<sup>7</sup>

$$D_{ij}^{\text{GM}} \sim \mathcal{N} \left( 2 \sum_{a \in \mathcal{A}} F^{\text{GM}}(a), 2 \sum_{a \in \mathcal{A}} F^{\text{GM}}(a)[1 - 2F^{\text{GM}}(a)] \right), \quad (117)$$

<sup>8</sup>where  $F^{\text{GM}}(a) = 2(1 - f_a)^3 f_a + 2(1 - f_a)f_a^3 + (1 - f_a)^2 f_a^2$  and  $f_a$  is the probability<sup>11</sup>  
<sup>12</sup>of a minor allele occurring at locus  $a$ . This GM distribution holds for random<sup>12</sup>  
<sup>13</sup>independent GWAS data with minor allele probabilities  $f_a$  and binomial samples<sup>13</sup>  
<sup>14</sup> $X_{ia} \sim \mathcal{B}(2, f_a)$  for all  $a \in \mathcal{A}$ . Next we consider the distance distribution for an<sup>14</sup>  
<sup>15</sup>AM metric, which incorporates differences at the allele level and contains more<sup>15</sup>  
<sup>16</sup>information than genotype differences.<sup>16</sup>

## <sup>17</sup> 4.2 AM distance distribution<sup>17</sup>

<sup>18</sup>As we have mentioned previously, the AM attribute diff metric (Eq. 108) is slightly<sup>18</sup>  
<sup>19</sup>more dynamic than the GM metric because the AM metric accounts for differences<sup>19</sup>  
<sup>20</sup>between the alleles of two genotypes. In this section, we derive moments of the<sup>20</sup>  
<sup>21</sup>AM distance metric (Eq. 111) that adequately define its corresponding asymptotic<sup>21</sup>  
<sup>22</sup>distribution.<sup>22</sup>

<sup>1</sup> The expected value of the AM attribute diff metric (Eq. 108) is given by the<sup>1</sup>  
<sup>2</sup>following<sup>2</sup>

$$\begin{aligned}
 \text{E} \left[ d_{ij}^{\text{AM}}(a) \right] &= \sum_{k \in \mathcal{D}} k \cdot \text{P} \left[ d_{ij}^{\text{AM}}(a) = k \right] \\
 &= 0 \cdot \text{P} \left[ d_{ij}^{\text{AM}}(a) = 0 \right] + \frac{1}{2} \cdot \text{P} \left[ d_{ij}^{\text{AM}}(a) = \frac{1}{2} \right] \\
 &\quad + 1 \cdot \text{P} \left[ d_{ij}^{\text{AM}}(a) = 1 \right] \\
 &= \frac{1}{2} (2\text{P} [X_{ia} = 0, X_{ja} = 1] + 2\text{P} [X_{ia} = 1, X_{ja} = 2]) \\
 &\quad + 2\text{P} [X_{ia} = 0, X_{ja} = 2] \quad (118)^9 \\
 &= \text{P} [X_{ia} = 0, X_{ja} = 1] + \text{P} [X_{ia} = 1, X_{ja} = 2] \\
 &\quad + 2\text{P} [X_{ia} = 0, X_{ja} = 2] \\
 &= 2(1 - f_a)^3 f_a + 2(1 - f_a) f_a^3 + 2(1 - f_a)^2 f_a^2 \\
 &= 2 \left[ (1 - f_a)^3 f_a + (1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2 \right] \\
 &= 2F^{\text{AM}}(a),
 \end{aligned}$$

<sup>16</sup> where  $F^{\text{AM}}(a) = (1 - f_a)^3 f_a + (1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2$ ,  $\mathcal{D} = \{0, 1/2, 1\}$ , and  $f_a$  is<sup>16</sup>  
<sup>17</sup> the probability of a minor allele occurring at locus  $a$ .<sup>17</sup>

<sup>18</sup> Using the expected AM attribute diff (Eq. 118), the expected pairwise AM dis-<sup>18</sup>  
<sup>19</sup> tance (Eq. 111) between instances  $i, j \in \mathcal{I}$  is given by<sup>19</sup>

$$\begin{aligned}
 \text{E} (D_{ij}^{\text{AM}}) &= \text{E} \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{AM}}(a) \right) \\
 &= \sum_{a \in \mathcal{A}} \text{E} \left[ d_{ij}^{\text{AM}}(a) \right] \quad (119)^{23} \\
 &= 2 \sum_{a \in \mathcal{A}} F^{\text{AM}}(a),
 \end{aligned}$$

<sup>26</sup> where  $F^{\text{AM}}(a) = (1 - f_a)^3 f_a + (1 - f_a) f_a^3 + (1 - f_a)^2 f_a^2$  and  $f_a$  is the probability<sup>26</sup>  
<sup>27</sup> of a minor allele occurring at locus  $a$ . Similar to GM distances, the expected AM<sup>27</sup>  
<sup>28</sup> distance (Eq. 119) depends only on the minor allele probabilities  $f_a$  for all  $a \in \mathcal{A}$ .<sup>28</sup>

<sup>29</sup> This is to be expected because, although the AM metric is more informative, it still<sup>29</sup>  
<sup>30</sup> only accounts for simple differences between nucleotides of two instances  $i, j \in \mathcal{I}$  at<sup>30</sup>  
<sup>31</sup> some locus  $a$ .<sup>31</sup>

<sup>32</sup>  
<sup>33</sup>

The second moment about the origin for the AM distance is computed as follows

$$\begin{aligned}
 \mathbb{E} \left[ (D_{ij}^{\text{AM}})^2 \right] &= \mathbb{E} \left[ \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{AM}}(a) \right)^2 \right] \\
 &= \mathbb{E} \left[ \sum_{a \in \mathcal{A}} \left( d_{ij}^{\text{AM}}(a) \right)^2 \right] + 2\mathbb{E} \left[ \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} d_{ij}^{\text{AM}}(r) \cdot d_{ij}^{\text{AM}}(s) \right] \\
 &= \sum_{a \in \mathcal{A}} \left( \sum_{k \in \mathcal{D}} k^2 \cdot \mathbb{P} \left[ d_{ij}^{\text{AM}}(a) = k \right] \right) \\
 &\quad + 2 \sum_{a \in \mathcal{A}} \sum_{s \leq r-1} \left( \sum_{k \in \mathcal{D}} k \cdot \mathbb{P} \left[ d_{ij}^{\text{AM}}(r) = k \right] \right) \cdot \left( \sum_{k \in \mathcal{D}} k \cdot \mathbb{P} \left[ d_{ij}^{\text{AM}}(s) = k \right] \right) \\
 &= \sum_{a \in \mathcal{A}} G^{\text{AM}}(a) + 8 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r, s\}} F^{\text{AM}}(\lambda),
 \end{aligned} \tag{120}$$

where  $G^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a) + 2(1 - f_a)^2 f_a^2$ ,  $F^{\text{AM}}(\lambda) = (1 - f_\lambda)^3 f_\lambda + f_\lambda^3 (1 - f_\lambda) + (1 - f_\lambda)^2 f_\lambda^2$ , and  $f_a$  is the probability of a minor allele occurring at locus  $a$ .

Using the first (Eq. 119) and second (Eq. 120) raw moments of the asymptotic AM distance distribution, the variance is given by

$$\begin{aligned}
 \text{Var} (D_{ij}^{\text{AM}}) &= \mathbb{E} \left[ (D_{ij}^{\text{AM}})^2 \right] - [\mathbb{E} (D_{ij}^{\text{AM}})]^2 \\
 &= \sum_{a \in \mathcal{A}} G^{\text{AM}}(a) + 8 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r, s\}} F^{\text{AM}}(\lambda) - 4 \left( \sum_{a \in \mathcal{A}} F^{\text{AM}}(a) \right)^2 \\
 &= \sum_{a \in \mathcal{A}} G^{\text{AM}}(a) - 4 \sum_{a \in \mathcal{A}} [F^{\text{AM}}(a)]^2 \\
 &= \sum_{a \in \mathcal{A}} \left( G^{\text{AM}}(a) - 4 [F^{\text{AM}}(a)]^2 \right),
 \end{aligned} \tag{121}$$

where  $G^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a) + 2(1 - f_a)^2 f_a^2$ ,  $F^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a) + (1 - f_a)^2 f_a^2$ , and  $f_a$  is the probability of a minor allele occurring at locus  $a$ . Similar to the mean (Eq. 119), the variance just depends on minor allele probabilities  $f_a$  for all  $a \in \mathcal{A}$ .

With the mean (Eq. 119) and variance (Eq. 121) estimates of AM distances, the asymptotic AM distance distribution is given by the following


$$D_{ij}^{\text{AM}} \sim \mathcal{N} \left( 2 \sum_{a \in \mathcal{A}} F^{\text{AM}}(a), \sum_{a \in \mathcal{A}} \left( G^{\text{AM}}(a) - 4 [F^{\text{AM}}(a)]^2 \right) \right), \tag{122}$$

<sup>1</sup>where  $G^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a) + 2(1 - f_a)^2 f_a^2$ ,  $F(a) = (1 - f_a)^3 f_a +$   
<sup>2</sup> $f_a^3 (1 - f_a) + (1 - f_a)^2 f_a^2$ , and  $f_a$  is the probability of a minor allele occurring at  
<sup>3</sup>locus  $a$ .

<sup>4</sup> This concludes our analysis of the AM metric in GWAS data when the inde-  
<sup>5</sup>pendence assumption holds for minor allele probabilities  $f_a$  and binomial samples  
<sup>6</sup> $\mathcal{B}(2, f_a)$  for all  $a \in \mathcal{A}$ . In the next section, we derive more complex asymptotic  
<sup>7</sup>results for the TiTv distance metric (Eq. 112).

#### <sup>9</sup>4.3 TiTv distance distribution <sup>9</sup>

<sup>10</sup>The TiTv metric allows for one to account for both genotype mismatch, allele mis-  
<sup>11</sup>match, transition, and transversion. However, this added dimension of information  
<sup>12</sup>requires knowledge of the nucleotide makeup at a particular locus. A sufficient condi-  
<sup>13</sup>tion to compute the TiTv metric between instances  $i, j \in \mathcal{I}$  is that we know whether  
<sup>14</sup>the nucleotides associated with a particular locus  $a$  are both purines (PuPu), purine  
<sup>15</sup>and pyrimidine (PuPy), or both pyrimidines (PyPy). We illustrate all possibilities  
<sup>16</sup>for transitions and transversions in a diagram (Fig. 2). Purines (A and G) and  
<sup>17</sup>pyrimidines (C and T) are shown at the top and bottom, respectively. Transitions  
<sup>18</sup>occur in the cases of PuPu and PyPy, while transversion occurs only with PuPy  
<sup>19</sup>encoding.



<sup>31</sup> This additional encoding is always given in a particular GWAS data set, which  
<sup>32</sup>leads us to consider the probabilities of PuPu, PuPy, and PyPy. These will be  
<sup>33</sup>necessary to determine asymptotics for the TiTv distance metric. Let  $\gamma_0$ ,  $\gamma_1$ , and  $\gamma_2$



<sup>1</sup>denote the probabilities of PuPu, PuPy, and PyPy, respectively, for the  $p$  loci of data<sup>1</sup>  
<sup>2</sup>matrix  $X$ . In real data, there are approximately twice as many transitions as there<sup>2</sup>  
<sup>3</sup>are transversions. That is, the probability of a transition  $P(\text{Ti})$  is approximately<sup>3</sup>  
<sup>4</sup>twice the probability of transversion  $P(\text{Tv})$ . It is likely that any particular data<sup>4</sup>  
<sup>5</sup>set will not satisfy this criterion exactly. In this general case, we have  $P(\text{Ti})$  being<sup>5</sup>  
<sup>6</sup>equal to some multiple  $\eta$  times  $P(\text{Tv})$ . In order to enforce this general constraint in<sup>6</sup>  
<sup>7</sup>simulated data, we define the following set of equalities<sup>7</sup>

$$\gamma_0 + \gamma_1 + \gamma_2 = 1, \quad (123)$$

$$P(\text{Ti}) - \eta P(\text{Tv}) = 0. \quad (124)$$

The sum-to-one constraint (Eq. 123) is natural in this context because there are<sup>11</sup>  
<sup>12</sup>only three possible genotype encodings at a particular locus, which are PuPu, PuPy,<sup>13</sup>  
<sup>14</sup>and PyPy. Solving the Ti/Tv ratio constraint (Eq. 124) for  $\eta$  gives<sup>14</sup>

$$\eta = \frac{P(\text{Ti})}{P(\text{Tv})}, \quad (125)$$

which is easily computed in a real data set by dividing the fraction of Ti out of the<sup>17</sup>  
<sup>18</sup>total  $p$  loci by the fraction of Tv out of the total  $p$  loci. We will use the simplified<sup>18</sup>  
<sup>19</sup>notation  $\eta = \text{Ti}/\text{Tv}$  to represent this factor for the remainder of this work.<sup>19</sup>

Using this PuPu, PuPy, and PyPy encoding, the probability of a transversion<sup>20</sup>  
<sup>21</sup>occurring at any fixed locus  $a$  is given by the following<sup>21</sup>

$$P(\text{Tv}) = \gamma_1. \quad (125)$$

Using the sum-to-one constraint (Eqs. 123) and the probability of<sup>24</sup>  
<sup>25</sup>transversion (Eq. 124), the probability of a transition occurring at locus  $a$  is com-<sup>25</sup>  
<sup>26</sup>puted as follows<sup>26</sup>

$$P(\text{Ti}) = \gamma_0 + \gamma_2. \quad (126)$$

Also using the sum-to-one constraint (Eq. 123) and the Ti/Tv ratio constraint<sup>29</sup>  
<sup>30</sup>(Eq. 124), it is clear that we have  $P(\text{Tv}) = \frac{1}{\eta+1}$  and  $P(\text{Ti}) = \frac{\eta}{\eta+1}$ . Without loss of<sup>30</sup>  
<sup>31</sup>generality, we then sample<sup>31</sup>

$$\gamma_0 \sim \mathcal{U}\left(\varepsilon, \frac{\eta}{\eta+1} - \varepsilon\right), \quad (127)$$

<sup>1</sup>where  $\varepsilon$  is some small positive real number.

<sup>2</sup> Then it immediately follows that we have

$$\gamma_2 = \frac{\eta}{\eta + 1} - \gamma_0. \quad (128)$$

However, we can derive the mean and variance of the distance distribution induced by the TiTv metric without specifying any relationship between  $\gamma_0$ ,  $\gamma_1$ , and  $\gamma_2$ . We proceed by computing  $P[d_{ij}^{\text{TiTv}}(a) = k]$  for each  $k \in \mathcal{D} = \{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}$ . Let  $y$  represent a random sample of size  $p$  from  $\{0, 1, 2\}$ , where

$$y_a = \begin{cases} 0 & \text{if locus } a \text{ is PuPu,} \\ 1 & \text{if locus } a \text{ is PuPy,} \\ 2 & \text{if locus } a \text{ is PyPy.} \end{cases} \quad (129)$$

We derive  $P[d_{ij}^{\text{TiTv}}(a) = 0]$  as follows

$$\begin{aligned} P[d_{ij}^{\text{TiTv}}(a) = 0] &= P[y_a = 0, X_{ia} = X_{ja}] \\ &\quad + P[y_a = 1, X_{ia} = X_{ja}] \\ &\quad + P[y_a = 2, X_{ia} = X_{ja}] \\ &= \gamma_0 [(1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2] \\ &\quad + \gamma_1 [(1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2] \\ &\quad + \gamma_2 [(1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2] \\ &= (\gamma_0 + \gamma_1 + \gamma_2) [(1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2] \\ &= (1 - f_a)^2 + 4f_a(1 - f_a) + f_a^2, \end{aligned} \quad (130)$$

where  $f_a$  is the probability of a minor allele occurring at locus  $a$ .

We derive  $P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{1}{4} \right]$  as follows

$$\begin{aligned}
 P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{1}{4} \right] &= 2P [y_a = 0, X_{ia} = 0, X_{ja} = 1] \\
 &\quad + 2P [y_a = 0, X_{ia} = 1, X_{ja} = 2] \\
 &\quad + 2P [y_a = 2, X_{ia} = 0, X_{ja} = 1] \\
 &\quad + 2P [y_a = 2, X_{ia} = 1, X_{ja} = 2] \\
 &= 4\gamma_0(1-f_a)^3 f_a + 4\gamma_0 f_a^3(1-f_a) + 4\gamma_2(1-f_a)^3 f_a \\
 &\quad + 4\gamma_2 f_a^3(1-f_a) \\
 &= 4\gamma_0 [(1-f_a)^3 f_a + f_a^3(1-f_a)] \\
 &\quad + 4\gamma_2 [(1-f_a)^3 f_a + f_a^3(1-f_a)] \\
 &= 4(\gamma_0 + \gamma_2) [(1-f_a)^3 f_a + f_a^3(1-f_a)],
 \end{aligned} \tag{131}$$

where  $f_a$  is the probability of a minor allele occurring at locus  $a$ ,  $\gamma_0$  is the probability of PuPu occurring at any locus  $a$ , and  $\gamma_2$  is the probability of PyPy occurring at any locus  $a$ .

We derive  $P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{1}{2} \right]$  as follows

$$\begin{aligned}
 P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{1}{2} \right] &= 2P [y_a = 1, X_{ia} = 0, X_{ja} = 1] \\
 &\quad + 2P [y_a = 1, X_{ia} = 1, X_{ja} = 2] \\
 &= 4\gamma_1(1-f_a)^3 f_a + 4\gamma_1 f_a^3(1-f_a) \\
 &= 4\gamma_1 [(1-f_a)^3 f_a + f_a^3(1-f_a)],
 \end{aligned} \tag{132}$$

where  $f_a$  is the probability of a minor allele occurring at locus  $a$  and  $\gamma_1$  is the probability of PuPy occurring at any locus  $a$ .

We derive  $P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{3}{4} \right]$  as follows

$$\begin{aligned}
 P \left[ d_{ij}^{\text{TiTv}}(a) = \frac{3}{4} \right] &= 2P [y_a = 0, X_{ia} = 0, X_{ja} = 2] \\
 &\quad + 2P [y_a = 2, X_{ia} = 0, X_{ja} = 2] \\
 &= 2\gamma_0(1-f_a)^2 f_a^2 + 2\gamma_2(1-f_a)^2 f_a^2 \\
 &= 2(\gamma_0 + \gamma_2)(1-f_a)^2 f_a^2,
 \end{aligned} \tag{133}$$

<sup>1</sup>where  $f_a$  is the probability of a minor allele occurring at locus  $a$ ,  $\gamma_0$  is the probability<sup>1</sup>  
<sup>2</sup>of PuPu occurring at any locus  $a$ , and  $\gamma_2$  is the probability of PyPy occurring at<sup>2</sup>  
<sup>3</sup>any locus  $a$ .<sup>3</sup>

<sup>4</sup> We derive  $P[d_{ij}^{\text{TiTv}}(a) = 1]$  as follows<sup>4</sup>

$$\begin{aligned} \text{P}[d_{ij}^{\text{TiTv}}(a) = 1] &= 2\text{P}[y_a = 1, X_{ia} = 0, X_{ja} = 2] \\ &= 2\gamma_1(1 - f_a)^2 f_a^2, \end{aligned} \quad (134)$$

<sup>8</sup> where  $f_a$  is the probability of a minor allele occurring at locus  $a$  and  $\gamma_1$  is the<sup>8</sup>  
<sup>9</sup>probability of PuPy occurring at any locus  $a$ .<sup>9</sup>

<sup>10</sup> Using the TiTv diff probabilities (Eqs. 130-134), we compute the expected TiTv<sup>10</sup>  
<sup>11</sup>distance between instances  $i, j \in \mathcal{I}$  as follows<sup>11</sup>

$$\begin{aligned} \text{E}(D_{ij}^{\text{TiTv}}) &= \sum_{a \in \mathcal{A}} \left( \sum_{k \in \mathcal{D}} k \cdot \text{P}[d_{ij}^{\text{TiTv}}(a) = k] \right) \\ &= (\gamma_0 + \gamma_2 + 2\gamma_1) \sum_{a \in \mathcal{A}} [(1 - f_a)^3 f_a + f_a^3 (1 - f_a)] \\ &\quad + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} (1 - f_a)^2 f_a^2 \\ &= (\gamma_0 + \gamma_2 + 2\gamma_1) \sum_{a \in \mathcal{A}} F^{\text{TiTv}}(a) \\ &\quad + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G^{\text{TiTv}}(a), \end{aligned} \quad (135)$$

<sup>21</sup> where  $F^{\text{TiTv}}(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a)$ ,  $G^{\text{TiTv}}(a) = (1 - f_a)^2 f_a^2$ ,  $f_a$  is the<sup>21</sup>  
<sup>22</sup>probability of a minor allele occurring at locus  $a$ ,  $\gamma_0$  is the probability of PuPu<sup>22</sup>  
<sup>23</sup>occurring at any locus  $a$ ,  $\gamma_1$  is the probability of PuPy occurring at any locus  $a$ ,<sup>23</sup>  
<sup>24</sup>and  $\gamma_2$  is the probability of PyPy occurring at any locus  $a$ . In contrast to the<sup>24</sup>  
<sup>25</sup>expected GM and AM distances (Eqs. 114 and 119), the expected TiTv distance<sup>25</sup>  
<sup>26</sup>(Eq. 135) depends on minor allele probabilities  $f_a$  for all  $a \in \mathcal{A}$  and the genotype<sup>26</sup>  
<sup>27</sup>encoding probabilities  $\gamma_0, \gamma_1$ , and  $\gamma_2$ .<sup>27</sup>

<sup>1</sup> The second moment about the origin for the TiTv distance is computed as follows<sup>1</sup>

$$\begin{aligned}
 & \text{E} \left[ \left( D_{ij}^{\text{TiTv}} \right)^2 \right] = \text{E} \left[ \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{TiTv}}(a) \right)^2 \right] \\
 & = \text{E} \left[ \sum_{a \in \mathcal{A}} \left( d_{ij}^{\text{TiTv}}(a) \right)^2 \right] + 2 \text{E} \left[ \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} d_{ij}^{\text{TiTv}}(r) \cdot d_{ij}^{\text{TiTv}}(s) \right] \\
 & = \sum_{a \in \mathcal{A}} \left( \sum_{k \in \mathcal{D}} k^2 \cdot \text{P} \left[ d_{ij}^{\text{TiTv}}(a) = k \right] \right) \\
 & + 2 \sum_{a \in \mathcal{A}} \sum_{s \leq r-1} \left( \sum_{k \in \mathcal{D}} k \cdot \text{P} \left[ d_{ij}^{\text{TiTv}}(r) = k \right] \right) \cdot \left( \sum_{k \in \mathcal{D}} k \cdot \text{P} \left[ d_{ij}^{\text{TiTv}}(s) = k \right] \right) \quad (136) \\
 & = \left[ \frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1 \right] \sum_{a \in \mathcal{A}} F^{\text{TiTv}}(a) + \left[ \frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G^{\text{TiTv}}(a) \\
 & + 2 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r, s\}} \left\{ \left( [\gamma_0 + \gamma_2 + 2\gamma_1] F^{\text{TiTv}}(\lambda) \right. \right. \\
 & \quad \left. \left. + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] G^{\text{TiTv}}(\lambda) \right) \right\},
 \end{aligned}$$

where  $F^{\text{TiTv}}(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a)$ ,  $G^{\text{TiTv}}(a) = (1 - f_a)^2 f_a^2$ ,  $f_a$  is the probability of a minor allele occurring at locus  $a$ ,  $\gamma_0$  is the probability of PuPu occurring at any locus  $a$ ,  $\gamma_1$  is the probability of PuPy occurring at any locus  $a$ , and  $\gamma_2$  is the probability of PyPy occurring at any locus  $a$ .


<sup>1</sup> Using the first (Eq. 135) and second (Eq. 136) raw moments of the TiTv distance,  
<sup>2</sup>the variance is given by

$$\begin{aligned}
 \text{Var}(D_{ij}^{\text{TiTv}}) &= \text{E} \left[ (D_{ij}^{\text{TiTv}})^2 \right] - [\text{E}(D_{ij}^{\text{TiTv}})]^2 \\
 &= \left[ \frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1 \right] \sum_{a \in \mathcal{A}} F^{\text{TiTv}}(a) \\
 &\quad + \left[ \frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G^{\text{TiTv}}(a) \\
 &\quad + 2 \sum_{r \in \mathcal{A}} \sum_{s \leq r-1} \prod_{\lambda \in \{r,s\}} \left\{ \left( [\gamma_0 + \gamma_2 + 2\gamma_1] F^{\text{TiTv}}(\lambda) \right. \right. \\
 &\quad \left. \left. + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] G^{\text{TiTv}}(\lambda) \right) \right\} \quad (137) \\
 &\quad - \left\{ \left( [\gamma_0 + \gamma_2 + 2\gamma_1] \sum_{a \in \mathcal{A}} F^{\text{TiTv}}(a) \right. \right. \\
 &\quad \left. \left. + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G^{\text{TiTv}}(a) \right) \right\}^2 \\
 &= \left[ \frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1 \right] \sum_{a \in \mathcal{A}} F^{\text{TiTv}}(a) + \left[ \frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] \sum_{a \in \mathcal{A}} G^{\text{TiTv}}(a) \\
 &\quad - \sum_{a \in \mathcal{A}} \left( [\gamma_0 + \gamma_2 + 2\gamma_1] F^{\text{TiTv}}(a) + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right] G^{\text{TiTv}}(a) \right)^2,
 \end{aligned}$$

where  $F^{\text{TiTv}}(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a)$ ,  $G^{\text{TiTv}}(a) = (1 - f_a)^2 f_a^2$ ,  $f_a$  is the  
probability of a minor allele occurring at locus  $a$ ,  $\gamma_0$  is the probability of PuPu  
occurring at any locus  $a$ ,  $\gamma_1$  is the probability of PuPy occurring at any locus  $a$ ,  
and  $\gamma_2$  is the probability of PyPy occurring at any locus  $a$ .




<sup>1</sup>slightly with increased  $\eta = \text{Ti}/\text{Tv}$ . As  $\eta \rightarrow 0^+$ , the data is approaching all Tv and<sup>1</sup>  
<sup>2</sup>no Ti, which means the TiTv distance is larger by definition. On the other hand, the<sup>2</sup>  
<sup>3</sup>TiTv distance decreases as  $\eta \rightarrow 2^-$  because the data is approaching approximately<sup>3</sup>  
<sup>4</sup>twice as many Ti as there are Tv, which is typical for GWAS data in humans.<sup>4</sup>



We also compared theoretical and sample moments as a function of  $\eta = \text{Ti}/\text{Tv}$  and  $\bar{f}_a$  for the TiTv distance metric (Fig. 4B and D). We fixed  $\bar{f}_a$  and computed the theoretical and simulated moments as a function of  $\eta$  (Fig. 4B). Theoretical average TiTv distance, given by Eq. 135, and simulated TiTv average distance are approximately equal as  $\eta$  increases. Theoretical standard deviation, given by Eq. 137, and simulated TiTv standard deviation differ slightly. We also fixed  $\eta$  and computed theoretical and sample moments as a function of  $\bar{f}_a$  (Fig. 4D). In this case, there is approximate agreement with simulated and theoretical moments as  $\bar{f}_a$  increases.

We summarize our moment estimates for GWAS distance metrics (Eqs. 110-112) (Table 4) organized by metric, statistic (mean or variance), and asymptotic formula. Next we consider the important case of distributions of GWAS distances projected onto a single attribute (Eqs. 107-109).





**Figure 4** Density curves and moments of TiTv distance as a function of average MAF  $\bar{f}_a$ , given by Eq. 139, and Ti/Tv ratio  $\eta$ , given by Eq. 124. We fix  $m = p = 100$  for all simulated TiTv distances. **(A)** For fixed  $\bar{f}_a = 0.055$ , TiTv distance density is plotted as a function of increasing  $\eta$ . TiTv distance decreases as  $\eta$  increases. For  $\eta = \text{Ti/Tv} = 0.5$ , there are twice as many transversions as there are transitions. On the other hand,  $\eta = \text{Ti/Tv} = 2$  indicates that there are half as many transversions as transitions. Since transversions encode a larger magnitude distance than transitions, this behavior is expected. **(B)** Simulated and predicted mean  $\pm$  SD are shown as a function of increasing Ti/Tv ratio  $\eta$ . Distance decreases as Ti/Tv increases. Theoretical and simulated moments are approximately the same. **(C)** For fixed  $\eta = 2$ , TiTv distance density is plotted as a function of increasing  $\bar{f}_a$ . TiTv distance increases as  $\bar{f}_a$  approaches maximum of 0.5, which means that there is about the same frequency of minor alleles as major alleles. **(D)** Simulated and predicted mean  $\pm$  SD as a function of increasing average MAF  $\bar{f}_a$ . Distance increases as the number of minor alleles increases. Theoretical and simulated moments are approximately the same.

#### 4.4 Distribution of one-dimensional projected GWAS distance onto a SNP

We previously derived the exact distribution of the one-dimensional projected distance onto an attribute in continuous data (Section 3.2.3), which is used as the predictor in NPDR to calculate relative attribute importance in the form of standardized beta coefficients. GWAS data and the metrics we have considered are dis-

<sup>1</sup>crete. Therefore, we derive the density function for each diff metric (Eqs. 107-109),<sup>1</sup>

<sup>2</sup>which also serves as the probability distribution for each metric, respectively.<sup>2</sup>

<sup>3</sup> The support of the GM metric (Eq. 107) is simply  $\{0, 1\}$ , so we derive the proba-<sup>3</sup>

<sup>4</sup>bility,  $P[d_{ij}^{GM}(a) = k]$ , of this diff taking on each of these two possible values. First,<sup>4</sup>

<sup>5</sup>the probability that the GM diff is equal to zero is given by<sup>5</sup>

$$f_{GM}(0; f_a) = P[d_{ij}^{GM}(a) = 0]$$

$$= P(X_{ia} = 0, X_{ja} = 0) + P(X_{ia} = 1, X_{ja} = 1)$$

$$+ P(X_{ia} = 2, X_{ja} = 2)$$

$$= (1 - f_a)^4 + 4f_a^2(1 - f_a)^2 + f_a^4,$$

<sup>11</sup>where  $f_a$  is the probability of a minor allele occurring at locus  $a$ .<sup>11</sup>

<sup>12</sup>Similarly, the probability that the GM diff is equal to 1 is derived as follows<sup>12</sup>

$$f_{GM}(1; f_a) = P[d_{ij}^{GM}(a) = 1]$$

$$= 2P(X_{ia} = 0, X_{ja} = 1) + 2P(X_{ia} = 1, X_{ja} = 2)$$

$$+ 2P(X_{ia} = 0, X_{ja} = 2)$$

$$= 4(1 - f_a)^3 f_a + 4f_a^3(1 - f_a) + 2f_a^2(1 - f_a)^2,$$

<sup>19</sup>where  $f_a$  is the probability of a minor allele occurring at locus  $a$ .<sup>19</sup>

<sup>20</sup> This leads us to the probability distribution of the GM diff metric, which is the<sup>20</sup>

<sup>21</sup>distribution of the one-dimensional GM distance projected onto a single SNP. This<sup>21</sup>

<sup>22</sup>distribution is given by<sup>22</sup>

$$f_{GM}(d; f_a) = \begin{cases} (1 - f_a)^4 + 4f_a^2(1 - f_a)^2 + f_a^4 & d = 0, \\ 4(1 - f_a)^3 f_a + 4f_a^3(1 - f_a) + 2f_a^2(1 - f_a)^2 & d = 1, \end{cases} \quad (142)$$

<sup>26</sup>where  $f_a$  is the probability of a minor allele occurring at locus  $a$ .<sup>26</sup>

<sup>27</sup> The mean and variance of this GM diff distribution can easily be derived using<sup>27</sup>

<sup>28</sup>this newly determined density function (Eq. 142). The average GM diff is given by<sup>28</sup>

<sup>29</sup>the following<sup>29</sup>

$$E[d_{ij}^{GM}(a)] = 2F^{GM}(a), \quad (143)$$

<sup>32</sup>where  $F^{GM} = 2(1 - f_a)^3 f_a + 2f_a^3(1 - f_a) + f_a^2(1 - f_a)^2$  and  $f_a$  is the probability<sup>32</sup>

<sup>33</sup>of a minor allele occurring at locus  $a$ .<sup>33</sup>

The variance of the GM diff metric is given by

$$\text{Var} \left[ d_{ij}^{\text{GM}}(a) \right] = 2F^{\text{GM}}(a) \left[ 1 - 2F^{\text{GM}}(a) \right], \quad (144)$$

where  $F^{\text{GM}} = 2(1 - f_a)^3 f_a + 2f_a^3(1 - f_a) + f_a^2(1 - f_a)^2$  and  $f_a$  is the probability of a minor allele occurring at locus  $a$ .

The support of the AM metric (Eq. 108) is  $\{0, 1/2, 1\}$ . Beginning with the probability of the AM diff being equal to 0, we have the following probability

$$\begin{aligned} f_{\text{AM}}(0; f_a) &= \text{P} \left[ d_{ij}^{\text{AM}}(a) = 0 \right] \\ &= \text{P}(X_{ia} = 0, X_{ja} = 0) + \text{P}(X_{ia} = 1, X_{ja} = 1) \\ &\quad + \text{P}(X_{ia} = 2, X_{ja} = 2) \\ &= (1 - f_a)^4 + 4f_a^2(1 - f_a)^2 + f_a^4, \end{aligned} \quad (145)$$

where  $f_a$  is the probability of a minor allele occurring at locus  $a$ .

The probability of the AM diff metric being equal to 1/2 is computed similarly as follows

$$\begin{aligned} f_{\text{AM}}(1/2; f_a) &= \text{P} \left[ d_{ij}^{\text{AM}}(a) = 1/2 \right] \\ &= 2\text{P}(X_{ia} = 0, X_{ja} = 1) + 2\text{P}(X_{ia} = 1, X_{ja} = 2) \\ &= 4(1 - f_a)^3 f_a + 4f_a^3(1 - f_a), \end{aligned} \quad (146)$$

where  $f_a$  the probability of a minor allele occurring at locus  $a$ .

Finally, the probability of the AM diff metric being equal to 1 is given by the following

$$\begin{aligned} f_{\text{AM}}(1; f_a) &= \text{P} \left[ d_{ij}^{\text{AM}}(a) = 1 \right] = 2\text{P}(X_{ia} = 0, X_{ja} = 2) \\ &= 2f_a^2(1 - f_a)^2, \end{aligned} \quad (147)$$

where  $f_a$  is the probability of a minor allele occurring at locus  $a$ .

As in the case of the GM diff metric, we now have the probability distribution of the AM diff metric. This also serves as the distribution of the one-dimensional AM

<sup>1</sup>distance projected onto a single SNP, and is given by the following

$$f_{\text{AM}}(d; f_a) = \begin{cases} (1 - f_a)^4 + 4f_a^2(1 - f_a)^2 + f_a^4 & d = 0, \\ 4(1 - f_a)^3 f_a + 4f_a^3(1 - f_a) & d = 1/2, \\ 2f_a^2(1 - f_a)^2 & d = 1, \end{cases} \quad (148)$$

<sup>7</sup>where  $f_a$  is the probability of a minor allele occurring at locus  $a$ .

<sup>8</sup>The mean and variance of this AM diff distribution is derived using the corre-  
<sup>9</sup>sponding density function (Eq. 148). The average AM diff is given by

$$\text{E} [d_{ij}^{\text{AM}}(a)] = 2F^{\text{AM}}(a), \quad (149)$$

<sup>12</sup>where  $F^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a) + f_a^2(1 - f_a)^2$  and  $f_a$  is the probability  
<sup>13</sup>of a minor allele occurring at locus  $a$ .

<sup>14</sup>The variance of the AM diff metric is given by

$$\text{Var} [d_{ij}^{\text{AM}}(a)] = G^{\text{AM}}(a) - 4 [F^{\text{AM}}(a)]^2, \quad (150)$$

<sup>17</sup>where  $G^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a) + 2(1 - f_a)^2 f_a^2$ ,  
<sup>18</sup> $F^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a) + f_a^2(1 - f_a)$ ,  $f_a$  is the probability of a minor  
<sup>19</sup>allele occurring at locus  $a$ .

<sup>20</sup>For the TiTv diff metric (Eq. 109), the support is  $\{0, 1/4, 1/2, 3/4, 1\}$ . We have  
<sup>21</sup>already derived the probability that the TiTv diff assumes each of the values of its  
<sup>22</sup>support (Eqs. 130-134). Therefore, we have the following distribution of the TiTv  
<sup>23</sup>diff metric

$$f_{\text{TiTv}}(d; f_a, \gamma_0, \gamma_1, \gamma_2) = \begin{cases} (1 - f_a)^4 + 4f_a^2(1 - f_a)^2 + f_a^4 & d = 0, \\ 4(\gamma_0 + \gamma_2) [(1 - f_a)^3 f_a + f_a^3(1 - f_a)] & d = 1/4, \\ 4\gamma_1 [(1 - f_a)^3 f_a + f_a^3(1 - f_a)] & d = 1/2, \\ 2(\gamma_0 + \gamma_2)(1 - f_a)^2 f_a^2 & d = 3/4, \\ 2\gamma_1(1 - f_a)^2 f_a^2 & d = 1, \end{cases} \quad (151)$$

<sup>31</sup>where  $f_a$  is the probability of a minor allele occurring at locus  $a$ ,  $\gamma_0$  is the probability  
<sup>32</sup>of PuPu at locus  $a$ ,  $\gamma_1$  is the probability of PuPy at locus  $a$ ,  $\gamma_2$  is the probability  
<sup>33</sup>of PyPy at locus  $a$ , and  $\eta$  is the Ti/Tv ratio (Eq. 124).

The mean and variance of this TiTv diff distribution is derived using the corresponding density function (Eq. 151). The average TiTv diff is given by

$$E[d_{ij}^{\text{TiTv}}(a)] = (\gamma_0 + \gamma_2 + 2\gamma_1)F^{\text{TiTv}}(a) + \left[\frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1\right]G^{\text{TiTv}}(a), \quad (152)$$

where  $F^{\text{TiTv}}(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a)$ ,  $G^{\text{TiTv}}(a) = f_a^2 (1 - f_a)^2$ ,  $f_a$  is the probability of a minor allele occurring at locus  $a$ ,  $\gamma_0$  is the probability of PuPu at locus  $a$ ,  $\gamma_1$  is the probability of PuPy at locus  $a$ , and  $\gamma_2$  is the probability of PyPy at locus  $a$ .

The variance of the TiTv diff metric is given by

$$\begin{aligned} \text{Var}[d_{ij}^{\text{TiTv}}(a)] = & \left[\frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1\right]F^{\text{TiTv}}(a) \\ & + \left[\frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1\right]G^{\text{TiTv}}(a) \\ & - \left\{ \left( (\gamma_0 + \gamma_2 + 2\gamma_1)F^{\text{TiTv}}(a) \right. \right. \\ & \left. \left. + \left[ \frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1 \right]G^{\text{TiTv}}(a) \right) \right\}^2, \end{aligned} \quad (153)$$

where  $F^{\text{TiTv}}(a) = (1 - f_a)^3 f_a + f_a^3 (1 - f_a)$ ,  $G^{\text{TiTv}}(a) = f_a^2 (1 - f_a)^2$ ,  $f_a$  is the probability of a minor allele occurring at locus  $a$ ,  $\gamma_0$  is the probability of PuPu at locus  $a$ ,  $\gamma_1$  is the probability of PuPy at locus  $a$ , and  $\gamma_2$  is the probability of PyPy at locus  $a$ .

These novel distribution results for the projection of pairwise GWAS distances onto a single genetic variant, as well as results for the full space of  $p$  variants, can inform NPDR and other nearest-neighbor distance-based feature selection algorithms. Next we introduce our new diff metric and distribution results for time series derived correlation-based data, with a particular application to resting-state fMRI.

## 5 Time series correlation-based distance distribution

For time series correlation-based data, we consider the case where there are  $m$  correlation matrices  $A^{(p \times p)}$  (one matrix for each subject). In particular, we have in mind the application of resting-state fMRI (rs-fMRI) data. The derivations that follow, however, are relevant to all correlation-based data with the assumptions we note. The attributes in rs-fMRI are commonly Regions of Interest (ROIs), which

<sup>1</sup>are collections of spatially proximal voxels [22]. Correlation in their time-series<sup>1</sup>  
<sup>2</sup>activity arise between different ROIs at the voxel level or for a given brain atlas<sup>2</sup>  
<sup>3</sup>[23]. Because the attributes of interest ( $a$ ) are the ROIs themselves, we propose the<sup>3</sup>  
<sup>4</sup>following attribute projection (diff) 4

$$d_{ij}^{\text{ROI}}(a) = \sum_{k \neq a} |A_{ak}^{(i)} - A_{ak}^{(j)}|, \quad (154)$$

where  $A_{ak}^{(i)}$  and  $A_{ak}^{(j)}$  are the correlations between ROI  $a$  and ROI  $k$  for instances<sup>5</sup>  
 $i, j \in \mathcal{I}$ , respectively. With this rs-fMRI diff, we define the pairwise distance between<sup>6</sup>  
two instances  $i, j \in \mathcal{I}$  as follows 10

$$D_{ij}^{\text{fMRI}} = \sum_{a \in \mathcal{A}} d_{ij}^{\text{ROI}}(a), \quad (155)$$

which is based on Manhattan ( $q = 1$ ). This metric may be expanded to general  $q$ ,  
but we only consider  $q = 1$ .

In order for comparisons between different correlations to be possible, we first  
perform a Fisher r-to-z transform on the correlations. This transformation makes the  
data approximately normally distributed with stabilized variance across different  
samples. After this transformation, we then load all of the transformed correlations  
into a  $p(p-1) \times m$  matrix  $X$  (Fig. 5). Each column of  $X$  represents a single instance  
(or subject) in rs-fMRI data. Contrary to a typical  $p \times m$  data set, each row does  
not represent a single attribute. Rather, each attribute (or ROI) is represented by  
 $p-1$  consecutive rows. The first  $p-1$  rows represent ROI<sub>1</sub>, the next  $p-1$  rows  
represent ROI<sub>2</sub>, and so on until the last  $p-1$  rows that represent ROI <sub>$p$</sub> . For a  
given column of  $X$ , we exclude pairwise correlations between an ROI and itself.  
Therefore, the matrix does not contain  $\hat{A}_{aa}^{(i)}$  for any  $i \in \mathcal{I}$  or  $a \in \mathcal{A}$ . Furthermore,  
symmetry of correlation matrices means that each column contains exactly two of  
each element of the upper triangle of an instance's transformed correlation matrix.  
For example,  $\hat{A}_{ka}^{(i)} = \hat{A}_{ak}^{(i)}$  for  $k \neq a$  and both will be contained in a given column  
of  $X$  for each  $a \in \mathcal{A}$ . Based on our rs-fMRI diff (Eq. 154), the organization of  $X$   
makes computation of each value of the diff very simple. In order to compute each  
value of the rs-fMRI diff, we just need to know the starting and ending row indices

<sup>1</sup>for a given ROI. Starting indices are given by

<sup>2</sup>


$$\text{start}_k = (k-1)(p-1) + 1, \quad \text{for } k = 1, 2, \dots, p$$

<sup>4</sup>

<sup>5</sup>and ending indices are given by

$$\text{end}_k = k(p-1), \quad \text{for } k = 1, 2, \dots, p.$$

<sup>8</sup> These indices allow us to extract just the rows necessary to compute the rs-fMRI<sup>8</sup>  
<sup>9</sup>diff for a fixed ROI.



<sup>24</sup> We further transform the data matrix  $X$  by standardizing so that each of the<sup>24</sup>  
<sup>25</sup> $m$  columns has zero mean and unit variance. Therefore, the data in matrix  $X$ <sup>25</sup>  
<sup>26</sup>are approximately standard normal. Recall that the mean (Eq. 38) and variance<sup>26</sup>  
<sup>27</sup>(Eq. 39) of the Manhattan ( $L_1$ ) distance distribution for standard normal data are<sup>27</sup>  
<sup>28</sup> $\frac{2p}{\sqrt{\pi}}$  and  $\frac{2(\pi-2)p}{\pi}$ , respectively. This allows us to easily derive the expected pairwise<sup>28</sup>

<sup>1</sup>distance between instances  $i, j \in \mathcal{I}$  in rs-fMRI data as follows

$$\begin{aligned}
 \mathbb{E}(D_{ij}^{\text{fMRI}}) &= \mathbb{E} \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{ROI}}(a) \right) \\
 &= \mathbb{E} \left( \sum_{a \in \mathcal{A}} \sum_{k \neq a} \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right| \right) \\
 &= \sum_{a \in \mathcal{A}} \sum_{k \neq a} \mathbb{E} \left( \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right| \right) \\
 &= \sum_{a \in \mathcal{A}} \sum_{k \neq a} \frac{2}{\sqrt{\pi}} \\
 &= \frac{2p(p-1)}{\sqrt{\pi}}.
 \end{aligned} \tag{156}$$

<sup>11</sup>The expected pairwise rs-fMRI distance (Eq. 156) grows on the order of  $p(p-1)$ ,  
<sup>12</sup>which is the total number of transformed pairwise correlations in each column of  $X$   
<sup>13</sup>(Fig. 5). This is similar to the case of a typical  $m \times p$  data matrix in which the data  
<sup>14</sup>is standard normal and Manhattan distances are computed between instances.

<sup>15</sup>We first derive the variance of the rs-fMRI distance by making an independence  
<sup>16</sup>assumption with respect to the magnitude differences  $|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|$  for all  $k \neq a \in \mathcal{A}$ .  
<sup>17</sup>We observe empirically that this assumption gives a reasonable estimate of the  
<sup>18</sup>actual variance of rs-fMRI distances in simulated data, but there is a consistent  
<sup>19</sup>discrepancy between predicted and simulated variances. We begin our derivation  
<sup>20</sup>of the variance of rs-fMRI distances by assuming that cross-covariances between  
<sup>21</sup>the diffs of different pairs of ROIs are negligible. This allows us to determine the  
<sup>22</sup>relationship between the predicted variance under the independence assumption  
<sup>23</sup>and the simulated variance. We proceed by applying the variance operator linearly  
<sup>24</sup>as follows

$$\begin{aligned}
 \text{Var}(D_{ij}^{\text{fMRI}}) &= \text{Var} \left( \sum_{a \in \mathcal{A}} d_{ij}^{\text{ROI}}(a) \right) \\
 &= \text{Var} \left( \sum_{a \in \mathcal{A}} \sum_{k \neq a} \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right| \right) \\
 &= \sum_{a \in \mathcal{A}} \sum_{k \neq a} \text{Var} \left( \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right| \right) \\
 &= \sum_{a \in \mathcal{A}} \sum_{k \neq a} \frac{2(\pi-2)}{\pi} \\
 &= \frac{2(\pi-2)(p-1)p}{\pi}.
 \end{aligned} \tag{157}$$



<sup>1</sup> Similar to the case of an  $m \times p$  data matrix containing standard normal data,<sup>1</sup>  
<sup>2</sup> we have an rs-fMRI distance variance that grows on the order of  $p(p-1)$ , which is<sup>2</sup>  
<sup>3</sup> the total number of pairwise associations in a column of data matrix  $X$  (Fig. 5).<sup>3</sup>  
<sup>4</sup> Therefore, the expected rs-fMRI distance (Eq. 156) and the variance of the rs-fMRI<sup>4</sup>  
<sup>5</sup> distance (Eq. 157) increase on the same order.<sup>5</sup>  
<sup>6</sup> The independence assumption used to derive the variance of our rs-fMRI distance<sup>6</sup>  
<sup>7</sup> metric (Eq. 157) is not satisfied because a single value of the diff (Eq. 154) includes<sup>7</sup>  
<sup>8</sup> the same fixed ROI,  $a$ , for each term in the sum for all  $k \neq a$ . Therefore, the linear<sup>8</sup>  
<sup>9</sup> application of the variance operator we have previously employed does not account<sup>9</sup>  
<sup>10</sup> for the additional cross-covariance that exists. However, we have seen empirically<sup>10</sup>  
<sup>11</sup> that the theoretical variance of the distance we computed for the rs-fMRI distance<sup>11</sup>  
<sup>12</sup> metric (Eq. 157) still reasonably approximates the sample variance, there is a slight<sup>12</sup>  
<sup>13</sup> discrepancy between our theoretical rs-fMRI distance metric variance (Eq. 157)<sup>13</sup>  
<sup>14</sup> and the sample variance. More precisely, the formula we have given for the variance<sup>14</sup>  
<sup>15</sup> (Eq. 157) consistently underestimates the sample variance of the rs-fMRI distance.<sup>15</sup>  
<sup>16</sup> To adjust for this discrepancy, we determine a corrected formula by assuming that<sup>16</sup>  
<sup>17</sup> there is dependence between the terms of the rs-fMRI diff and estimate the cross-<sup>17</sup>  
<sup>18</sup> covariance between rs-fMRI diffs of different pairs of ROIs.<sup>18</sup>  
<sup>19</sup> We begin the derivation of our corrected formula by writing the variance as a two-<sup>19</sup>  
<sup>20</sup> part sum, where the first term in the sum involves the variance of the magnitude<sup>20</sup>  
<sup>21</sup> difference  $|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|$  and then second term involves the cross-covariance of the rs-<sup>21</sup>  
<sup>22</sup> fMRI diff for distinct pairwise ROI-ROI associations. This formulation is implied in<sup>22</sup>  
<sup>23</sup> our previous derivation of the variance, but our independence assumption allowed<sup>23</sup>  
<sup>24</sup> us to assume that all terms in the second part of the two-part sum were zero. Our<sup>24</sup>  
<sup>25</sup>  
<sup>26</sup>  
<sup>27</sup>  
<sup>28</sup>  
<sup>29</sup>  
<sup>30</sup>  
<sup>31</sup>  
<sup>32</sup>  
<sup>33</sup>

<sup>1</sup>formulation of the variance is given by the following

$$\begin{aligned}
 \text{Var}(D_{ij}^{\text{fMRI}}) &= \text{Var} \left( \sum_{a \in \mathcal{A}} \sum_{k \neq a} \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right| \right) \\
 &= \sum_{a=1}^{p-1} \text{Var} \left( \sum_{k=a+1}^p 2 \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right| \right) \\
 &\quad + 2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2 \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right|, \sum_{s=r+1}^p 2 \left| \hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)} \right| \right) \\
 &= \sum_{a=1}^{p-1} \sum_{k=a+1}^p \text{Var} \left( 2 \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right| \right) \\
 &\quad + 2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2 \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right|, \sum_{s=r+1}^p 2 \left| \hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)} \right| \right) \quad (158) \\
 &= \sum_{a=1}^{p-1} \sum_{k=a+1}^{p-1} \frac{4(\pi-2)}{\pi} \\
 &\quad + 2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2 \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right|, \sum_{s=r+1}^p 2 \left| \hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)} \right| \right) \\
 &= \frac{2p(\pi-2)(p-1)}{\pi} \\
 &\quad + 2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2 \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right|, \sum_{s=r+1}^p 2 \left| \hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)} \right| \right).
 \end{aligned}$$

In order to have a formula in terms of the number of ROIs  $p$  only, we estimate the double sum on the right-hand side of the equation of rs-fMRI distance variance (Eq. 158). Through simulation, it can be seen that the difference between sample variance  $S_{D_{ij}}^2$  and  $\frac{2p(\pi-2)(p-1)}{\pi}$  has a quadratic relationship with  $p$ . More explicitly, we have the following relationship

$$S_{D_{ij}}^2 - \frac{2p(\pi-2)(p-1)}{\pi} = \beta_1 p^2 + \beta_0 p. \quad (159)$$

where  $\beta_0$  and  $\beta_1$  are the coefficients we must estimate in order to approximate the cross-covariance term on the right-hand side of the rs-fMRI distance variance equation (Eq. 158).

The coefficient estimates found through least squares fitting are  $\beta_1 = -\beta_0 \approx 0.08$ .

These estimates allow us to arrive at a functional form for the double sum on the right-hand side of the rs-fMRI distance variance equation (Eq. 158) that is proportional to  $\frac{2p(\pi-2)(p-1)}{\pi}$ . That is, we have the following formula for approximating the

$$\begin{aligned}
& \text{double sum} \\
& 2 \sum_{a=1}^{p-1} \sum_{r=a+1}^{p-1} \text{Cov} \left( \sum_{k=a+1}^p 2 \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right|, \sum_{s=r+1}^p 2 \left| \hat{A}_{rs}^{(i)} - \hat{A}_{rs}^{(j)} \right| \right) \\
& \approx \frac{p(\pi - 2)(p - 1)}{4\pi}.
\end{aligned} \tag{160}$$

Therefore, the variance of the rs-fMRI distances is approximated well by the following

$$\text{Var}(D_{ij}^{\text{fMRI}}) \approx \frac{9p(\pi - 2)(p - 1)}{4\pi}. \tag{161}$$

With the mean (Eq. 156) and variance (Eq. 161) estimates, we have the following asymptotic distribution for rs-fMRI distances

$$D_{ij}^{\text{fMRI}} \sim \mathcal{N} \left( \frac{2p(p - 1)}{\sqrt{\pi}}, \frac{9p(\pi - 2)(p - 1)}{4\pi} \right). \tag{162}$$

### 5.1 Max-min normalized time series correlation-based distance distribution

Previously (Section 3.4) we determined the asymptotic distribution of the sample maximum of size  $m$  from a standard normal distribution. We can naturally extend these results to our transformed rs-fMRI data because  $X$  (Fig. 5) is approximately standard normal. We proceed with the definition of the max-min normalized rs-fMRI pairwise distance.

Consider the max-min normalized rs-fMRI distance given by the following equation

$$D_{ij}^{\text{fMRI}*} = \sum_{a \in \mathcal{A}} \sum_{k \neq a} \frac{|A_{ak}^{(i)} - A_{ak}^{(j)}|}{\max(a) - \min(a)}. \tag{163}$$

Assuming that the data  $X$  has been r-to-z transformed and standardized, we can easily compute the expected attribute range and variance of the attribute range.

The expected maximum of a given attribute in data matrix  $X$  is estimated by the following

$$\begin{aligned}
\mathbb{E}(X_a^{\max} - X_a^{\min}) &= 2\mu_{\max}^{(1)}(m, p) \\
&= 2 \left[ \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m(p-1)}\right)} - \Phi^{-1}\left(\frac{1}{m(p-1)}\right) \right].
\end{aligned} \tag{164}$$

The variance can be estimated with the following

$$\text{Var}(X_a^{\max} - X_a^{\min}) = \frac{\pi^2}{6\log[m(p-1)]}. \quad (165)$$

Let  $\mu_{D_{ij}^{\text{fMRI}}}$  and  $\sigma_{D_{ij}^{\text{fMRI}}}^2$  denote the mean and variance of the rs-fMRI distance distribution given by Eqs. 156 and 161. Using the formulas for the mean and variance of the max-min normalized distance distribution given in Eq. 89, we have the following asymptotic distribution for the max-min normalized rs-fMRI distances

$$D_{ij}^{\text{fMRI}*} \sim \mathcal{N}\left(\frac{\mu_{D_{ij}^{\text{fMRI}}}}{2\mu_{\max}^{(1)}(m,p)}, \frac{6\sigma_{D_{ij}^{\text{fMRI}}}^2 \log[m(p-1)]}{\pi^2 + 24[\mu_{\max}^{(1)}(m,p)]^2 \log[m(p-1)]}\right). \quad (166)$$

## 5.2 One-dimensional projection of rs-fMRI distance onto a single ROI

Just as in previous sections (Sections. 3.2.3 and 4.4), we now derive the distribution of our rs-fMRI diff metric (Eq. 154). Unlike what we have seen in previous sections, we do not derive the exact distribution for this diff metric. We have determined empirically that the rs-fMRI diff is approximately normal. Although the rs-fMRI diff is a sum of  $p-1$  magnitude differences, the Classical Central Limit Theorem does not apply because of the dependencies that exist between the terms of the sum. Examination of histograms and quantile-quantile plots of simulated values of the rs-fMRI diff easily indicate that the normality assumption is safe. Therefore, we derive the mean and variance of the approximately normal distribution of the rs-fMRI diff. As we have seen previously, this normality assumption is reasonable even for small values of  $p$ .

The mean of the rs-fMRI diff is derived by fixing a single ROI  $a$  and considering all pairwise associations with other ROIs  $k \neq a$ . This is done as follows

$$\begin{aligned} \mathbb{E}[d_{ij}^{\text{ROI}}(a)] &= \mathbb{E}\left(\sum_{k \neq a} |\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|\right) \\ &= \sum_{k \neq a} \mathbb{E}\left(|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|\right) \\ &= \sum_{k \neq a} \frac{2}{\sqrt{\pi}} \\ &= \frac{2(p-1)}{\sqrt{\pi}}, \end{aligned} \quad (167)$$

where  $a$  is a single fixed ROI.

Considering the variance of the rs-fMRI diff metric, we have two estimates. The first estimate uses the variance operator in a linear fashion, while the second will simply be a direct implication of the corrected formula of the variance of rs-fMRI pairwise distances (Eq. 161). Our first estimate is derived as follows

$$\begin{aligned}
 \text{Var} \left[ d_{ij}^{\text{ROI}}(a) \right] &= \text{Var} \left( \sum_{k \neq a} \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right| \right) \\
 &= \sum_{k \neq a} \text{Var} \left( \left| \hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)} \right| \right) \\
 &= \sum_{k \neq a} \frac{2(\pi - 2)}{\pi} \\
 &= \frac{2(\pi - 2)(p - 1)}{\pi},
 \end{aligned} \tag{168}$$

where  $a$  is a single fixed ROI.

Using the corrected rs-fMRI distance variance formula (Eq. 161), our second estimate of the rs-fMRI diff variance is given directly by the following

$$\text{Var} \left[ d_{ij}^{\text{ROI}}(a) \right] = \frac{9(\pi - 2)(p - 1)}{4\pi}, \tag{169}$$

where  $a$  is a single fixed ROI.

Empirically, the first estimate (Eq. 168) of the variance of our rs-fMRI diff is closer to the sample variance than the second estimate (Eq. 169). This is due to the fact that we are considering only a fixed ROI  $a \in \mathcal{A}$ , so the cross-covariance between the magnitude differences  $|\hat{A}_{ak}^{(i)} - \hat{A}_{ak}^{(j)}|$  for different pairs of ROIs ( $a$  and  $k \neq a$ ) is negligible here. When considering all ROIs  $a \in \mathcal{A}$ , these cross-covariances are no longer negligible. Using the first variance estimate (Eq. 168) and the estimate of the mean (Eq. 167), we have the following asymptotic distribution of the rs-fMRI diff

$$d_{ij}^{\text{ROI}}(a) \sim \mathcal{N} \left( \frac{2(p - 1)}{\sqrt{\pi}}, \frac{2(\pi - 2)(p - 1)}{\pi} \right), \tag{170}$$

where  $a$  is a single fixed ROI.

### 15.3 Normalized Manhattan ( $q = 1$ ) for rs-fMRI

<sup>2</sup>Substituting the non-normalized mean (Eq. 156) into the equation for the mean of<sup>2</sup>

<sup>3</sup>the max-min normalized rs-fMRI metric (Eq. 166), we have the following<sup>3</sup>

$$\begin{aligned} \text{E} \left( D_{ij}^{\text{fMRI}*} \right) &= \frac{\mu_{D_{ij}^{\text{fMRI}}}^{\text{fMRI}}}{2\mu_{\max}^{(1)}(m, p)} \\ &= \frac{p(p-1)}{\sqrt{\pi}\mu_{\max}^{(1)}(m, p)}, \end{aligned} \quad (171)$$

<sup>8</sup>where  $\mu_{\max}^{(1)}(m, p)$  (Eq. 164) is the expected maximum of a single ROI in a data set<sup>8</sup>

<sup>9</sup>with  $m$  instances and  $p$  ROIs.<sup>9</sup>

<sup>10</sup> Similarly, the variance of  $D_{ij}^{\text{fMRI}*}$  is given by<sup>10</sup>

$$\begin{aligned} \text{Var} \left( D_{ij}^{\text{fMRI}*} \right) &= \frac{6\sigma_{D_{ij}^{\text{fMRI}}}^2 \log[m(p-1)]}{\pi^2 + 24 \left[ \mu_{\max}^{(1)}(m, p) \right]^2 \log[m(p-1)]} \\ &= \frac{27(\pi-2)\log[m(p-1)](p-1)p}{2\pi \left( \pi^2 + 24 \left[ \mu_{\max}^{(1)}(m, p) \right]^2 \log[m(p-1)] \right)}, \end{aligned} \quad (172)$$

<sup>16</sup>where  $\mu_{\max}^{(1)}(m, p)$  (Eq. 164) is the expected maximum of a single ROI in a data set<sup>16</sup>

<sup>17</sup>with  $m$  instances and  $p$  ROIs.<sup>17</sup>

<sup>18</sup> We summarize the moment estimates for the rs-fMRI metrics for correlation-based<sup>18</sup>  
<sup>19</sup>data derived from time series (Table 5). We organize this summary by standard and<sup>19</sup>  
<sup>20</sup>attribute range-normalized rs-fMRI distance metric, statistic (mean or variance),<sup>20</sup>  
<sup>21</sup>and asymptotic formula.<sup>21</sup>

## 23 6 Comparison of theoretical and sample moments

<sup>24</sup>We compare our analytical asymptotic estimates of sample moments for distri-<sup>24</sup>

<sup>25</sup>butions of pairwise distances in high attribute dimension by generating random<sup>25</sup>

<sup>26</sup>data for various dimensions  $m$  and  $p$  (Fig. 6). We fix  $m = 100$  samples and<sup>26</sup>

<sup>27</sup>compute Manhattan (Eq. 1) distance matrices from standard normal data for<sup>27</sup>

<sup>28</sup> $p = 1000, 2000, 3000, 4000$ , and  $5000$  attributes. For each value of  $p$ , we generate<sup>28</sup>

<sup>29</sup>20 random datasets and compute the mean and standard deviation of pairwise dis-<sup>29</sup>


<sup>30</sup>tances. We then average these 20 simulated means and standard deviations. For<sup>30</sup>

<sup>31</sup>comparison, we compute the theoretical moments (Eqs. 38 and 39) for each value of<sup>31</sup>

<sup>32</sup> $p$  and fixed  $m = 100$  from the theoretical formulas. Scatter plots of theoretical ver-<sup>32</sup>

<sup>33</sup>sus simulated mean (Fig. 6A) and theoretical versus simulated standard deviation<sup>33</sup>

<sup>1</sup>(Fig. 6B) indicate that our theoretical asymptotic formulas for sample moments are <sup>1</sup>  
<sup>2</sup>reliable for both large and relatively small numbers of attributes. For other com- <sup>2</sup>  
<sup>3</sup>binations of data type, distance metric, sample size  $m$ , and number of attributes <sup>3</sup>  
<sup>4</sup> $p$ , we find similar agreement between theoretical formulas and simulated moments <sup>4</sup>  
<sup>5</sup>(not shown). <sup>5</sup>



## <sup>24</sup>7 Effects of correlation on distances <sup>24</sup>

<sup>25</sup>All of the derivations presented in previous sections were for the cases where there <sup>25</sup>  
<sup>26</sup>is no correlation between instances or between attributes. We assumed that any <sup>26</sup>  
<sup>27</sup>pair  $(X_{ia}, X_{ja})$  of data points for instances  $i$  and  $j$  and fixed attribute  $a$  were <sup>27</sup>  
<sup>28</sup>independent and identically distributed. This was assumed in order to determine <sup>28</sup>  
<sup>29</sup>asymptotic estimates in null data. That is, data with no main effects, interaction <sup>29</sup>  
<sup>30</sup>effects, or pairwise correlations between attributes. Within this simplified context, <sup>30</sup>  
<sup>31</sup>our asymptotic formulas for distributional moments are reliable. However, in real <sup>31</sup>  
<sup>32</sup>data are numerous statistical effects that impact distance distributional properties. <sup>32</sup>  
<sup>33</sup>We find that deviation from normality is caused primarily by large magnitude pair- <sup>33</sup>

wise correlation between attributes. Pairwise attribute correlation can be the result<sup>1</sup>  
of main effects, where attributes have different within-group means. On the other<sup>2</sup>  
hand, there could be an underlying interaction network in which there are strong<sup>3</sup>  
associations between attributes. If attributes are differentially correlated between<sup>4</sup>  
phenotype groups, then interactions exist that change the distance distribution. In<sup>5</sup>  
the following few sections, we consider particular cases of the  $L_q$  metric for contin-<sup>6</sup>  
uous and discrete data under the effects of pairwise attribute correlation.<sup>7</sup>

### 7.1 Continuous data

Without loss of generality, suppose we have  $X^{(m \times p)}$  where  $X_{ia} \sim \mathcal{N}(0, 1)$  for all<sup>10</sup>  
 $i = 1, 2, \dots, m$  and  $a = 1, 2, \dots, p$ , and let  $m = p = 100$ . We consider only the  $L_2$ <sup>11</sup>  
(Euclidean) metric (Eq. 1,  $q = 2$ ). We explore the effects of correlation on these<sup>12</sup>  
distances by generating simulated data sets with increasing strength of pairwise<sup>13</sup>  
attribute correlation and then plotting the density curve of the induced distances<sup>14</sup>  
(Fig. 7A). Deviation from normality in the distance distribution is directly related<sup>15</sup>  
to the average absolute pairwise correlation that exists in the simulated data. This<sup>16</sup>  
measure is given by<sup>17</sup>

$$\bar{r}_{\text{abs}} = \frac{2}{p(p-1)} \sum_{a \in \mathcal{A}} \sum_{k \neq a}^{p-1} r_{ak} \quad (173)$$

where  $r_{ak}$  is the correlation between attributes  $a, k \in \mathcal{A}$  across all instances  $m$ .<sup>20</sup>  
Distances generated on data without correlation closely approximate a Gaussian.<sup>21</sup>  
The mean (Eq. 50) and variance (Eq. 49) of the uncorrelated distance distribution<sup>22</sup>  
are given by substituting  $p = 100$  for the mean. As  $\bar{r}_{\text{abs}}$  increases, positive skewness<sup>23</sup>  
and increased variability in distances emerges. The predicted and sample means,<sup>24</sup>  
however, are approximately the same between correlated and uncorrelated distances<sup>25</sup>  
due to linearity of the expectation operator. Because of the dependencies between<sup>26</sup>  
attributes, the predicted variance of 1 for  $L_2$  on standard normal data obviously no<sup>27</sup>  
longer holds.<sup>28</sup>

In order to introduce a controlled level of correlation between attributes, we cre-<sup>29</sup>  
ated correlation matrices based on a random graph with specified connection prob-<sup>30</sup>  
ability, where attributes correspond to the vertices in each graph. We assigned high<sup>31</sup>  
correlations to connected attributes from the random graph and low correlations to<sup>32</sup>  
all non-connections. Using the upper-triangular Cholesky factor  $U$  for uncorrelated<sup>33</sup>



<sup>1</sup>data matrix  $X$ , we computed the following product to create correlated data matrix<sup>1</sup>

$$\begin{array}{l} \text{2 } X^{\text{corr}} \qquad \qquad \qquad \text{2} \\ \text{3} \qquad \qquad \qquad \text{3} \\ \text{4 } \quad X^{\text{corr}} = XU^T. \qquad \qquad \qquad (174) \text{4} \end{array}$$


<sup>5</sup>The new data matrix  $X^{\text{corr}}$  has approximately the same correlation structure as the<sup>5</sup>  
<sup>6</sup>randomly generated correlation matrix created from a random graph.<sup>6</sup>

## <sup>8</sup>7.2 GWAS data<sup>8</sup>

<sup>9</sup>Analogous to the previous section, we explore the effects of pairwise attribute corre-<sup>9</sup>  
<sup>10</sup>lation in the context of GWAS data. Without loss of generality, we let  $m = p = 100$ <sup>10</sup>  
<sup>11</sup>and consider only the TiTv metric (Eq. 112). To create correlated GWAS data, we<sup>11</sup>  
<sup>12</sup>first generated standard normal data with random correlation structure, just as in<sup>12</sup>  
<sup>13</sup>the previous section. We then applied the standard normal cumulative distribution<sup>13</sup>  
<sup>14</sup>function (CDF) to this correlated data in order transform the correlated standard<sup>14</sup>  
<sup>15</sup>normal variates into uniform data with preserved correlation structure. We then<sup>15</sup>  
<sup>16</sup>subsequently applied the inverse binomial CDF to the correlated uniform data with<sup>16</sup>  
<sup>17</sup>random success probabilities  $f_a$  for all  $a \in \mathcal{A}$ . Each attribute  $a \in \mathcal{A}$  corresponds<sup>17</sup>  
<sup>18</sup>to an individual SNP in the data matrix. The resulting GWAS data set is bino-<sup>18</sup>  
<sup>19</sup>mial with  $n = 2$  trials and has roughly the same correlation matrix as the original<sup>19</sup>  
<sup>20</sup>correlated standard normal data with which we started. Average absolute pairwise<sup>20</sup>  
<sup>21</sup>correlation  $\bar{r}_{\text{abs}}$  induces positive skewness in GWAS data at lower levels than in<sup>21</sup>  
<sup>22</sup>correlated standard normal data (Fig. 7B). This could have important implications<sup>22</sup>  
<sup>23</sup>in nearest neighborhoods in NPDR and similar methods.<sup>23</sup>

## <sup>25</sup>7.3 Time-series derived correlation-based datasets<sup>25</sup>

<sup>26</sup>For our correlation-based metric (Eq. 155), we consider additional effects of correla-<sup>26</sup>  
<sup>27</sup>tion between attributes. Without loss of generality, we let  $m = 100$  and  $p = 30$ . We<sup>27</sup>  
<sup>28</sup>illustrate the effects of correlated attributes in this context (Fig. 7C). Based on the<sup>28</sup>  
<sup>29</sup>density estimates, it appears that correlation between attributes introduces positive<sup>29</sup>  
<sup>30</sup>skewness at low values of  $\bar{r}_{\text{abs}}$ . We introduced correlation to the transformed data<sup>30</sup>  
<sup>31</sup>matrix (Fig. 5) with the Cholesky method discussed above.<sup>31</sup>



## 8 Discussion

Nearest-neighbor distance-based feature selection is a class of methods that are relatively simple to implement, and they perform well at detecting interaction effects in high dimensional data. Theoretical analysis of the limiting behavior of distance distributions for various data types and dimensions may lead to improved hyperparameter estimates of these feature selection methods. Furthermore, these theoretical results may help guide the choice of distance metric for a given dataset. Most often, distance-based feature selection methods use the  $L_q$  metric (Eq. 1) with  $q = 1$  or  $q = 2$ . However, these two realizations of the  $L_q$  metric have considerably different expressions for the mean and variance of their respective limiting distributions. For instance, the expected distance for  $L_1$  and  $L_2$  for standard normal data is on the order of  $p$  (Eq. 38 and Table 2) and  $\sqrt{p}$  (Eq. 48 and Table 2), respectively. In addition,  $L_1$  and  $L_2$  on standard normal data have asymptotic variances on the order of  $p$  and 1, respectively (Eqs. 39 and 49).

These results can inform the choice of  $L_1$  or  $L_2$  depending context. For instance, distances become harder to distinguish from one another in high dimensions, which is one of the curses of dimensionality. In the case of  $L_2$ , the asymptotic distribution

<sup>1</sup> $(\mathcal{N}(\sqrt{2p-1}, 1))$  indicates that the limiting  $L_2$  distribution can be thought of simply<sup>1</sup>  
<sup>2</sup>as a positive translation of the standard normal distribution  $(\mathcal{N}(0, 1))$ . The  $L_2$ <sup>2</sup>  
<sup>3</sup>distribution  $(\mathcal{N}(\sqrt{2p-1}, 1))$  also indicates that most neighbors are contained in a<sup>3</sup>  
<sup>4</sup>thin shell far from the instance in high dimension ( $p \gg 1$ ). On the other hand, the<sup>4</sup>  
<sup>5</sup> $L_1$  distances become more dispersed due to the fact that the variance of the limiting<sup>5</sup>  
<sup>6</sup>distribution is proportional to the attribute dimension  $p$ . This  $L_1$  dispersion could<sup>6</sup>  
<sup>7</sup>be more desirable when determining nearest neighbors because instances may be<sup>7</sup>  
<sup>8</sup>easier to distinguish with this metric. If using  $L_1$ , then it may be best to use a<sup>8</sup>  
<sup>9</sup>fixed-k algorithm instead of fixed-radius. This is because fixed-radius neighborhood<sup>9</sup>  
<sup>10</sup>order could vary quite a bit considering the  $L_1$  variance is proportional to attribute<sup>10</sup>  
<sup>11</sup>dimension  $p$ , which in turn could affect the quality of selected attributes. If  $L_2$  is<sup>11</sup>  
<sup>12</sup>being used, then perhaps either fixed-k or fixed-radius may perform equally well<sup>12</sup>  
<sup>13</sup>because most distances will be within 1 standard deviation away from the mean.<sup>13</sup>  
<sup>14</sup> In our analysis, we derived distance asymptotics for some of the most commonly<sup>14</sup>  
<sup>15</sup>used metrics in nearest-neighbor distance-based feature selection, as well as two new<sup>15</sup>  
<sup>16</sup>metrics for GWAS (Eq. 112) and time series correlation-based data (Eqs. 155 and<sup>16</sup>  
<sup>17</sup>163) like resting-state fMRI. We also extended the asymptotic results for the mean<sup>17</sup>  
<sup>18</sup>and variance of the attribute range-normalized  $L_q$  (max-min) distance for standard<sup>18</sup>  
<sup>19</sup>normal (Eq. 89) and standard uniform (Eq. 97) data using extreme value theory.<sup>19</sup>  
<sup>20</sup>Our derivations provide an important reference for those using nearest-neighbor<sup>20</sup>  
<sup>21</sup>feature selection or classification methods in common bioinformatics data.<sup>21</sup>  
<sup>22</sup> In this work, we expanded nearest-neighbor distance-based feature selection into<sup>22</sup>  
<sup>23</sup>the context of time series correlation-based data. Our motivation for this is partly<sup>23</sup>  
<sup>24</sup>based on the application to resting-state fMRI data. In order for this to be possible,<sup>24</sup>  
<sup>25</sup>we created a new metric (Eq. 154) that allows us to use regions of interest (ROIs)<sup>25</sup>  
<sup>26</sup>as attributes. Not all ROIs will be relevant to a particular phenotype in case-control<sup>26</sup>  
<sup>27</sup>studies, and nearest-neighbor feature selection would be a useful tool to determine<sup>27</sup>  
<sup>28</sup>important ROIs due to interactions and to help elucidate the network structure of<sup>28</sup>  
<sup>29</sup>the brain as it relates to the phenotype of interest.<sup>29</sup>  
<sup>30</sup> The recently introduced transition-transversion metric (Eq. 109) provides an addi-<sup>30</sup>  
<sup>31</sup>tional dimension to the commonly used discrete metrics in GWAS nearest-neighbor<sup>31</sup>  
<sup>32</sup>distance-based feature selection. In this work, we have provided the asymptotic<sup>32</sup>  
<sup>33</sup>mean and variance of the limiting TiTv distance distribution. This novel result,<sup>33</sup>

<sup>1</sup>as well as asymptotic estimates for the GM (Eq. 107) and AM (Eq. 108) metrics,<sup>1</sup>  
<sup>2</sup>provides an important reference to aid in neighborhood parameter selection in this<sup>2</sup>  
<sup>3</sup>context. We have also shown how the Ti/TV ratio  $\eta$  (Eq. 124) and minor allele<sup>3</sup>  
<sup>4</sup>frequency (or success probability)  $f_a$  affects these discrete distances. For the GM<sup>4</sup>  
<sup>5</sup>and AM metrics, the distance is solely determined by the minor allele frequencies<sup>5</sup>  
<sup>6</sup>because the genotype encoding is not taken into account. We showed how both mi-<sup>6</sup>  
<sup>7</sup>nor allele frequency and Ti/Tv ratio uniquely affects the TiTv distance (Figs. 4A<sup>7</sup>  
<sup>8</sup>and 4C). Because transversions are more drastic forms of mutation than transitions,<sup>8</sup>  
<sup>9</sup>this additional dimension of information is important to consider, which is why we<sup>9</sup>  
<sup>10</sup>have provided asymptotic results for this metric. 10

<sup>11</sup> In addition to asymptotic  $L_q$  distance distributions, we have also provided the<sup>11</sup>  
<sup>12</sup>exact distributions for the one-dimensional projection of the  $L_q$  distance onto indi-<sup>12</sup>  
<sup>13</sup>vidual attributes (Sections. 3.2.3, 4.4, and 5.2). These distributions are important<sup>13</sup>  
<sup>14</sup>for all nearest-neighbor distance-based feature selection algorithms, such as Relief<sup>14</sup>  
<sup>15</sup>or NPDR, because the  $L_q$  distance is a function of the one-dimensional attribute<sup>15</sup>  
<sup>16</sup>projection (diff). In particular, these projected distance distributions are impor-<sup>16</sup>  
<sup>17</sup>tant for improving inference for predictors in NPDR, which are one-dimensional<sup>17</sup>  
<sup>18</sup>attribute projections. 18

<sup>19</sup> Correlations between attributes and instances can cause significant deviations<sup>19</sup>  
<sup>20</sup>from the asymptotic results for uncorrelated data we have derived in this work. To<sup>20</sup>  
<sup>21</sup>illustrate this behavior, we showed how strong correlations lead to positive skew-<sup>21</sup>  
<sup>22</sup>ness in the distance distribution of random normal, binomial, and rs-fMRI data<sup>22</sup>  
<sup>23</sup>(Figs. 7A, 7B, and 7C). Pairwise correlation between attributes does not change<sup>23</sup>  
<sup>24</sup>the average distance, so our asymptotic results for uncorrelated data also apply<sup>24</sup>  
<sup>25</sup>when attributes are not independent. In contrast, the sample variance of distances<sup>25</sup>  
<sup>26</sup>deviates from the uncorrelated case substantially as the average absolute pairwise<sup>26</sup>  
<sup>27</sup>attribute correlation increases (Eq. 173). For fixed-radius neighborhood methods,<sup>27</sup>  
<sup>28</sup>this deviation increases the probability of including neighbors for a given instance<sup>28</sup>  
<sup>29</sup>and may reduce the power to detect interactions. The increased variability for dis-<sup>29</sup>  
<sup>30</sup>tances with correlated data may inform the choice of metric and optimization of<sup>30</sup>  
<sup>31</sup>neighborhoods in nearest-neighbor feature selection. 31

32

### Competing interests

33

The authors declare that they have no competing interests.

33

# <sup>1</sup>Author's contributions

<sup>2</sup>BAD, TTL, and BAM conceived the project. BAD carried out all mathematical derivations. All authors read and approved the final manuscript.

## Funding

<sup>4</sup>This work was supported in part by the National Institute of Health Grant No. P20GM121312 (to BAM).

## <sup>5</sup>Author details

<sup>1</sup>Department of Mathematics, University of Tulsa, 74104 Tulsa, United States. <sup>2</sup>Department of Biostatistics Epidemiology and Informatics, University of Pennsylvania, 19104 Philadelphia, United States. <sup>3</sup>Tandy School of Computer Science, University of Tulsa, 74104 Tulsa, United States.

## 8References

1. Urbanowicz, R.J., Olson, R.S., Schmitt, P., Meeker, M., Moore, J.H.: Benchmarking relief-based feature selection methods for bioinformatics data mining. *Journal of Biomedical Informatics* **85**, 168–188 (2018). doi:[10.1016/j.jbi.2018.07.015](https://doi.org/10.1016/j.jbi.2018.07.015)
2. Urbanowicz, R.J., Meeker, M., Cava, W.L., Olson, R.S., Moore, J.H.: Relief-based feature selection: Introduction and review. *Journal of Biomedical Informatics* (2018). doi:[10.1016/j.jbi.2018.07.014](https://doi.org/10.1016/j.jbi.2018.07.014)
3. Robnik-Šikonja, M., Igor Kononenko: Theoretical and Empirical Analysis of ReliefF and RReliefF. *Machine Learning* **53**, 23–69 (2003)
4. Le, T.T., Bryan A. Dawkins, Brett A. McKinney: Nearest-neighbor Projected-Distance Regression (NPDR) detects network interactions and controls for confounding and multiple testing. Under Review (2019)
5. Le, T.T., Urbanowicz, R.J., Moore, J.H., McKinney, B.A.: Statistical inference relief (stir) feature selection. *Bioinformatics*, 788 (2018). doi:[10.1093/bioinformatics/bty788](https://doi.org/10.1093/bioinformatics/bty788)
6. McKinney, B.A., White, B.C., Grill, D.E., Li, P.W., Kennedy, R.B., Poland, G.A., Oberg, A.L.: ReliefSeq: a gene-wise adaptive-K nearest-neighbor feature selection tool for finding gene-gene interactions and main effects in mRNA-Seq gene expression data. *PloS one* **8**(12), 81527 (2013)
7. Arabnejad, M., Dawkins, B.A., Bush, W.S., White, B.C., Harkness, A.R., McKinney, B.A.: Transition-transversion encoding and genetic relationship metfc in ReliefF feature selection improves pathway enrichment in GWAS. *BioData Mining* **11**(23) (2018)
8. Venkataraman, A., Marek Kubicki, Carl-Fredrik Westin, Polina Golland: Robust Feature Selection in Resting-State fMRI Connectivity Based on Population Studies. *Conf Comput Vis Pattern Recognit Workshops*, 63–70 (2010). doi:[10.1109/CVPRW.2010.5543446](https://doi.org/10.1109/CVPRW.2010.5543446)
9. Hay, E., Petra Ritter, Nancy J. Lobaugh, Anthony R. McIntosh: Multiregional integration in the brain during resting-state fMRI activity. *PLOS Computational Biology* (2017). doi:[10.1371/journal.pcbi.1005410](https://doi.org/10.1371/journal.pcbi.1005410)
10. Sundermann, B., Mona Olde lütke Beverborg, Bettina Pflöderer: Toward literature-based feature selection for diagnostic classification: a meta-analysis of resting-state fMRI in depression. *Frontiers in Human Neuroscience* (2014). doi:[10.3389/fnhum.2014.00692](https://doi.org/10.3389/fnhum.2014.00692)
11. Vergun, S., Alok S. Deshpande, Timothy B. Meier, Jie Song, Dana L. Tudorascu, Veena A. Nair, Vikas Singh, Bharat B. Biswal, Elizabeth Meverand, M., Rasmus M. Birn, Vivek Prabhakaran: Characterizing functional connectivity differences in aging adults using machine learning on resting state fMRI data. *Frontiers in Computational Neuroscience* (2013). doi:[10.3389/fncom.2013.00038](https://doi.org/10.3389/fncom.2013.00038)
12. Le, T.T., Simmons, W.K., Misaki, M., Bodurka, J., White, B.C., Savitz, J., McKinney, B.A.: Differential privacy-based evaporative cooling feature selection and classification with relief-f and random forests. *Bioinformatics* **33**(18), 2906–2913 (2017)
13. Gotts, S.J., Kyle Simmons, W., Lydia A. Milbury, Gregory L. Wallace, Robert W. Cox, Alex Martin: Fractionation of social brain circuits in autism spectrum disorders. *Brain* **135**, 2711–2725 (2012). doi:[10.1093/brain/aws160](https://doi.org/10.1093/brain/aws160)
14. Wasserman, L.: All of Statistics: A Concise Course in Statistical Inference. Springer, ??? (2004)
15. Miller, I., Marylees Miller: John E. Freund's Mathematical Statistics with Applications, 7th edn. Pearson Prentice Hall, ??? (2004)
16. Brazma, A., Jaak Vilo: Gene expression data analysis. *FEBS Letters* **480**, 17–24 (2000)

17. Wang, D., Jin Gu: VASC: Dimension Reduction and Visualization of Single-cell RNA-seq Data by Deep Variational Autoencoder. *Genomics Proteomics Bioinformatics* **16**, 320–331 (2018)
18. Gumbel, E.J.: The Distribution of the Range. *The Annals of Mathematical Statistics* **18**(3), 384–412 (1947)
19. Chatterjee, S.: *Superconcentration and Related Topics*, 1st edn. 1439–7382. Springer International Publishing, ??? (2014). doi:[10.1007/978-3-319-03886-5](https://doi.org/10.1007/978-3-319-03886-5)
20. Cramér, H.: *Mathematical Methods of Statistics* vol. 1, Reprint, revised edn. Princeton University Press, ??? (1999)
21. Li, P., Maozu Guo, Chunyu Wang, Xiaoyan Liu, Quan Zou: An overview of SNP interactions in genome-wide association studies. *Briefings in Functional Genomics* **14**(2), 143–155 (2014). doi:[10.1093/bfgp/elu036](https://doi.org/10.1093/bfgp/elu036)
22. Lee, M.H., Christopher D. Smyser, Joshua S. Shimony: Resting state fMRI: A review of methods and clinical applications. *AJNR Am J Neuroradiol.* **34**(10), 1866–1872 (2013). doi:[10.3174/ajnr.A3263](https://doi.org/10.3174/ajnr.A3263)
23. Alexander Dickie, D., Susan D. Shenkin, Devasuda Anblagan, Juyoung Lee, Manuel Blesa Cabez, David Rodriguez, James P. Boardman, Adam Waldman, Dominic E. Job, Joanna M. Wardlaw: Whole Brain Magnetic Resonance Image Atlases: A Systematic Review of Existing Atlases and Caveats for Use in Population Imaging. *Frontiers in Neuroinformatics* (2017). doi:[10.3389/fninf.2017.00001](https://doi.org/10.3389/fninf.2017.00001)

#### Additional Files

- Additional file 1 — Supplementary figures
- Contains figures regarding Gaussian convergence, theoretical versus simulated moment comparison, and density curves for one-dimensional projected distances onto a single attribute.

**Table 1** Summary of distance distribution derivations for standard normal ( $\mathcal{N}(0, 1)$ ) and standard uniform ( $\mathcal{U}(0, 1)$ ) data. Asymptotic estimates are given for both standard (Eq. 1) and max-min normalized (Eq. 55) q-metrics. These estimates are relevant for all  $q \in \mathbb{N}$  and  $p \gg 1$  for which the normality assumption of distances holds.

q-Metric	Data	Stat	Formula (Eq. #)
standard (Eq. 1)	$\mathcal{N}(0, 1)$	mean	$\left( \frac{2^q \Gamma(\frac{q+1}{2}) p}{\sqrt{\pi}} \right)^{1/q} \quad (25)$
	$\mathcal{N}(0, 1)$	variance	$\frac{4^q p}{q^2 \left( \frac{2^q \Gamma(\frac{1}{2} q + \frac{1}{2})}{\sqrt{\pi}} p \right)^{2(1-\frac{1}{q})}} \left[ \frac{\Gamma(q+\frac{1}{2})}{\sqrt{\pi}} - \frac{\Gamma^2(\frac{1}{2} q + \frac{1}{2})}{\pi} \right] \quad (26)$
	$\mathcal{U}(0, 1)$	mean	$\left( \frac{2p}{(q+2)(q+1)} \right)^{1/q} \quad (35)$
	$\mathcal{U}(0, 1)$	variance	$\frac{p}{q^2 \left( \frac{2p}{(q+2)(q+1)} \right)^{2(1-\frac{1}{q})}} \left[ \frac{1}{(q+1)(2q+1)} - \left( \frac{2}{(q+2)(q+1)} \right)^2 \right] \quad (36)$
max-min normalized (Eq. 55)	$\mathcal{N}(0, 1)$	mean	$\frac{\mu_{D_{ij}^{(q)}}}{2\mu_{\max}^{(1)}(m)} \quad (89)$ where $\mu_{D_{ij}^{(q)}}$ and $\mu_{\max}^{(1)}(m)$ are given by Eqs. 25 and 83 respectively.
	$\mathcal{N}(0, 1)$	variance	$\frac{6\log(m)\sigma_{D_{ij}^{(q)}}^2}{\pi^2 + 24[\mu_{\max}^{(1)}(m)]^2 \log(m)} \quad (89)$ where $\sigma_{D_{ij}^{(q)}}^2$ and $\mu_{\max}^{(1)}(m)$ are given by Eqs. 25 and 83 respectively.
	$\mathcal{U}(0, 1)$	mean	$\frac{(m+1)\mu_{D_{ij}^{(q)}}}{m-1} \quad (97)$ where $\mu_{D_{ij}^{(q)}}$ is given by Eq. 35
	$\mathcal{U}(0, 1)$	variance	$\frac{(m+2)(m+1)^2 \sigma_{D_{ij}^{(q)}}^2}{m^3 - m + 2} \quad (97)$ where $\sigma_{D_{ij}^{(q)}}^2$ is given by Eq. 36

**Table 2** Asymptotic estimates of means and variances for the standard  $L_1$  and  $L_2$  ( $q = 1$  and  $q = 2$  in Table 1) distance distributions. Estimates for both standard normal ( $\mathcal{N}(0, 1)$ ) and standard uniform ( $\mathcal{U}(0, 1)$ ) data are given.

<i>q</i> -Metric	Data	Stat	Formula (Eq. #)
standard $L_1$ (Eq. 1)	$\mathcal{N}(0, 1)$	mean	$\frac{2p}{\sqrt{\pi}}$ (38)
		variance	$\frac{2(\pi-2)p}{\pi}$ (39)
	$\mathcal{U}(0, 1)$	mean	$\frac{p}{3}$ (40)
		variance	$\frac{p}{18}$ (41)
standard $L_2$ (Eq. 1)	$\mathcal{N}(0, 1)$	mean	$\sqrt{2p-1}$ (48)
		variance	1 (49)
	$\mathcal{U}(0, 1)$	mean	$\sqrt{\frac{p}{6} - \frac{7}{120}}$ (51)
		variance	$\frac{7}{120}$ (52)



**Table 3** Asymptotic estimates of means and variances for the max-min normalized  $L_1$  and  $L_2$  distance distributions commonly used in Relief-based algorithms. Estimates for both standard normal  $\mathcal{N}(0, 1)$  and standard uniform  $\mathcal{U}(0, 1)$  data are given. The cumulative distribution function of the standard normal distribution is represented by  $\Phi$ . Furthermore,  $\mu_{\max}^{(1)}(m)$  (Eq. 83) is the asymptotic median of the sample maximum from  $m$  standard normal random samples.

$q$ -Metric	Data	Stat	Formula (Eq. #)
max-min normalized $L_1$ (Eq. 55)	$\mathcal{N}(0, 1)$	mean	$\frac{p}{\sqrt{\pi}\mu_{\max}^{(1)}(m)} \quad (98)$ where $\mu_{\max}^{(1)}(m) = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right)$
		variance	$\frac{12p(\pi-2)\log(m)}{\pi\left(\pi^2+24\left[\mu_{\max}^{(1)}(m)\right]^2\log(m)\right)} \quad (99)$ where $\mu_{\max}^{(1)}(m) = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right)$
	$\mathcal{U}(0, 1)$	mean	$\frac{(m+1)p}{3(m-1)} \quad (100)$
		variance	$\frac{(m+2)(m+1)^2p}{18(m^3-m+2)} \quad (100)$
max-min normalized $L_2$ (Eq. 55)	$\mathcal{N}(0, 1)$	mean	$\frac{\sqrt{2p-1}}{2\mu_{\max}^{(1)}(m)} \quad (102)$ where $\mu_{\max}^{(1)}(m) = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right)$
		variance	$\frac{6\log(m)}{\pi^2+24\left[\mu_{\max}^{(1)}(m)\right]^2\log(m)} \quad (103)$ where $\mu_{\max}^{(1)}(m) = \frac{\log(\log(2))}{\Phi^{-1}\left(\frac{1}{m}\right)} - \Phi^{-1}\left(\frac{1}{m}\right)$
	$\mathcal{U}(0, 1)$	mean	$\sqrt{\frac{p}{6} - \frac{7}{120} \left(\frac{m+1}{m-1}\right)} \quad (104)$
		variance	$\frac{7(m+2)(m+1)^2}{120(m^3-m+2)} \quad (105)$

**Table 4** Asymptotic estimates of means and variances of genotype mismatch (GM) (Eq. 110), allele mismatch (AM) (Eq. 111), and transition-transversion (TiTv) (Eq. 112) distance metrics in GWAS data ( $p \gg 1$ ). GWAS data  $X_{ia} \sim \mathcal{B}(2, f_a)$ , where  $f_a$  for all  $a \in \mathcal{A}$  are the probabilities of a minor allele occurring at locus  $a$ . For the TiTv distance metric, we have the additional encoding that uses  $\gamma_0 = P(\text{PuPu})$ ,  $\gamma_1 = P(\text{PuPy})$ , and  $\gamma_2 = P(\text{PyPy})$ .

GWAS-Metric	Stat	Formula (Eq. #)
GM (Eq. 110)	mean	$2 \sum_{a \in \mathcal{A}} F^{\text{GM}}(a) \quad (114)$ where $F^{\text{GM}}(a) = 2(1 - f_a)^3 f_a + 2f_a^3(1 - f_a) + (1 - f_a)^2 f_a^2$
	variance	$2 \sum_{a \in \mathcal{A}} F^{\text{GM}}(a)[1 - 2F^{\text{GM}}(a)] \quad (116)$ where $F^{\text{GM}}(a) = 2(1 - f_a)^3 f_a + 2f_a^3(1 - f_a) + (1 - f_a)^2 f_a^2$
AM (Eq. 111)	mean	$2 \sum_{a \in \mathcal{A}} F^{\text{AM}}(a) \quad (119)$ where $F^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a) + (1 - f_a)^2 f_a^2$
	variance	$\sum_{a \in \mathcal{A}} [G^{\text{AM}}(a) - 4(F^{\text{AM}}(a))^2] \quad (121)$ where $F^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a) + (1 - f_a)^2 f_a^2$ and $G^{\text{AM}}(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a) + 2(1 - f_a)^2 f_a^2$
TiTv (Eq. 112)	mean	$(\gamma_0 + \gamma_2 + 2\gamma_1) \sum_{a \in \mathcal{A}} F^{\text{TiTv}}(a) + \left[\frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1\right] \sum_{a \in \mathcal{A}} G^{\text{TiTv}}(a) \quad (135)$ where $F^{\text{TiTv}}(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a)$ and $G^{\text{TiTv}}(a) = (1 - f_a)^2 f_a^2$
	variance	$\left[\frac{1}{4}(\gamma_0 + \gamma_2) + \gamma_1\right] \sum_{a \in \mathcal{A}} F^{\text{TiTv}}(a) + \left[\frac{9}{8}(\gamma_0 + \gamma_2) + 2\gamma_1\right] \sum_{a \in \mathcal{A}} G^{\text{TiTv}}(a)$ $+ \sum_{a \in \mathcal{A}} \left( [\gamma_0 + \gamma_2 + 2\gamma_1] F^{\text{TiTv}}(a) + \left[\frac{3}{2}(\gamma_0 + \gamma_2) + 2\gamma_1\right] G^{\text{TiTv}}(a) \right)^2 \quad (137)$ where $F^{\text{TiTv}}(a) = (1 - f_a)^3 f_a + f_a^3(1 - f_a)$ and $G^{\text{TiTv}}(a) = (1 - f_a)^2 f_a^2$

**Table 5** Asymptotic means and variances for the new standard (Eq. 155) and max-min normalized (Eq. 163) rs-fMRI distance metrics.

rs-fMRI - Metric	Stat	Formula (Eq. #)
standard (Eq. 155)	mean	$\frac{2p(p-1)}{\sqrt{\pi}} \quad (156)$
	variance	$\frac{9p(\pi-2)(p-1)}{4\pi} \quad (158)$
max-min normalized (Eq. 163)	mean	$\frac{\mu_{D_{ij}}}{2\mu_{\max}^{(1)}(m,p)} \quad (171)$ where $\mu_{D_{ij}}$ and $\mu_{\max}^{(1)}(m,p)$ are given by Eqs. 156 and 164
	variance	$\frac{6\sigma_{D_{ij}}^2 \log[m(p-1)]}{\pi^2 + 24 \left[ \mu_{\max}^{(1)}(m,p) \right]^2 \log[m(p-1)]} \quad (172)$ where $\sigma_{D_{ij}}^2$ and $\mu_{\max}^{(1)}(m,p)$ are given by Eqs. 156 and 164