

Mathematical Statements

(Section 0.2)

Overview

- 1 Warm-up
- 2 Atomic and Molecular Statements
- 3 Implications
- 4 Predicates and Quantifiers

Warm-up

While walking through a fictional forest, you encounter three trolls guarding a bridge. Each is either a *knight*, who always tells the truth, or a *knave*, who always lies. The trolls will not let you pass until you correctly identify each as either a knight or a knave. Each troll makes a single statement: Which troll is which?

Statements: atomic and molecular

- A **statement** is any declarative sentence which is either true or false.
- A statement is **atomic** if it cannot be divided into smaller statements, otherwise it is called **molecular**.

Example (0.2.1)

These are statements (in fact, *atomic* statements):

- Telephone numbers in the USA have 10 digits.
- The moon is made of cheese.
- 42 is a perfect square.
- Every even number greater than 2 can be expressed as the sum of two primes.
- $3 + 7 = 12$

Example (0.2.1)

And these are not statements:

- Would you like some cake?
- The sum of two squares.
- $1 + 3 + 5 + 7 + \cdots + 2n + 1$.
- Go to your room!
- $3 + x = 12$

Molecular statements

You can build more complicated (molecular) statements out of simpler (atomic or molecular) ones using **logical connectives**. For example, this is a molecular statement:

Telephone numbers in the USA have 10 digits and 42 is a perfect square.

we can break this down into two smaller statements. The two shorter statements are *connected* by an “and.”

Logical Connectives

There are five logical connectives we will consider:

Logical Connectives

There are five logical connectives we will consider:

- $P \wedge Q$ is read “ P and Q ,” and called a **conjunction**.

Logical Connectives

There are five logical connectives we will consider:

- $P \wedge Q$ is read “ P and Q ,” and called a **conjunction**.
- $P \vee Q$ is read “ P or Q ,” and called a **disjunction**.

Logical Connectives

There are five logical connectives we will consider:

- $P \wedge Q$ is read “ P and Q ,” and called a **conjunction**.
- $P \vee Q$ is read “ P or Q ,” and called a **disjunction**.
- $P \rightarrow Q$ is read “if P then Q ,” and called an **implication** or **conditional**.

Logical Connectives

There are five logical connectives we will consider:

- $P \wedge Q$ is read “ P and Q ,” and called a **conjunction**.
- $P \vee Q$ is read “ P or Q ,” and called a **disjunction**.
- $P \rightarrow Q$ is read “if P then Q ,” and called an **implication** or **conditional**.
- $P \leftrightarrow Q$ is read “ P if and only if Q ,” and called a **biconditional**.

Logical Connectives

There are five logical connectives we will consider:

- $P \wedge Q$ is read “ P and Q ,” and called a **conjunction**.
- $P \vee Q$ is read “ P or Q ,” and called a **disjunction**.
- $P \rightarrow Q$ is read “if P then Q ,” and called an **implication** or **conditional**.
- $P \leftrightarrow Q$ is read “ P if and only if Q ,” and called a **biconditional**.
- $\neg P$ is read “not P ,” and called a **negation**.

Truth Conditions for Connectives

Key point: the truth-value of a molecular statement is determined by the truth values of its parts and the type of connective.

Truth Conditions for Connectives

Key point: the truth-value of a molecular statement is determined by the truth values of its parts and the type of connective.

- $P \wedge Q$ is true when both P and Q are true

Truth Conditions for Connectives

Key point: the truth-value of a molecular statement is determined by the truth values of its parts and the type of connective.

- $P \wedge Q$ is true when both P and Q are true
- $P \vee Q$ is true when P or Q or both are true.

Truth Conditions for Connectives

Key point: the truth-value of a molecular statement is determined by the truth values of its parts and the type of connective.

- $P \wedge Q$ is true when both P and Q are true
- $P \vee Q$ is true when P or Q or both are true.
- $P \rightarrow Q$ is true when P is false or Q is true or both.

Truth Conditions for Connectives

Key point: the truth-value of a molecular statement is determined by the truth values of its parts and the type of connective.

- $P \wedge Q$ is true when both P and Q are true
- $P \vee Q$ is true when P or Q or both are true.
- $P \rightarrow Q$ is true when P is false or Q is true or both.
- $P \leftrightarrow Q$ is true when P and Q are both true, or both false.

Truth Conditions for Connectives

Key point: the truth-value of a molecular statement is determined by the truth values of its parts and the type of connective.

- $P \wedge Q$ is true when both P and Q are true
- $P \vee Q$ is true when P or Q or both are true.
- $P \rightarrow Q$ is true when P is false or Q is true or both.
- $P \leftrightarrow Q$ is true when P and Q are both true, or both false.
- $\neg P$ is true when P is false.

Implications

An **implication** or **conditional** is a molecular statement of the form

$$P \rightarrow Q$$

where P and Q are statements. We say that

- P is the **hypothesis** (or **antecedent**).
- Q is the **conclusion** (or **consequent**).

An implication is *true* provided P is false or Q is true (or both), and *false* otherwise. In particular, the only way for $P \rightarrow Q$ to be false is for P to be true *and* Q to be false.

Example (0.2.2)

Consider the state-

ment: *If Bob gets a 90 on the final, then Bob will pass the class.*

Example (0.2.2)

Consider the state-

ment: *If Bob gets a 90 on the final, then Bob will pass the class.*

- This is definitely an implication: P is the statement “Bob gets a 90 on the final,” and Q is the statement “Bob will pass the class.”

Example (0.2.2)

Consider the state-

ment: *If Bob gets a 90 on the final, then Bob will pass the class.*

- This is definitely an implication: P is the statement “Bob gets a 90 on the final,” and Q is the statement “Bob will pass the class.”
- Is it true or false? What would it take for the statement to be *false*?

Example (0.2.2)

Consider the state-

ment: *If Bob gets a 90 on the final, then Bob will pass the class.*

- This is definitely an implication: P is the statement “Bob gets a 90 on the final,” and Q is the statement “Bob will pass the class.”
- Is it true or false? What would it take for the statement to be *false*?
- The only way to be false: Bob does get a 90 on the final *AND* Bob still does not pass the class.

Example (0.2.2)

Consider the state-

If Bob gets a 90 on the final, then Bob will pass the class.

ment:

- This is definitely an implication: P is the statement “Bob gets a 90 on the final,” and Q is the statement “Bob will pass the class.”
- Is it true or false? What would it take for the statement to be *false*?
- The only way to be false: Bob does get a 90 on the final *AND* Bob still does not pass the class.
- In particular, if Bob does not get a 90 on the final (P is false), then whether or not he passes the class, the statement is true.

Example (0.2.3)

Decide which of the following statements are true and which are false. Briefly explain.

- 1 If $1 = 1$, then most horses have 4 legs.
- 2 If $0 = 1$, then $1 = 1$.
- 3 If 8 is a prime number, then the 7624th digit of π is an 8.
- 4 If the 7624th digit of π is an 8, then $2 + 2 = 4$.

Direct Proofs of Implications

The truth conditions for an implication tell us one way we could *prove* that an implication is true.

Direct Proofs of Implications

The truth conditions for an implication tell us one way we could *prove* that an implication is true. To prove an implication $P \rightarrow Q$, it is enough to *assume* P , and from it, *deduce* Q .

Example (0.2.4)

Prove: If two numbers a and b are even, then their sum $a + b$ is even.

Proof:

Example (0.2.4)

Prove: If two numbers a and b are even, then their sum $a + b$ is even.

Proof: Suppose the numbers a and b are even. This means that $a = 2k$ and $b = 2j$ for some integers k and j . The sum is then $a + b = 2k + 2j = 2(k + j)$. Since $k + j$ is an integer, this means that $a + b$ is even. \square

Converse and Contrapositive

How does $P \rightarrow Q$ relate to $Q \rightarrow P$?

Converse and Contrapositive

How does $P \rightarrow Q$ relate to $Q \rightarrow P$?

- The **converse** of an implication $P \rightarrow Q$ is the implication $Q \rightarrow P$.

Converse and Contrapositive

How does $P \rightarrow Q$ relate to $Q \rightarrow P$?

- The **converse** of an implication $P \rightarrow Q$ is the implication $Q \rightarrow P$.
- The converse is *NOT* logically equivalent to the original implication.
That is, whether the converse of an implication is true is independent of the truth of the implication.

Converse and Contrapositive

How does $P \rightarrow Q$ relate to $Q \rightarrow P$?

- The **converse** of an implication $P \rightarrow Q$ is the implication $Q \rightarrow P$.
- The converse is *NOT* logically equivalent to the original implication.
That is, whether the converse of an implication is true is independent of the truth of the implication.
- The **contrapositive** of an implication $P \rightarrow Q$ is the statement $\neg Q \rightarrow \neg P$.

Converse and Contrapositive

How does $P \rightarrow Q$ relate to $Q \rightarrow P$?

- The **converse** of an implication $P \rightarrow Q$ is the implication $Q \rightarrow P$.
- The converse is *NOT* logically equivalent to the original implication. That is, whether the converse of an implication is true is independent of the truth of the implication.
- The **contrapositive** of an implication $P \rightarrow Q$ is the statement $\neg Q \rightarrow \neg P$.
- An implication and its contrapositive are logically equivalent (they are either both true or both false).

Example (0.2.5)

True or false: If you draw any nine playing cards from a regular deck, then you will have at least three cards all of the same suit.

Example (0.2.5)

True or false: If you draw any nine playing cards from a regular deck, then you will have at least three cards all of the same suit. Try looking at the contrapositive!

Example (0.2.5)

True or false: If you draw any nine playing cards from a regular deck, then you will have at least three cards all of the same suit. Try looking at the contrapositive! Is the converse true?

Example (0.2.6)

Suppose I tell Sue that if she gets a 93% on her final, then she will get an A in the class. Assuming that what I said is true, what can you conclude in the following cases:

- 1 Sue gets a 93% on her final.
- 2 Sue gets an A in the class.
- 3 Sue does not get a 93% on her final.
- 4 Sue does not get an A in the class.

If and only if

$P \leftrightarrow Q$ is logically equivalent to $(P \rightarrow Q) \wedge (Q \rightarrow P)$.

If and only if

$P \leftrightarrow Q$ is logically equivalent to $(P \rightarrow Q) \wedge (Q \rightarrow P)$.

Example: Given an integer n , it is true that n is even if and only if n^2 is even. That is, if n is even, then n^2 is even, as well as the converse: if n^2 is even, then n is even.

Example (0.2.7)

Suppose it is true that I sing if and only if I'm in the shower. We know this means both that if I sing, then I'm in the shower, and also the converse, that if I'm in the shower, then I sing. Let P be the statement, "I sing," and Q be, "I'm in the shower." So $P \rightarrow Q$ is the statement "if I sing, then I'm in the shower." Which part of the if and only if statement is this?

Example (0.2.8)

Rephrase the implication, “if I dream, then I am asleep” in as many different ways as possible. Then do the same for the converse.

Necessary and Sufficient

A common way to express the relationship between statements in mathematics is to say that one is necessary or sufficient for the other.

Necessary and Sufficient

A common way to express the relationship between statements in mathematics is to say that one is necessary or sufficient for the other.

- “ P is necessary for Q ” means $Q \rightarrow P$.

Necessary and Sufficient

A common way to express the relationship between statements in mathematics is to say that one is necessary or sufficient for the other.

- “ P is necessary for Q ” means $Q \rightarrow P$.
- “ P is sufficient for Q ” means $P \rightarrow Q$.

Necessary and Sufficient

A common way to express the relationship between statements in mathematics is to say that one is necessary or sufficient for the other.

- “ P is necessary for Q ” means $Q \rightarrow P$.
- “ P is sufficient for Q ” means $P \rightarrow Q$.
- If P is necessary and sufficient for Q , then $P \leftrightarrow Q$.

Example (0.2.9)

Recall from calculus, if a function is differentiable at a point c , then it is continuous at c , but that the converse of this statement is not true (for example, $f(x) = |x|$ at the point 0). Restate this fact using “necessary and sufficient” language.

Predicates

How could we claim that if n is prime, then $n + 7$ is not prime? This looks like an implication. I would like to write something like

$$P(n) \rightarrow \neg P(n + 7)$$

where $P(n)$ means “ n is prime.” But this is not quite right.

Predicates

How could we claim that if n is prime, then $n + 7$ is not prime? This looks like an implication. I would like to write something like

$$P(n) \rightarrow \neg P(n + 7)$$

where $P(n)$ means “ n is prime.” But this is not quite right. For one thing, this sentence has a **free variable** (that is, a variable that we have not specified anything about), so it is not a statement. A sentence that contains variables is called a **predicate**.

Predicates

How could we claim that if n is prime, then $n + 7$ is not prime? This looks like an implication. I would like to write something like

$$P(n) \rightarrow \neg P(n + 7)$$

where $P(n)$ means “ n is prime.” But this is not quite right. For one thing, this sentence has a **free variable** (that is, a variable that we have not specified anything about), so it is not a statement. A sentence that contains variables is called a **predicate**. If we plug in a specific value for n , we do get a statement. What we really want to say is that *for all* values of n , if n is prime, then $n + 7$ is not. We need to *quantify* the variable.

Universal and Existential Quantifiers

The existential quantifier is \exists and is read “there exists” or “there is.” For example,

$$\exists x(x < 0)$$

asserts that there is a number less than 0.

Universal and Existential Quantifiers

The existential quantifier is \exists and is read “there exists” or “there is.” For example,

$$\exists x(x < 0)$$

asserts that there is a number less than 0. The universal quantifier is \forall and is read “for all” or “every.” For example,

$$\forall x(x \geq 0)$$

asserts that every number is greater than or equal to 0.

Quantifiers and Negation

When is a quantified statement false? $\neg\forall xP(x)$ is equivalent to $\exists x\neg P(x)$.
 $\neg\exists xP(x)$ is equivalent to $\forall x\neg P(x)$.